

T2K: Status and Prospects

Ali Ajmi
(on behalf of **T2K Collaboration**)

Kyoto University



*Exploration of Particle Physics and Cosmology with
Neutrinos Workshop 2019*




14 June, 2019





The T2K Experiment:



The T2K Experiment:



ν -Physics @ Far and Near Detectors:




Phys. Rev. D 88, 032002

Far Detector:

ν -Oscillation parameters

Near Detectors: Pre-Osc. Flux-measurement, ν -Interaction/Cross-section measurement

$$\frac{N^{\text{far}}}{N^{\text{near}}} \sim f(\sigma, \Phi, Det)$$



Measuring neutrino Cross Sections

$$N(\vec{x}) = \Phi(E_\nu) \cdot \sigma(E_\nu, \vec{x}) \cdot \epsilon(E_\nu) \cdot P_{ab}(E_\nu, \Delta m_{ij}^2, \theta_{ij})$$

- Systematic uncertainty important for precision in oscillation results.

- **Events** ratios at **Near** and **Far** detectors do not cancel systematics: owing to different

Flux, **Cross-sections**, **Detector-smearing**, and **Osc. Probability**.

- $\sigma(E_\nu, \vec{x})$ relate E_ν with observables



✠ Measurement of σ with minimal uncertainty

⇒ Reduced Systematics in Osc. analysis @FD

⇒ Better goals-designing of future OA experiments





⇒ Several measurements needed to constrain and develop ν -interaction models



Flux-U.Details @poster: Lukas





The Cross-section Strategy:

- Nuclear & Detector effects obscure true interaction mode
- Sample w.r.t visible outgoing hadron (π) content






- Sample w.r.t. to #reconstructed tracks
- Measure in: Reconstructed muon momentum and direction.
- Multi-targets: CH and CH+H2O separately, Fe, H2O
- Multi-beam energy peaks: Off-axis: $2.5^\circ \sim 0.6$ GeV; On-axis: $0^\circ \sim 1$ GeV
- Phase space coverage: fwd, bwd, 4π .





Sample Selection:

- ND280 flux favors ν -CCQE as a major fraction
- Appropriate selections to ensure high purity of an int.mode. Altered criteria enrich other modes
- Signal selection efficiency and purity ensuring crucial.
- Primarily: Event quality, Total multiplicity, Quality and fiducial origin cut, μ -CID multiplicity check etc.
- Other cuts applied depending on the considered target.



As for example in the following sample in FGD1:

- Vetoing with TPC and FGD2 applied
- Additional constraints from the ECAL improve the selection, by reducing the p-bkg.
- Max. Signal efficiency/purity in the p_μ and $\cos\theta_\mu$ ensured



Unfolding the Reconstructed Events:

- An unfolding method used aiming to remove the detector effects in the measurement
- The flux-avg./int. xsec. is calculated from the **unfolded** events N in each bin, $N_{\alpha\beta}$

$$\text{For e.g. , } \left\langle \frac{\delta^2 \sigma}{\delta p_\mu \delta \cos \theta_\mu} \right\rangle_{\alpha\beta} = \frac{\hat{N}_{\alpha\beta}}{T \Phi \Delta p_{\mu,\alpha} \Delta \cos \theta_{\mu,\beta}}$$



And finally also obtain total cross section σ .

- So choice of the **Unfolding method** to be used must be checked to ensure min. MC bias, and convergence of the unfolding process.
- **Unfolding Techniques:** D'Agostini, SVD, TUnfold or binned-likelihood fit and many more available.
- Checks with lots of pseudo data sets. After lot of checks and reviews: unblind the data.
- Forward-folding techniques also used by some.

Cross Section Results from T2K:





CC Inclusive on CH (Off-axis):

- Fwd, Bwd, & High angle selections
- CCQE dominance, Flux-integrated Cross Section ($\sim 6.9 \pm 0.7\%$ stat. $\pm 8.7\%$ fluxU. $\pm 1.8\%$ syst. $\times 10^{-39} \text{cm}^2 \text{nucleon}^{-1}$)
- Binned likelihood fit, unregularised. Fit with both NEUT and GENIE as priors.




CC 0π on CH, H₂O (Off-axis):

- CC $0\pi \sim 80\%$ CCQE + 12% 2p2h
- Measurements in P0D (2018) (1-track), Compared with FGD1(2016)
- Unfolded with D'Agostini method, MC truth used as prior
- Flux-int. $\sigma_{\nu\mu}^{CC0\pi} = 0.95 \pm 8\%$ stat. $\pm 8\%$ fluxU. $\pm 10\%$ syst. $\times 10^{-38} \text{cm}^2 \text{nuc}^{-1}$
- Compared with other models. Better if 2p2h included.




Many More Cross Section results:

- ☑ CC $0\pi+p$ on CH (Off-axis, phase space constraints boosting the efficiency; Standard Transverse variables introduced for probing p): PRD 98, 032003 (2018)
- ☑ CC Inclusive on CH, Fe, H2O (On-axis): Paper under prep.
- ☑ CC- $1\pi^+$ on CH, H2O (Off-axis): PRD 95 (2017) 012010
- ☑ CC 0π for $\bar{\nu}_\mu$ on H2O (Off-axis): Paper under prep.
- ☐ NC $1\pi^0$ and CC $1\pi^0$ (rate measured: PRD 97 032002 (2018), cross section ongoing).
- ☐ $\nu_e/\bar{\nu}_e$ in the FHC/RHC mode: coming soon.
- ☐ Several more analyses ongoing: $\bar{\nu}_\mu$ CC-Inc., NC 1γ etc. etc.



Newer Near Detectors: (Off-Axis)



- The **WAGASCI** detector at 1.5° Off-Axis taking data since 2017
- Cross Section measured with $\bar{\nu}_\mu$ on H₂O, CH.

$$\sigma_{\text{H}_2\text{O}}^{\bar{\nu}_\mu} = [1.082 \pm 0.068(\text{stat.})_{-0.128}^{+0.145}(\text{syst.})] \times 10^{-39} \text{cm}^2$$


$$\sigma_{\text{CH}}^{\bar{\nu}_\mu} = [1.096 \pm 0.054(\text{stat.})_{-0.117}^{+0.132}(\text{syst.})] \times 10^{-39} \text{cm}^2$$

$$\sigma_{\text{H}_2\text{O}}^{\bar{\nu}_\mu} / \sigma_{\text{CH}}^{\bar{\nu}_\mu} = 0.987 \pm 0.078(\text{stat.})_{-0.090}^{+0.093}(\text{syst.})$$


$$\sigma_{\text{H}_2\text{O}}^{\bar{\nu}_\mu + \nu_\mu} = [1.155 \pm 0.064(\text{stat.})_{-0.129}^{+0.148}(\text{syst.})] \times 10^{-39} \text{cm}^2 \cdot \text{nucleon}^{-1}$$

$$\sigma_{\text{CH}}^{\bar{\nu}_\mu + \nu_\mu} = [1.159 \pm 0.049(\text{stat.})_{-0.115}^{+0.129}(\text{syst.})] \times 10^{-39} \text{cm}^2 \cdot \text{nucleon}^{-1}$$

$$\sigma_{\text{H}_2\text{O}}^{\bar{\nu}_\mu + \nu_\mu} / \sigma_{\text{CH}}^{\bar{\nu}_\mu + \nu_\mu} = 0.996 \pm 0.069(\text{stat.})_{-0.078}^{+0.083}(\text{syst.})$$




Oscillation Studies at the FAR detector: T2K-SK



The Oscillation Measurements: Super-Kamiokande

(General details already covered in Roger's talk)




Phys. Rev. D 96, 011102

- ν_μ Disappearance: Precision measurement of θ_{23} and Δm_{32}^2
- $\nu_e/\bar{\nu}_e$ Appearance: δ_{CP}

Octant degeneracy? CP violation?

Order of Mass Hierarchy ?

The Analysis Strategy (Conceptual):




Analyses & Results:


- Improved reconstruction & classifier
- Expanded Fiducial Volume
- Select events with no pions (& $\nu_e 1\pi^+$)

$$E_{\nu}^{QE} = \frac{m_p^2 - m_n^2 - m_{\mu}^2 + 2m_n E_{\mu}}{2(m_n - E_{\mu} + p_{\mu} \cos \theta_{\mu})}$$

- Fit is done simultaneously to all five data samples





Sample	Predicted				Observed	Systematic uncertainty for prediction
	$\delta_{CP} = -\pi/2$	$\delta_{CP} = 0$	$\delta_{CP} = +\pi/2$	$\delta_{CP} = \pi$		
ν mode μ -like	272.4	272.0	272.4	272.8	243	5.1%
$\bar{\nu}$ mode μ -like	139.5	139.2	139.5	139.9	140	4.5%
ν mode e-like	74.4	62.2	50.6	62.7	75	8.8%
$\bar{\nu}$ mode e-like	17.1	19.4	21.7	19.3	15	7.1%
ν mode e-like + $1\pi^+$	7.0	6.1	4.9	5.9	15	18.4%





Results from T2K-SK data:

- Favoring $\delta_{CP} \sim -\pi/2$, 2σ confidence interval
- Including reactor constrain, matches PDG2018 value: $\sin^2 \theta_{13} = 0.0212$



- Maximal mixing favored



Status Summary and Prospects:


- ✦ T2K has added more new $\bar{\nu}$ -data in the analyses.
- ✦ Oscillation Results updated; 2σ in deciding CP-violation
- ✦ T2K-near detectors: on-axis, off-axis: Cross sections measured for several samples, several still on-going.
- ✦ T2K near detectors (ND280/WAGASCI/BABYMIND) and beamline upgrade to allow 750 kW–1.3 MW: to set for obtaining 3σ in deciding CP-violation.

Status Summary and Prospects: (contd...)


- ⊠ New off-axis WAGASCI already started
- ⊠ BABYMIND being integrated, to add charge id-capability to WAGASCI
- ⊠ Ongoing efforts to install the ND280-Upgrade very soon



Details talked at
Kenji/Giorgio posters



More@posters:
Soichiro,
Takuji, Yuki



Thank you !

Back ups:

We define for every bin k :

$$\hat{N}_k = \frac{\sum_j^{n_r} U_{jk} (N_j^{sel} - B_j)}{\epsilon_k}. \quad (1)$$

- the selected events in reconstructed bin $j = 1, \dots, n_r$: N_j^{sel} .
- the background events in reconstructed bin j : B_j .
- the generated events in true bin $k = 1, \dots, n_t$: N_k


The smearing matrix S_{jk} = the number of events generated in the k -th bin and reconstructed in the j -th bin.

- the probability to have an event reconstructed in the bin j when it has been generated in the true bin k : $P(j|k) = \frac{S_{jk}}{N_k}$
- the probability that an event was generated in the truth bin k given that it was reconstructed in bin j : U_{jk} (unfolding matrix)

$$U_{jk} = \frac{\frac{S_{jk}}{N_k} \frac{N_k}{\sum_k N_k}}{\sum_{\alpha}^{n_t} \left(\frac{S_{j\alpha}}{N_{\alpha}} \frac{N_{\alpha}}{\sum_{\alpha} N_{\alpha}} \right)} = \frac{S_{jk}}{\sum_{\alpha}^{n_t} S_{j\alpha}} \quad (2)$$

$$\epsilon_k = \frac{\sum_j^{n_r} S_{jk}}{N_k} \quad (3)$$

Effect of ND280-fit on SK-analyses:



Sample	w/o ND280	w/ND280
FHC $1R_\mu$	14.6%	5.1%
RHC $1R_\mu$	12.5%	4.5%
FHC $1R_e$	16.9%	8.8%
RHC $1R_e$	14.4%	7.1%

The Oscillation Probability:

Appearance

$$P(\nu_\mu \rightarrow \nu_e) = 4c_{13}^2 s_{13}^2 s_{23}^2 \sin^2 \Delta_{31} \times \left(1 \pm \frac{2a}{\Delta m_{31}^2} (1 - s_{13}^2) \right)$$

Leading term

$$+ 8c_{13}^2 s_{12} s_{13} s_{23} (c_{12} c_{23} \cos \delta - s_{12} s_{13} s_{23}) \cos \Delta_{32} \sin \Delta_{31} \sin \Delta_{21}$$

CP Conserving

ν vs. $\bar{\nu}$
sign
change

$$\mp 8c_{13}^2 s_{13}^2 s_{23}^2 \cos \Delta_{32} \sin \Delta_{31} \frac{aL}{4E} (1 - 2s_{13}^2)$$

Matter effect

$$\mp 8c_{13}^2 c_{12}^2 c_{23} s_{12} s_{13} s_{23} \sin \delta \sin \Delta_{32} \sin \Delta_{31} \sin \Delta_{21}$$

CP Violating

$$+ 4s_{12}^2 c_{13}^2 (c_{12}^2 c_{23}^2 + s_{12}^2 s_{13}^2 s_{23}^2 - 2c_{12} c_{23} s_{12} s_{13} s_{23} \cos \delta) \sin^2 \Delta_{21}$$

Solar term

$$c_{ij} = \cos \theta_{ij}, \quad s_{ij} = \sin \theta_{ij}, \quad \Delta_{ij} = \Delta m_{ij}^2 \frac{L}{4E_\nu}, \quad a = 2\sqrt{2} G_F n_e E$$

Disappearance

$$P(\nu_\mu \rightarrow \nu_\mu) \approx 1 - \left(\cos^4 \theta_{13} \sin^2 2\theta_{23} + \sin^2 2\theta_{13} \sin^2 \theta_{23} \right) \sin^2 \frac{\Delta m_{32}^2 \cdot L}{4E_\nu}$$