

Neutrinoless Double-Beta Decay and Xe Gas TPC Development

S. Obara

for the AXEL Collaboration




京都大学
KYOTO UNIVERSITY

Outline

- INTRODUCTION;
 - Double-Beta Decay and Majorana Neutrinos
 - Recent $0\nu 2\beta$ Search Experiments
 - Importance for Future Detectors
- AXEL EXPERIMENT;
 - Conceptual Design of AXEL
 - Results from Xe Gas 10L size prototype (HP10L)
 - Overview of Recent Study in Xe Gas 180L size prototype (HP180L)
- SUMMARY

Neutrino Mysteries




We know $m_\nu \neq 0$

via ν -oscillation

However, we still have some neutrino mysteries...

- Mass Hierarchy ?
- Majorana ? Dirac ?
- Why so small mass ?

and also CP-phase, sterile, absolute mass, ...



Neutrino is Dirac ? Majorana ?




Dirac

$$\nu \neq \bar{\nu}$$

Majorana

$$\nu = \bar{\nu}$$



$$\mathcal{L}_{D+M} = m_D (\bar{\psi}_L \psi_R + \bar{\psi}_L^c \psi_R^c) + \underbrace{m_L \bar{\psi}_L^c \psi_L^c + m_R \bar{\psi}_R^c \psi_R^c}_{\text{Majorana mass term}} + h.c.$$

Majorana mass term


In Majorana case,,,

See-Saw mechanism^[1-3] explain why so small mass

$$m_\nu \sim \frac{m_D^2}{m_R}$$

$$\mathcal{L}_{D+M} = (\bar{\nu} \quad \bar{N}) \begin{pmatrix} m_\nu & 0 \\ 0 & m_N \end{pmatrix} \begin{pmatrix} \nu \\ N \end{pmatrix} \longrightarrow (\bar{\nu} \quad \bar{N}) \begin{pmatrix} \frac{m_D^2}{m_R} & 0 \\ 0 & m_R \end{pmatrix} \begin{pmatrix} \nu \\ N \end{pmatrix}$$


- [1] Yanagida 1979,
- [2] Gell-Mann, Ramond and Slansky 1979,
- [3] Minkowski 1977



What's the double-beta decay ?


Neutrinoless Double-Beta Decay is ...

- Two beta-rays only emission
- Monochromatic energy @ Q-value
- Evidence of Majorana Neutrino



For some nuclei, single-beta decay is prohibited by the energy level but simultaneously double-beta is allowed


Energy spectra for $\beta\beta$ decay



Nucl	Q [keV]	Nat. [%]	Cost* [M\$/ton]	$T^{2\nu_{1/2}}$ [yr]	Merit 😊	Demerit 😓	Experiments
^{48}Ca	4271	0.187	> 1000	4.4×10^{19}	Largest Q-value	Low NA, Difficult enrich	CANDLES
^{76}Ge	2040	7.7	~80	1.6×10^{21}	Long T2v		GERDA, MAJORANA, (LEGEND)
^{82}Se	2995	9.2	~120	9.6×10^{19}			NEMO-3, (Super-NEMO)
^{96}Zr	3350	2.8		2.4×10^{19}	Large Q-value		(ZICOS)
^{100}Mo	3034	9.6	~80	7.1×10^{18}	Large Q-value	Short T2v	AMoRE, NEMO-3
^{116}Cd	2802	7.5	~180	2.8×10^{19}			COBRA
^{130}Te	2533	34.5	20	7.0×10^{20}	High NA		CUORE, (SNO+), (CUPID)
^{136}Xe	2457	8.9	5~10	2.3×10^{21}	Long T2v, Easy enrich		KamLAND-Zen, PANDAX-III, EXO, NEXT, AXEL, and etc...
^{150}Nd	3367	5.6	> 300	9.1×10^{18}	Large Q-value	ShortT2v, Difficult enrich	DCBA, NEMO

Recent $0\nu 2\beta$ search


Searching the monochromatic $\beta\beta$ signal...



$$\left(T_{1/2}^{0\nu}\right)^{-1} = G^{0\nu} |M^{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$

[PhaseSpace] [NME] [EffectiveMass]

$$\langle m_{\beta\beta} \rangle = \left| |U_{e1}^L|^2 m_1 + |U_{e2}^L|^2 m_2 e^{i\phi_2} + |U_{e3}^L|^2 m_3 e^{i\phi_3} \right|$$



KamLAND-Zen (Liq.Scintillator)

$T_{1/2} > 1.07 \times 10^{26}$ yr (90%C.L.)

$|m_{\beta\beta}| < 61-165$ meV (90%C.L.)

Thanks to their

large volume and **low radioactivity**


GERDA (Germanium)

$T_{1/2} > 8.0 \times 10^{25}$ yr (90%C.L.)


$|m_{\beta\beta}| < 120-260$ meV (90%C.L.)

Thanks to their

good energy resolution and **background-free in ROI**




Neutrino Hierarchy is Normal ? or Inverted ?




Recent $0\nu 2\beta$

We have to explore 0.01 ~ 0.001 meV region !!

Search



$$\left(T_{1/2}^{0\nu}\right)^{-1} = G^{0\nu} |M^{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$



KamLAND-Zen (Liq.Scintillator)

$T_{1/2} > 1.07 \times 10^{26}$ yr (90% C.L.)
 $|m_{\beta\beta}| < 61-165$ meV (90% C.L.)

Thanks to their

large volume and **low radioactivity**

GERDA (Germanium)

$T_{1/2} > 8.0 \times 10^{25}$ yr (90% C.L.)
 $|m_{\beta\beta}| < 120-260$ meV (90% C.L.)

Thanks to their

high energy resolution and **background-free in ROI**

for Future $0\nu 2\beta$ Detectors

Gaseous Xenon TPC has three advantages !!

High-Pressure Gas



- 90% enrich ^{136}Xe
- Smaller than Super-K even if 1-ton Xe

Xenon Gas

- Small W-value
- Electroluminescence process


Tracking

- Identify two-electrons



AXEL

(A Xenon ElectroLuminescence detector)



**Cockcroft-Walton
Voltage multiplier**
for HV supply

ELCC
for energy & tracking


- ^{136}Xe for $0\nu 2\beta$
- 1ton xenon gas
- 6-8 bar

VUV PMTs
for T_0 signal

Drift Field Cage

AXEL


(A Xenon ElectroLuminescence detector)



Pure Xenon Gas TPC


- $^{136}\text{Xe} = 0\nu 2\beta$ nuclei
- Ionization $W = 21\text{eV}$
- X Easy to discharge

Energy resolution of the Xenon Ion chamber
A. Bolotnikov, B. Ramsey Nucl. Instr. And Meth. A396(1997) 360




AXEL

(A Xenon ElectroLuminescence detector)




AXEL

(A Xenon ElectroLuminescence detector)



Use **ElectroLuminescence Process** for the ionization electrons tracking





- Linear Amplification
- Good Energy Resolution

cf1; NEXT also uses EL-process but with mesh plane
cf2; PandaX-III uses MM for tracking readout

ELCC

(Electroluminescence Light Collection Cell)

Cell-Type Tracking Plane








- Collect the drift electrons into cells
- Linear Amp. via EL-process
- Detect EL-photons by MPPCs

ELCC

(Electroluminescence Light Collection Cell)

- Optimize Parameter; hole-size, hole-pitch
- Max; Electron collection efficiency with ΔE
- Use; Finite Element Method
- Gmsh, ElmerSolver, and Garfield++
- => $\Phi 5.5\text{mm}$, 10mm-pitch



AXEL prototypes & Future prospects

2015



2018

202?

20??

10L - size (10g xenon)

- Establish ELCC
- Basic R&Ds
- Check ΔE




3000L - size (100 kg xenon)

- Underground




Large - size

180L - size (8.6kg xenon)

- Evaluate ΔE @ Q-value
- Estimate Radioactivity, Backgrounds
- Now constructing ...




10L-prototype





10L-prototype

- Using ^{57}Co , ^{133}Ba source for ΔE evaluation
- Q-value (2.46MeV) β -ray is not fully contained in this small size




- Estimate with extrapolation; 0.8~1.8% FWHM @ Q-value

- ※ This worse ΔE is caused by ELCC design
- ※ Not so 100% efficiency for EL-photon collection
- ※ Next HP180L prototype has the fixed ELCC






180L prototype detector






10-times larger size



- Evaluate Energy Resolution at Q-value
- Establish the Scalability for larger detector
- Now Constructing ...
(Fin. 3unit/27unit=10%)



ELCC-Unit, FieldCage





Firstly, we start with
3-Units



First Light in HP180L !!!!!!!!!!!!!!!!

An environmental- γ event in a ELCC-Unit (56ch)




- Xe 1atm
- Field Region $\Phi 15\text{cm} \times 10\text{cm}$
- No source (Environmental- γ)
- EL region; 3.0 kV/cm/atm
- Drift region; 100V/cm/atm

Now on detail analysis with source-run

Electronics board & MPPC gain correction



New boards for HP180L



- Waveform readout from MPPCs; 56ch/board
- Dynamics range of Flash ADC; 1-10⁴p.e./μsec (~2.5MeV)
- Monitor & Adjust the gain with feedback logic
- HV-supply to each MPPC (in the 0.2mV)

Talk by Nakamura

MPPC gain correction




- Non linearity after large light intensity
- We need the correction factor of saturation curve
- Measure with LED

FieldCage and Cockcroft-Walton Circuit for Drift Electric Field Supply

Develop "non-discharge field cage"

- Pure Xenon gas is easy to discharge
- Block the electric path



Talk by Yoshida



Develop "CW circuit for E_{dfirt} "

- Prevent to discharge at feedthrough
- HV is generated in the Chamber with Cockcroft-Walton Circuit

Prototype CW circuit



BG for $0\nu 2\beta$

From the Chamber material

Even if we achieve good $\Delta E = 0.5\%$ -FWHM,


^{214}Bi γ -ray 2,447keV

is dominant BG for **^{136}Xe - $0\nu 2\beta$ (2,458keV)**

Topology identification !!

By DeepLearning; 75evt/yr \Rightarrow 0.8evt/yr

arXiv:1106.3630v1 [physics.ins-det] from NEXT paper




Positive Ion maybe useful for topology used BG rejection

Talk by Nakamura

$0\nu 2\beta$ (double- β)



^{214}Bi (gamma-ray)



Very Recent Status (1-good, 3-bad)

😄 First Light Observation & Well-working New Components

- Successfully worked !!
- New ELCC, New ReadoutFPC, New Electronics, ...




😓 Discharge ; ELCC-anode to GND-mesh

- Now investigating
- Find the electric path of discharge

😓 Dead Circulation Pump !!

- Not working, now being repaired



Anode plate to GND mesh through between the Units ?

😓 Aeration by Booster Pump Trouble



- Another pump (booster pump) has also trouble and Aeration !!
- How to purify our Xenon ?



Very Recent Status

😄 First Light Observation & Well-working New Components

- Successfully worked !!
- New ELCC, New ReadoutFPC, New Electronics, ...




Very Recent Status

😓 Discharge ; New ELCC-anode to GND-mesh


- Now investigating
- Find the electric path of discharge

Talk by Yoshida





Anode plate to GND mesh
through between the Units ?

Very Recent Status



😓 **Dead Circulation Pump !!**




- Not working, now being repaired

Very Recent Status

😓 **Aeration** by Booster Pump Trouble



- Another pump (booster pump) has also trouble and Aeration !!
- No any scintillation light nor EL light
- Found High H₂O contamination by dew-point meter
- **How to purify our Xenon ?**



Very Recent Status (1-good, 3-bad)

😄 First Light Observation & Well-working New Components

- Successfully worked !!
- New ELCC, New ReadoutFPC, New Electronics, ...




😓 Discharge ; ELCC-anode to GND-mesh

- Now investigating
- Find the electric path of discharge

😓 Dead Circulation Pump !!

- Not working, now being repaired



Anode plate to GND mesh through between the Units ?

😓 Aeration by Booster Pump Trouble

- Another pump (booster pump) has also trouble and Aeration !!
- How to purify our Xenon ?



SUMMARY

- $0\nu 2\beta$ search is important to reveal the **Majorana nature of Neutrinos**
- For future double-beta decay search experiments (towards NH),
 - Large Target Mass, High Energy Resolution, Topology Tracking
 - Xe Gas TPC satisfy above !!
- **AXEL** detector
 - 1ton High-Pressure Xenon TPC
 - Unique readout system “**ELCC**” using electroluminescence process
 - **Constructing HP180L prototype**
 - Now some pump troubles ...
- Various R&Ds
 - Optimization of ELCC design
 - Non-discharge Field cage & Cockcroft-Walton circuit
 - Topology Identification w/ Deep Learning & Positive-Ion detection
 - (Xenon gas purification)

Talk by Yoshida

Talk by Nakamura

