

PSI

THEORETICAL OVERVIEW FOR FUTURE LHC HIGGS PHYSICS

Michael Spira (PSI)

I Introduction

II Higgs Boson Decays

III Higgs Boson Production

IV Conclusions

I INTRODUCTION

- Higgs Boson Production







LHC Higgs XS WG

- Discovery: LHC [Tevatron]
  - → Higgs mass
    couplings
    spin
    $\mathcal{CP}$
    λ ?



II HIGGS BOSON DECAYS

Partial Width	QCD	Electroweak	Total	on-shell Higgs
$H \to b\bar{b}/c\bar{c}$	$\sim 0.2\%$	$\sim 0.5\%$	$\sim 0.5\%$	NNNNLO / NLO
$H \to \tau^+\tau^-/\mu^+\mu^-$		$\sim 0.5\%$	$\sim 0.5\%$	NLO
$H \to gg$	$\sim 3\%$	$\sim 1\%$	$\sim 3\%$	NNNLO approx. / NLO
$H \to \gamma\gamma$	$< 1\%$	$< 1\%$	$\sim 1\%$	NLO / NLO
$H \to Z\gamma$	$< 1\%$	$\sim 5\%$	$\sim 5\%$	(N)LO / LO $\leftarrow$ NLO
$H \to WW/ZZ \to 4f$	$< 0.5\%$	$\sim 0.5\%$	$\sim 0.5\%$	(N)NLO

- QCD: variation $\mu_R = [1/2, 2]\mu_0$
  elw: missing HO estimated from known structure at NLO

- parametric uncertainties:
  $m_t = 172.5 \pm 1$ GeV $\alpha_s(M_Z) = 0.118 \pm 0.0015$
  $m_b(m_b) = 4.18 \pm 0.03$ GeV $m_c(3\text{GeV}) = 0.986 \pm 0.025$ GeV
  different uncertainties added quadratically for each channel

- total uncertainties: parametric & theor. uncertainties added linearly

YR3 HDECAY & Prophecy4f YR4

$\sim \sim$ 10% $\rightarrow$
$\sim$ 10% $\rightarrow$



$\longrightarrow$



Denner, Heinemeyer, Puljak, Rebuzzi, S.

- refinements input parameters
- full NLO elw. corrections to $H \to f\bar{f}$
- NLO quark-mass effects in $H \to gg$

III HIGGS BOSON PRODUCTION

(i) $gg \to H$



Georgi,. . .

S., Djouadi, Graudenz, Zerwas
Dawson, Kauffman

- NLO QCD corrections: $\sim 100\%$

- NNLO calculated for $m_t \gg M_\phi \Rightarrow$ further increase by 20–30%
  [top mass effects small in SM]

Harlander, Kilgore
Anastasiou, Melnikov
Ravindran, Smith, van Neerven

Marzani, Ball, Del Duca, Forte, Vicini
Harlander, Ozeren
Pak, Rogal, Steinhauser
Czakon, Harlander, Klappert, Niggetied

- N^{3}LO for $m_t \gg M_\phi \quad \Rightarrow$ scale stabilization
scale dependence: $\Delta \lesssim 5\%$

Moch, Vogt
Ravindran
de Florian, Mazzitelli, Moch, Vogt
Anastasiou, Duhr, Dulat, Furlan, Gehrmann, Herzog, Mistlberger
Ball, Bonvini, Forte, Marzani, Ridolfi

- N^{3}LL soft gluon resummation: $\lesssim 1\%$

Catani, de Florian, Grazzini, Nason
Ravindran
Ahrens, Becher, Neubert, Yang
Ball, Bonvini, Forte, Marzani, Ridolfi
Bonvini, Marzani
Schmidt, S.

- impl. of $gg \to \phi$ in POWHEG including mass effects @ NLO
  (QCD also valid for 2HDM and other Higgs extensions)

Bagnaschi, Degrassi, Slavich, Vicini

- elw. corrections: $\sim 5\%$

Aglietti,. . .
Degrassi, Maltoni
Actis, Passarino, Sturm, Uccirati

- $\sigma(gg \to H) = \left(52.17^{+4.6\%}_{-6.7\%}(TH) \pm 3.2\%(PDF, \alpha_s)\right)\ pb$ @ $\sqrt{s} = 13.6$ TeV

Anastasiou,. . .

- uncertainties: PDF+α_s, renormalization/factorization scale
  top/bottom masses: $\sim \pm 0.8\%$ $\leftarrow$ scale/scheme dependence



stolen from M. Grazzini@Higgs10



stolen from M. Grazzini@Higgs10



stolen from M. Grazzini@Higgs10



stolen from M. Grazzini@Higgs10



stolen from M. Grazzini@Higgs10

$\sigma(gg \to H)_{LO} = 18.43^{+0.8\%}_{-1.1\%}$ pb $\qquad$ $\sigma(gg \to H)^{QCD}_{NLO} = 42.17^{+0.4\%}_{-0.5\%}$ pb



Jones, S.

m_t scheme/scale uncertainties only:

• LO:

$\sigma(gg \to H^*)|_{Q=125\ \text{GeV}} = 18.43^{+0.8\%}_{-1.1\%}\ \text{pb}, \qquad \sigma(gg \to H^*)|_{Q=300\ \text{GeV}} = 4.88^{+23.1\%}_{-1.1\%}\ \text{pb}$

$\sigma(gg \to H^*)|_{Q=400\ \text{GeV}} = 4.94^{+1.2\%}_{-1.8\%}\ \text{pb}, \qquad \sigma(gg \to H^*)|_{Q=600\ \text{GeV}} = 1.13^{+0.0\%}_{-26.2\%}\ \text{pb}$

$\sigma(gg \to H^*)|_{Q=900\ \text{GeV}} = 0.139^{+0.0\%}_{-36.0\%}\ \text{pb}, \quad \sigma(gg \to H^*)|_{Q=1200\ \text{GeV}} = 0.0249^{+0.0\%}_{-41.1\%}\ \text{pb}$

• NLO QCD:

$\sigma(gg \to H^*)|_{Q=125\ \text{GeV}} = 42.17^{+0.4\%}_{-0.5\%}\ \text{pb}, \qquad \sigma(gg \to H^*)|_{Q=300\ \text{GeV}} = 9.85^{+7.5\%}_{-0.3\%}\ \text{pb}$

$\sigma(gg \to H^*)|_{Q=400\ \text{GeV}} = 9.43^{+0.1\%}_{-0.9\%}\ \text{pb}, \qquad \sigma(gg \to H^*)|_{Q=600\ \text{GeV}} = 1.97^{+0.0\%}_{-15.9\%}\ \text{pb}$

$\sigma(gg \to H^*)|_{Q=900\ \text{GeV}} = 0.230^{+0.0\%}_{-22.3\%}\ \text{pb}, \quad \sigma(gg \to H^*)|_{Q=1200\ \text{GeV}} = 0.0402^{+0.0\%}_{-26.0\%}\ \text{pb}$

(ii) W/Z fusion: $pp \to W^*W^*/Z^*Z^* \to h/H$



Cahn, Dawson
Hikasa
Atarelli, Mele, Pitolli

- QCD corrections $\leftarrow$ DIS: $\sim 10\%$

Han, Valencia,
Willenbrock
Figy, Oleari, Zeppenfeld
Berger, Campbell

[approx] 2–loop: $\lesssim 1\%$

Bolzano, Maltoni, Moch, Zaro
Cacciari, Dreyer, Karlberg, Salam, Zanderighi

[approx] 3–loop: $\lesssim 0.3\%$

Dreyer, Karlberg

- elw. corrections: $\sim 10\%$

Ciccolini, Denner, Dittmaier

- fully exclusive @ NNLO QCD, NLO elw.

- $\sigma(VBF) = \left(4.075^{+0.5\%}_{-0.3\%}(TH) \pm 2.1\%(PDF, \alpha_s)\right)\ pb$ @ $\sqrt{s} = 13.6$ TeV

(iii) Higgs–strahlung: $pp \to W^*/Z^* \to W/Z + h/H$



Glashow,. . .
Kunszt,. . .

- QCD corrections ← DY: $\sim 30\%$
  2–loop: $\lesssim 5\%$ — Han, Willenbrock; Brein, Djouadi, Harlander
- electroweak corrections: $\sim -10\%$ — Ciccolini, Dittmaier, Krämer
- $W/Z + H$: fully exclusive @ NNLO QCD — Ferrera, Grazzini, Tramantano



$$\sigma(WH) = \left(1.453^{+0.4\%}_{-0.7\%}(TH) \pm 1.8\%(PDF, \alpha_s)\right)\ pb$$

$$\sigma(ZH) = \left(0.9422^{+3.8\%}_{-3.2\%}(TH) \pm 1.6\%(PDF, \alpha_s)\right)\ pb$$

@ $\sqrt{s} = 13.6$ TeV

(iv) Bremsstrahlung: $pp \to t\bar{t} + h/H/A$





Kunszt
Gunion
Marciano, Paige

dominant

- $t\bar{t}h \to t\bar{t}b\bar{b}$ important @ LHC $\to$ top Yukawa cplg.

- QCD corrections [SM]: $\sim 20\%$ Beenakker, Dittmaier, Krämer, Plümper, S., Zerwas
  Dawson, Orr, Reina, Wackeroth
  [threshold suppressed: $\sigma_{LO} \sim \beta^4$] Broggio, Ferroglia, Pecjak, Signer, Yang

- link to parton showers: aMC@NLO, PowHel Frederix et al.
  Garzelli, Kardos, Papadopoulos, Trócsányi

- important work on backgrounds $t\bar{t}b\bar{b}, t\bar{t}jj$, etc.
  Bredenstein, Denner, Dittmaier, Pozzorini
  Bevilacqua, Czakon, Papadopoulos, Pittau, Worek
  Cascioli, Maierhofer, Pozzorini

- $\sigma(t\bar{t}H) = \left(0.5688^{+6.0\%}_{-9.3\%}(TH) \pm 3.5\%(PDF, \alpha_s)\right)\ pb$ @ $\sqrt{s} = 13.6$ TeV

$pp \to t\bar{t}H$: approx. NNLO



$m_t = 173.3$ GeV



sensitive to top Yukawa coupling

Catani, Devoto, Grazzini, Kallweit, Mazzitelli, Savoini

- 2-loop virtual corrections: soft Higgs approximation
  single + double real corrections: exact

(v) $gg \to HH$



- threshold region: sensitive to λ
  large M_{HH}: sensitive to $c_{tt/bb}$ [e.g. boosted Higgs pairs]



$$gg \to HH : \frac{\Delta\sigma}{\sigma} \sim -\frac{\Delta\lambda}{\lambda}$$

[decreasing with M^2_{HH}]

Baglio, Djouadi, Gröber, Mühlleitner, Quevillon, S.

Overview

(Slide design shamelessly stolen from G. Salam)



[1] LO



[2] NLO HTL



[5,6] NNLO HTL



[8] Full reals ($\mathrm{FT}_{\mathrm{approx}}$)

88

98

13

14

15

16

17

18

19

21

22

23

24

25

$+\mathcal{O}(1/m_T)$

[3] NLO HTL + NNLL

[4] NLO $1/m_T^8$

[7] NLO $1/m_T^{12}$

[9] NNLO virt $1/m_T^4$

$+\mathcal{O}(1/m_T)$

[10] NNLO HTL (Fully Diff)

[11,12] NLO numerical



[13] NLO + NLL

[14,15] NLO + PS

[16] $1/m_T$ + thres.

[17,28] NLO $\overline{\mathrm{MS}}$ m_T

[18] NNLO ($\mathrm{FT}_{\mathrm{approx}}$)

[19] NNLO + NNLL

[20] NLO small-p_T

[21,22] NLO small-m_T

[23] NLO small-p_T + thres.

[24] NLO num. + small-m_T

[25,27] NNLO $1/m_T^2$



[26] N^{3}LO HTL



[29] small-p_T + small-m_T

[30] EW: leading Yukawa vs λ_3^{eff}

[31] EW: leading Yukawa

[32] N^{3}LO + N^{3}LL + NLO m_t

[33] small-t

[34] NNLO: n_f contrib.

[35] EW: $1/m_T$

[36] NLO comb expansions

[37] NLO EW corrections

[38] NLO + $b\bar{b}\gamma\gamma$ decays

[39] NNLO: reducible

[40] EW: Yukawa/self-coupling

[42] EW: self-coupling

[43] EW: Factorizable

[44] EW: Yukawa/self-coupling small-m_t

[45] NNLO: Large-N_c small-p_T

[46] EW: Light quark induced

[1] Glover, van der Bij 88; [2] Dawson, Dittmaier, Spira 98; [3] Shao, Li, Li, Wang 13; [4] Grigo, Hoff, Melnikov, Steinhauser 13; [5] de Florian, Mazzitelli 13; [6] Grigo, Melnikov, Steinhauser 14; [7] Grigo, Hoff 14; [8] Maltoni, Vryonidou, Zaro 14; [9] Grigo, Hoff, Steinhauser 15; [10] de Florian, Grazzini, Hanga, Kallweit, Lindert, Maierhöfer, Mazzitelli, Rathlev 16; [11] Borowka, Greiner, Heinrich, SPJ, Kerner, Schlenk, Schubert, Zirke 16; [12] Borowka, Greiner, Heinrich, SPJ, Kerner, Schlenk, Zirke 16; [13] Ferrera, Pires 16; [14] Heinrich, SPJ, Kerner, Luisoni, Vryonidou 17; [15] SPJ, Kuttimalai 17; [16] Gröber, Maier, Rauh 17; [17] Baglio, Campanario, Glaus, Mühlleitner, Spira, Streicher 18; [18] Grazzini, Heinrich, SPJ, Kallweit, Kerner, Lindert, Mazzitelli 18; [19] de Florian, Mazzitelli 18; [20] Bonciani, Degrassi, Giardino, Gröber 18; [21] Davies, Mishima, Steinhauser, Wellmann 18, 18; [22] Mishima 18; [23] Gröber, Maier, Rauh 19; [24] Davies, Heinrich, SPJ, Kerner, Mishima, Steinhauser, David Wellmann 19; [25] Davies, Steinhauser 19; [26] Chen, Li, Shao, Wang 19, 19; [27] Davies, Herren, Mishima, Steinhauser 19, 21; [28] Baglio, Campanario, Glaus, Mühlleitner, Ronca, Spira 21; [29] Bellafronte, Degrassi, Giardino, Gröber, Vitti 22; [30] Mühlleitner, Schlenk, Spira 22; [31] Davies, Mishima, Schönwald, Steinhauser, Zhang 22; [32] Ajjath, Shao 22; [33] Davies, Mishima, Schönwald, Steinhauser 23; [34] Davies, Schönwald, Steinhauser 23; [35] Davies, Schönwald, Steinhauser, Zhang 23; [36] Bagnaschi, Degrassi, Gröber 23; [37] Bi, Huang, Huang, Ma Yu 23 [38] Li, Si, Wang, Zhang, Zhao 24; [39] Davies, Schönwald, Steinhauser, Vitti 24; [40] Heinrich, SPJ, Kerner, Stone, Vestner [41] Li, Si, Wang, Zhang, Zhao 24; [42] Davies, Schönwald, Steinhauser, Zhang 24; [43] Davies, Schönwald, Steinhauser, Zhang 25; [44] Davies, Schönwald, Steinhauser 25; [45] Bonetti, Rendler, Bobadilla 25;

S. Jones


σ (HH) [fb]
60
50
40
30
20
10
0
LO
NLO HTL
NNLO HTL
N3LO HTL
NNLO FTapprox
NNLO FTapprox + mtNLO
N3LO HTL + mtNLO

Full NLO calculation: top only, numerical integration

Borowka *et al.*	Baglio *et al.*
tensor reduction	no tensor reduction
sector decomposition	IR, end-point subtraction
contour deformation	IBP, Richardson extrapolation
$m_t = 173$ GeV	$m_t = 172.5$ GeV

Borowka, Greiner, Heinrich, Jones, Kerner, Schlenk, Schubert, Zirke
Baglio, Campanario, Glaus, Mühlleitner, Ronca, S., Streicher



Borowka, Greiner, Heinrich, Jones, Kerner
Schlenk, Schubert, Zirke



Baglio, Campanario, Glaus,
Mühlleitner, Ronca, S., Streicher

$$\sigma_{NLO} = 32.91(10)^{+13.8\%}_{-12.8\%}\ fb \qquad 32.81(7)^{+13.5\%}_{-12.5\%}\ fb$$

$$\sigma^{HTL}_{NLO} = 38.75^{+18\%}_{-15\%}\ fb \qquad 38.66^{+18\%}_{-15\%}\ fb$$

$$m_t = 173\ \text{GeV} \qquad 172.5\ \text{GeV}$$

$\Rightarrow$ -15% mass effects on top of LO


σ (HH) [fb]
60
50
40
30
20
10
0
LO
NLO HTL
NNLO HTL
N3LO HTL
NNLO FTapprox
NNLO FTapprox + mtNLO
N3LO HTL + mtNLO

uncertainties due to m_t

- use m_t, $\overline{m}_t(\overline{m}_t)$ and scan $Q/4 < \mu < Q \rightarrow$ uncertainty = envelope:

$$\frac{d\sigma(gg \to HH)}{dQ}|_{Q=300\ \text{GeV}} = 0.02978(7)^{+6\%}_{-34\%}\ \text{fb/GeV},$$

$$\frac{d\sigma(gg \to HH)}{dQ}|_{Q=400\ \text{GeV}} = 0.1609(4)^{+0\%}_{-13\%}\ \text{fb/GeV},$$

$$\frac{d\sigma(gg \to HH)}{dQ}|_{Q=600\ \text{GeV}} = 0.03204(9)^{+0\%}_{-30\%}\ \text{fb/GeV},$$

$$\frac{d\sigma(gg \to HH)}{dQ}|_{Q=1200\ \text{GeV}} = 0.000435(4)^{+0\%}_{-35\%}\ \text{fb/GeV}$$

- bin-by-bin interpolation:

$$\sigma(gg \to HH) = 32.81^{+4\%}_{-18\%}\ \text{fb}$$

- why a dynamical scale $\sim Q$?
  large momentum expansion ($\hat{s} = Q^2 \gg m_t^2$), two FF:

← Davies, Mishima, Steinhauser, Wellmann

pole mass m_t:

$$\Delta F_{1,mass} \rightarrow \frac{\alpha_s}{\pi}\left\{2F_{1,LO}\log\frac{m_t^2}{\hat{s}} + \frac{m_t^2}{\hat{s}}G_1(\hat{s},\hat{t})\right\},$$

$$\Delta F_{2,mass} \rightarrow \frac{\alpha_s}{\pi}\left\{2F_{2,LO}\log\frac{m_t^2}{\hat{s}} + \frac{m_t^2}{\hat{s}}G_2(\hat{s},\hat{t})\right\}$$

$\overline{\text{MS}}$ mass $\overline{m}_t(\mu_t)$:

$$\Delta F_{1,mass} \rightarrow \frac{\alpha_s}{\pi}\left\{2F_{1,LO}\left[\log\frac{\mu_t^2}{\hat{s}} + \frac{4}{3}\right] + \frac{\overline{m}_t^2(\mu_t)}{\hat{s}}G_1(\hat{s},\hat{t})\right\},$$

$$\Delta F_{2,mass} \rightarrow \frac{\alpha_s}{\pi}\left\{2F_{2,LO}\left[\log\frac{\mu_t^2}{\hat{s}} + \frac{4}{3}\right] + \frac{\overline{m}_t^2(\mu_t)}{\hat{s}}G_2(\hat{s},\hat{t})\right\}$$

$\Rightarrow$ scale $\mu_t \sim Q$ preferred at large Q



Baglio, Campanario, Glaus, Mühlleitner, Ronca, S.

- renormalization/factorization scale uncertainties @ NLO:

$$\begin{aligned}
\sqrt{s} = 13 \text{ TeV}: \quad \sigma_{tot} &= 27.73(7)^{+13.8\%}_{-12.8\%} \text{ fb} \\
\sqrt{s} = 14 \text{ TeV}: \quad \sigma_{tot} &= 32.81(7)^{+13.5\%}_{-12.5\%} \text{ fb} \\
\sqrt{s} = 27 \text{ TeV}: \quad \sigma_{tot} &= 127.0(2)^{+11.7\%}_{-10.7\%} \text{ fb} \\
\sqrt{s} = 100 \text{ TeV}: \quad \sigma_{tot} &= 1140(2)^{+10.7\%}_{-10.0\%} \text{ fb}
\end{aligned}$$

- m_t scale/scheme uncertainties @ NLO:

$$\begin{aligned}
\sqrt{s} = 13 \text{ TeV}: \quad \sigma_{tot} &= 27.73(7)^{+4\%}_{-18\%} \text{ fb} \\
\sqrt{s} = 14 \text{ TeV}: \quad \sigma_{tot} &= 32.81(7)^{+4\%}_{-18\%} \text{ fb} \\
\sqrt{s} = 27 \text{ TeV}: \quad \sigma_{tot} &= 127.8(2)^{+4\%}_{-18\%} \text{ fb} \\
\sqrt{s} = 100 \text{ TeV}: \quad \sigma_{tot} &= 1140(2)^{+3\%}_{-18\%} \text{ fb}
\end{aligned}$$

- renormalization/factorization scale uncertainties @ NNLO$_{FTapprox}$:

$$\begin{aligned}
\sqrt{s} = 13 \text{ TeV}: \quad \sigma_{tot} &= 31.05^{+2.2\%}_{-5.0\%} \text{ fb} \\
\sqrt{s} = 14 \text{ TeV}: \quad \sigma_{tot} &= 36.69^{+2.1\%}_{-4.9\%} \text{ fb} \\
\sqrt{s} = 27 \text{ TeV}: \quad \sigma_{tot} &= 139.9^{+1.3\%}_{-3.9\%} \text{ fb} \\
\sqrt{s} = 100 \text{ TeV}: \quad \sigma_{tot} &= 1224^{+0.9\%}_{-3.2\%} \text{ fb}
\end{aligned}$$

- HO corrections: dominated by universal S+V+C corrections

$\Rightarrow \sim$ rescaling of rel. m_t scale/scheme uncertainties
combination $\rightarrow$ envelope $\sim$ linear sum (rel. err.)

final combined ren./fac. scale and m_t scale/scheme unc. @ NNLO$_{FTapprox}$:

$$\sqrt{s} = 13 \text{ TeV}: \quad \sigma_{tot} = 31.05^{+6\%}_{-23\%} \text{ fb}$$

$$\sqrt{s} = 14 \text{ TeV}: \quad \sigma_{tot} = 36.69^{+6\%}_{-23\%} \text{ fb}$$

$$\sqrt{s} = 27 \text{ TeV}: \quad \sigma_{tot} = 139.9^{+5\%}_{-22\%} \text{ fb}$$

$$\sqrt{s} = 100 \text{ TeV}: \quad \sigma_{tot} = 1224^{+4\%}_{-21\%} \text{ fb}$$

λ dependence



Baglio, Campanario, Glaus, Mühlleitner, Ronca, S.

- final combined uncertainties @ NNLO$_{FTapprox}$ ($\sqrt{s} = 14$ TeV):

$$
\begin{aligned}
\kappa_\lambda = -10: \quad \sigma_{tot} &= 1680^{+13\%}_{-14\%} \text{ fb} \\
\kappa_\lambda = -5: \quad \sigma_{tot} &= 598.9^{+13\%}_{-15\%} \text{ fb} \\
\kappa_\lambda = -1: \quad \sigma_{tot} &= 131.9^{+11\%}_{-16\%} \text{ fb} \\
\kappa_\lambda = 0: \quad \sigma_{tot} &= 70.38^{+8\%}_{-18\%} \text{ fb} \\
\kappa_\lambda = 1: \quad \sigma_{tot} &= 31.05^{+6\%}_{-23\%} \text{ fb} \\
\kappa_\lambda = 2: \quad \sigma_{tot} &= 13.81^{+3\%}_{-28\%} \text{ fb} \\
\kappa_\lambda = 2.4: \quad \sigma_{tot} &= 13.10^{+6\%}_{-27\%} \text{ fb} \\
\kappa_\lambda = 3: \quad \sigma_{tot} &= 18.67^{+12\%}_{-22\%} \text{ fb} \\
\kappa_\lambda = 5: \quad \sigma_{tot} &= 94.82^{+18\%}_{-13\%} \text{ fb} \\
\kappa_\lambda = 10: \quad \sigma_{tot} &= 672.2^{+16\%}_{-13\%} \text{ fb}
\end{aligned}
$$

reduction of these uncertainties: full NNLO corrs. [not available]

[partial virtual NNLO corr. known]

Grigo, Hoff, Steinhauser
Davies, Schönwald, Steinhauser

reduction of these uncertainties: full NNLO corrs. [not available]
[partial virtual NNLO corr. known]

Grigo, Hoff, Steinhauser
Davies, Schönwald, Steinhauser

- resummation of $\log(M_{HH}^2/m_t^2)$ [SCET]:

Jaskiewicz, Jones, Szafron, Ulrich

$$\overline{m}(\mu) = M z_m(\mu) \qquad \mu \sim M_{HH}$$

$$m^{LL}(\mu) = M \ \exp[a_{\gamma_m}^{LL}(\mu)] \ z_m(M) \qquad \sim \overline{m}(\mu) \text{ @ LL}$$

$$a_{\gamma_m}^{LL}(\mu) = \frac{3C_F}{2\beta_0} \log\left(1 - \frac{\alpha_s(\mu)}{2\pi}\beta_0 \log\frac{\mu^2}{M^2}\right) \qquad \left[\beta_0 = \frac{33-2N_F}{6}\right]$$

$$z_m(\mu) = 1 + \frac{\alpha_s(\mu)}{2\pi}\left(-2C_F - \frac{3}{2}C_F \log\frac{\mu^2}{M^2}\right)$$

scheme dependence $\rightarrow$ matching scale dependence (@ LL):

$$m^{LL}(\mu_t, \mu_m) = M \ \exp[a_{\gamma_m}^{LL}(\mu_t, \mu_m)] \ z_m(\mu_m)$$

$$a_{\gamma_m}^{LL}(\mu_t, \mu_m) = \frac{3C_F}{2\beta_0} \log\left(1 - \frac{\alpha_s(\mu_t)}{2\pi}\beta_0 \log\frac{\mu_t^2}{\mu_m^2}\right) = -\frac{3C_F}{2\beta_0} \log\left(1 + \frac{\alpha_s(\mu_m)}{2\pi}\beta_0 \log\frac{\mu_t^2}{\mu_m^2}\right)$$

$$z_m(\mu_m) = 1 + \frac{\alpha_s(\mu)}{2\pi}\left(-2C_F - \frac{3}{2}C_F \log\frac{\mu_m^2}{M^2}\right)$$

μ_t dependence still significant [16% $\rightarrow$ 8% for virt], μ_m?

OPEN FOR DISCUSSIONS!

- pole mass $\leftrightarrow$ $\overline{\text{MS}}$ mass:

$$\overline{m}_t(M_t) = \frac{M_t}{1 + \frac{4}{3}\frac{\alpha_s(M_t)}{\pi} + 10.9\left(\frac{\alpha_s(M_t)}{\pi}\right)^2}$$

$$\overline{m}_t(\mu) \;=\; \overline{m}_t(M_t)\,\frac{c\left[\alpha_s(\mu)/\pi\right]}{c\left[\alpha_s(M_t)/\pi\right]}$$

$$c(x) = (2\beta_0\, x)^{\frac{2\gamma_0}{\beta_0}}\,[1 + 1.398x + 1.793\,x^2 - 0.6834\,x^3] \qquad [\gamma_0 = 3C_F/4 = 1]$$

$$M_t = 172.5 \text{ GeV} \qquad \overline{m}_t(\overline{m}_t) = 163.0 \text{ GeV}$$

$$\alpha_s(\mu) \;=\; \frac{\alpha_s(M_t)}{1 + \beta_0\frac{\alpha_s(M_t)}{2\pi}\log\frac{\mu^2}{M_t^2}} \qquad @\ \text{LL}$$

$$\overline{m}_t(\mu) \;=\; \overline{m}_t(M_t)\left(1 + \beta_0\frac{\alpha_s(M_t)}{2\pi}\log\frac{\mu^2}{M_t^2}\right)^{-2/\beta_0} \qquad @\ \text{LL}$$

Electroweak corrections

(i) y_t: HTL for $ggH(H)$ coupling + full corrections to HHH vertex

Mühlleitner, Schlenk, S.

(ii) y_t: analytical results for $ggHH$ coupling in the HEL

Davies, Mishima, Schönwald, Steinhauser, Zhang

and close to the production threshold

Davies, Schönwald, Steinhauser, Zhang

(iii) λ: elw. corrections due to the Higgs self-int. [SMEFT @ D8]

Borowka, Duhr, Maltoni, Pagani, Shivaji, Zhao

(iv) g_t, λ: elw. corrections due to the top Yukawa and Higgs self-interactions [only Higgs exchange diagrams]

Heinrich, Jones, Kerner, Stone, Vestner

(v) full elw. corrections ($\leftarrow$ to be checked & understood)

Bi, Huang, Huang, Ma, Yu

Full electroweak corrections





- -4.2% for total cross section

Bi, Huang, Huang, Ma, Yu

Top-Yukawa-induced elw. corrections

Mühlleitner, Schlenk, S.



(i) effective $ggH(H)$ couplings:

$$\mathcal{L}_{eff} = C_1 \frac{\alpha_s}{12\pi} G^{a\mu\nu} G^a_{\mu\nu} \log\left(1 + C_2 \frac{H}{v}\right)$$



(C_1) (C_2)

- $C_1 = 1 - 3x_t$: genuine vertex corrections $\quad [x_t = G_F m_t^2/(8\sqrt{2}\pi^2)]$

Djoaudi, Gambino
Chetyrkin, Kniehl, Steinhauser

- $C_2 = 1 + 7x_t/2$ $[= 1 + \delta Z_H/2 - \delta v/v]$: universal corrections

Kniehl, Spira
Kwiatkowski, Steinhauser

$$\mathcal{L}_{eff} = \frac{\alpha_s}{12\pi} G^{a\mu\nu} G^a_{\mu\nu} \left\{ (1+\delta_1)\frac{H}{v} + (1+\eta_1)\frac{H^2}{2v^2} + \mathcal{O}(H^3) \right\}$$

$$\delta_1 = \frac{x_t}{2} + \mathcal{O}(x_t^2) \qquad \eta_1 = 4x_t + \mathcal{O}(x_t^2)$$

elw. gaugeless limit + QCD = top-Yukawa model + QCD

$$\phi = \begin{pmatrix} G^+ \\ \dfrac{v + H + iG^0}{\sqrt{2}} \end{pmatrix}$$

$$\begin{aligned}
\mathcal{L} &= -\frac{1}{4} G^{a\mu\nu} G^a_{\mu\nu} + \bar{t} i \not{D} t + |\partial_\mu \phi|^2 - V(\phi) - g_t \bar{Q}_L \phi^c t_R \\
V(\phi) &= -\mu^2 |\phi|^2 + \frac{\lambda}{2} |\phi|^4 \\
&= -\frac{M_H^2}{8} v^2 + \frac{M_H^2}{2} H^2 + \frac{M_H^2}{v} \left[\frac{H^3}{2} + \frac{H}{2} (G^0)^2 + H G^+ G^- \right] \\
&\quad + \frac{M_H^2}{2v^2} \left[\frac{H^4}{4} + \frac{H^2}{2} (G^0)^2 + H^2 G^+ G^- + (G^+ G^-)^2 + (G^0)^2 G^+ G^- + \frac{(G^0)^4}{4} \right]
\end{aligned}$$

$$\Rightarrow M_{G^0} = M_{G^\pm} = 0$$

INTERMEZZO

Full top-mass dependence (wave-function ren. adjusted appropriately)



Light-quark loops



Aglietti, Bonciani, Degrassi, Vicini

→ Bonetti, Rendler, Torres Bobadilla

$$\delta_1 = \delta_{2loop} + \delta Z_H/2 - \delta v/v$$



Bhattacharya, Campanario, Carlotti, Chang, Mazzitelli, Mühlleitner, Ronca, S.

- box diagrams in the making. . .

BOXES

Topology 1



Topology 2



TOPOLOGY 3





Bhattacharya, Campanario, Carlotti, Chang, Mazzitelli, Mühlleitner, Ronca, S.

(ii) effective $HHH(H)$ couplings:

- effective Higgs potential: Coleman, Weinberg

$$
\begin{aligned}
V_{eff} &= V_0 + V_1 \\
V_0 &= \mu_0^2|\phi|^2 + \frac{\lambda_0}{2}|\phi|^4 \\
V_1 &= \frac{3\overline{m}_t^4}{16\pi^2}\Gamma(1+\epsilon)(4\pi^2)^\epsilon\left(\frac{1}{\epsilon} + \log\frac{\bar{\mu}^2}{\overline{m}_t^2} + \frac{3}{2}\right) \\
\phi &= \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v+H \end{pmatrix} \qquad \overline{m}_t = m_t\left(1+\frac{H}{v}\right)
\end{aligned}
$$

- after renormalization

$$
\begin{aligned}
\lambda_{HHH}^{eff} &= 3\frac{M_H^2}{v} + \Delta\lambda_{HHH}\,, & \lambda_{HHHH}^{eff} &= 3\frac{M_H^2}{v^2} + \Delta\lambda_{HHHH} \\
\Delta\lambda_{HHH} &= -\frac{3m_t^4}{\pi^2 v^3}\,, & \Delta\lambda_{HHHH} &= -\frac{12m_t^4}{\pi^2 v^4}
\end{aligned}
$$

$$
\boxed{\lambda_{HHH}^{eff} = 3\frac{M_H^2}{v} - \frac{3m_t^4}{\pi^2 v^3} \approx 0.91 \times 3\frac{M_H^2}{v}}
$$



$$\sigma = 1.002 \times \sigma_{LO} \qquad (\lambda_{HHH})$$

$$\sigma = 0.938 \times \sigma_{LO} \qquad (\lambda^{eff}_{HHH}) \leftarrow \text{disfavoured}$$

Mühlleitner, Schlenk, S.

IV *CONCLUSIONS*

- Higgs boson searches/studies at LHC belong to major endeavours
- important to develop NLO event generators [$\leftarrow$ backgrounds]
- scale and scheme uncertainties due to m_t relevant for large momenta
- significant uncertainties for off-shell Higgs production and decays
- Higgs pair production: m_t effects on top of LO $\sim -15\%$ for σ_{tot}
  [larger for distributions]
- factorization/renormalization scale uncertainties @NNLO$_{FTapprox}$ $\lesssim 5\%$
- uncertainties due to m_t scale/scheme choice sizeable $\lesssim 20\%$
  $\rightarrow$ linear combination of rel. uncertainties
- analogous issues in $gg \rightarrow H + 2jet, ZH, ZZ$ etc.
- small top-induced electroweak corrections

BACKUP SLIDES

$$\Gamma[H \to b\bar{b}] = \frac{3 G_F M_H}{4\sqrt{2}\pi}\, \overline{m}_b^2(M_H)\, \Delta_{\rm QCD}$$

$\uparrow$

log resummation $\to \sim$ factor 1/2

(larger than BSM effects!)



Braaten, Leveille
Drees, Hikasa
Kataev,. . .
Chetyrkin,. . .
etc.

→ HDECAY

Djouadi, Kalinowski, Mühlleitner, S.

- ATLAS: $\mu_{bb}/\mu_{ZZ} = \Gamma(H \to bb)/\Gamma(H \to ZZ)|_{SM-norm} = 0.87^{+0.28}_{-0.21}$

- CMS: $\mu_{bb}/\mu_{ZZ} = \Gamma(H \to bb)/\Gamma(H \to ZZ)|_{SM-norm} = 0.84^{+0.37}_{-0.27}$

$\Rightarrow \overline{m}_b(M_H) = 2.60^{+0.36}_{-0.31}$ GeV
(BLUE) Nisius

RG-evolution: REvolver

Hoang, Lepenik, Mateu



Aparisi, Fuster, Irles, Rodrigo, Vos, Yamamoto, Hoang, Lepenik, S., Tarafune, Yonamine

- Higgs + jet production: $gg \to H + j$



LO: $\mu_t = H_T/2 = (\sqrt{M_H^2 + p_T^2} + p_{Tj})/2$



→ NLO? Jones, Kerner, Luisoni

Jones, S.

(v) $b\bar{b}$+Higgs production



NLO



$[\leftarrow \Delta_b]$

NNLO

exact $g \to b\bar{b}$ splitting & mass/off-shell effects

no resummation of $\log M_H^2/m_b^2$ terms

massless/on-shell b's, no p_{Tb}

resummation of $\log M_H^2/m_b^2$ terms



Santander matching:

$$\sigma = \frac{\sigma^{4FS} + w\sigma^{5FS}}{1+w}$$

$$w = \log\frac{M_H}{m_b} - 2$$

Harlander, Krämer, Schumacher

Dittmaier, Krämer, S.
Dawson, Jackson, Reina, Wackeroth
Harlander, Kilgore

matching



Bonvini, Papanastasiou, Tackmann

Forte, Napoletano, Ubiali

Flavour-scheme comparison

$$H_T = \sum_{i=b,\bar{b},H} \sqrt{m^2(i) + p_T^2(i)}$$

NNLO corrections solve the FS issue.

At NNLO QCD, the two predictions agree within the scale variation when using the most natural choice (m_H) as the central scale, without the need for artificial factors to improve the comparison.

5FS $\mathrm{NLO_{PS}}(m_H)$	$0.676(4)^{+27\%}_{-15\%}$ pb
4FS $\mathrm{NLO_{PS}}(m_H)$	$0.355(3)^{+20\%}_{-16\%}$ pb
4FS $\mathrm{NLO_{PS}}(\frac{H_T}{4})$	$0.385(4)^{+16\%}_{-14\%}$ pb

5FS $\mathrm{MiNNLO_{PS}}(m_H)$	$0.509(4)^{+2.9\%}_{-5.3\%}$ pb
4FS $\mathrm{MiNNLO_{PS}}(m_H)$	$0.465(4)^{+16\%}_{-14\%}$ pb
4FS $\mathrm{MiNNLO_{PS}}(\frac{H_T}{4})$	$0.492(3)^{+16\%}_{-14\%}$ pb





CB, Mazzitelli, Sankar, Wiesemann, Zanderighi [to appear]

- m_t scale/scheme uncertainties @ NLO:

$$\begin{aligned}
\kappa_\lambda = -10: \quad \sigma_{tot} &= 1438(1)^{+10\%}_{-6\%}\ \text{fb} \\
\kappa_\lambda = -5: \quad \sigma_{tot} &= 512.8(3)^{+10\%}_{-7\%}\ \text{fb} \\
\kappa_\lambda = -1: \quad \sigma_{tot} &= 113.66(7)^{+8\%}_{-9\%}\ \text{fb} \\
\kappa_\lambda = 0: \quad \sigma_{tot} &= 61.22(6)^{+6\%}_{-12\%}\ \text{fb} \\
\kappa_\lambda = 1: \quad \sigma_{tot} &= 27.73(7)^{+4\%}_{-18\%}\ \text{fb} \\
\kappa_\lambda = 2: \quad \sigma_{tot} &= 13.2(1)^{+1\%}_{-23\%}\ \text{fb} \\
\kappa_\lambda = 2.4: \quad \sigma_{tot} &= 12.7(1)^{+4\%}_{-22\%}\ \text{fb} \\
\kappa_\lambda = 3: \quad \sigma_{tot} &= 17.6(1)^{+9\%}_{-15\%}\ \text{fb} \\
\kappa_\lambda = 5: \quad \sigma_{tot} &= 83.2(3)^{+13\%}_{-4\%}\ \text{fb} \\
\kappa_\lambda = 10: \quad \sigma_{tot} &= 579(1)^{+12\%}_{-4\%}\ \text{fb}
\end{aligned}$$

- renormalization/factorization scale uncertainties @ NNLO$_{FTapprox}$:

$$\begin{aligned}
\kappa_\lambda = -10: \quad \sigma_{tot} &= 1680^{+3.0\%}_{-7.7\%}\ \text{fb} \\
\kappa_\lambda = -5: \quad \sigma_{tot} &= 598.9^{+2.7\%}_{-7.5\%}\ \text{fb} \\
\kappa_\lambda = -1: \quad \sigma_{tot} &= 131.9^{+2.5\%}_{-6.7\%}\ \text{fb} \\
\kappa_\lambda = 0: \quad \sigma_{tot} &= 70.38^{+2.4\%}_{-6.1\%}\ \text{fb} \\
\kappa_\lambda = 1: \quad \sigma_{tot} &= 31.05^{+2.2\%}_{-5.0\%}\ \text{fb} \\
\kappa_\lambda = 2: \quad \sigma_{tot} &= 13.81^{+2.1\%}_{-4.9\%}\ \text{fb} \\
\kappa_\lambda = 2.4: \quad \sigma_{tot} &= 13.10^{+2.3\%}_{-5.1\%}\ \text{fb} \\
\kappa_\lambda = 3: \quad \sigma_{tot} &= 18.67^{+2.7\%}_{-7.3\%}\ \text{fb} \\
\kappa_\lambda = 5: \quad \sigma_{tot} &= 94.82^{+4.9\%}_{-8.8\%}\ \text{fb} \\
\kappa_\lambda = 10: \quad \sigma_{tot} &= 672.2^{+4.2\%}_{-8.5\%}\ \text{fb}
\end{aligned}$$

$$
\begin{aligned}
F_i &= F_{i,LO} + \Delta F_i \\
\Delta F_i &= \Delta F_{i,HTL} + \Delta F_{i,mass}
\end{aligned}
$$

- pole mass:

$$
\begin{aligned}
F_{1,LO} &\rightarrow 4\frac{m_t^2}{\hat{s}} \\
F_{2,LO} &\rightarrow -\frac{m_t^2}{\hat{s}\hat{t}(\hat{s}+\hat{t})}\{(\hat{s}+\hat{t})^2 L_{1ts}^2 + \hat{t}^2 L_{ts}^2 + \pi^2[(\hat{s}+\hat{t})^2 + \hat{t}^2]\}
\end{aligned}
$$

- $\overline{\mathrm{MS}}$ mass:

$$
\begin{aligned}
F_{1,LO} &\rightarrow 4\frac{\overline{m}_t^2(\mu_t)}{\hat{s}} \\
F_{2,LO} &\rightarrow -\frac{\overline{m}_t^2(\mu_t)}{\hat{s}\hat{t}(\hat{s}+\hat{t})}\{(\hat{s}+\hat{t})^2 L_{1ts}^2 + \hat{t}^2 L_{ts}^2 + \pi^2[(\hat{s}+\hat{t})^2 + \hat{t}^2]\}
\end{aligned}
$$