

# Testing indirect experimental method for constraining the $^{193,194}\text{Ir}(n,\gamma)$ cross sections

Friday 31 May 2024 11:40 (20 minutes)

As far as nucleosynthesis or element formation is concerned, almost all the nuclei heavier than iron have been made in part by the slow neutron capture and the rapid neutron capture processes ( $\approx 50\%$  each), respectively known as the s- and r- processes [1].

The neutron capture reactions  $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$  and  $^{193}\text{Ir}(n,\gamma)^{194}\text{Ir}$  are indirectly studied by analysing data obtained from the Oslo Cyclotron Laboratory (OCL). These data will allow for the study of  $^{193,194}\text{Ir}$  isotopes, from the  $^{192}\text{Os}(\alpha,\gamma)$  and  $^{192}\text{Os}(\alpha,\text{d}\gamma)$  reactions, respectively. The  $^{193}\text{Ir}(n,\gamma)^{194}\text{Ir}$  cross sections which our measurement will constrain will provide a comparison to existing (n, $\gamma$ ) measurement data [2].

In addition, the  $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$  reaction is a branching point in the s-process making it very interesting, but it is challenging to measure the (n, $\gamma$ ) cross section directly since  $^{192}\text{Ir}$  is unstable. Therefore the OCL data may provide very valuable information on the  $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$  cross section by indirectly constraining it with the experimental nuclear level density (NLD) and  $\gamma$ -strength function ( $\gamma\text{SF}$ ).

An array of Sodium Iodine (NaI)Tl detectors, called CACTUS, detected  $\gamma$ -rays and the silicon particle telescope array, called SiRi, was used to detect charged particles in coincidence. The NLDs and  $\gamma\text{SF}$ s are being extracted below the neutron separation energy,  $S_n$ , using the Oslo Method [3]. Furthermore, the NLDs and  $\gamma\text{SF}$ s will be used as inputs in the open-source code called TALYS to calculate cross-sections of  $^{193,194}\text{Ir}$ . I will provide preliminary results of the measured NLDs and  $\gamma\text{SF}$ s from the  $^{192}\text{Os}(\alpha,\text{d}\gamma)^{194}\text{Ir}$  reaction which will be used as inputs in the code TALYS to calculate cross-sections of  $^{193,194}\text{Ir}$ .

[1] Arnould, M., Goriely, S., and Takahashi, K. (2007). Physics Reports, 450(4-6), 97-213.

[2] Zerkov, V. V., and Pritychenko, B. (2018). The experimental nuclear reaction data (EXFOR) 888, 31-43.

[3] Schiller, A., Bergholt, L., Guttormsen, M., Melby, E., Rekstad, J., and Siem, S. (2000). Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, 447(3), 498-511.

This work is based on the research supported by the National Research Foundation of South Africa (Grant Number:PMDS22070734847) and SAINTS Prestigious Doctoral Scholarship.

**Author:** MAGAGULA, Sebenzile (University of the Witwatersrand and iThemba LABS)

**Co-authors:** LARSEN, Ann-Cecilie (University of Oslo (NO)); WIEDEKING, Mathis (Lawrence Berkeley National Laboratory)

**Presenter:** MAGAGULA, Sebenzile (University of the Witwatersrand and iThemba LABS)