



Contribution ID: 58

Type: **Non-Invited Talk**

Neutrino-Anti-neutrino beaming across the Earth to test CPT asymmetry

Tuesday 27 November 2018 14:45 (25 minutes)

Here we considered how the longest baseline neutrino oscillation available, crossing most of Earth's diameter, may improve the measurement and at best disentangle any hypothetical CPT violation occurring between neutrino-antineutrino, while testing τ neutrino and even the appearance of anti tau one at the highest rate. The ν_μ and anti ν_μ disappearance correlated with the tau appearance is considered for those events at the largest distances. We thus propose a beam of ν_μ and anti ν_μ crossing through the Earth, within an OPERA-like experiment from CERN (or Fermilab), beaming in the direction of the IceCube-DeepCore or future Pingu detector at the South Pole. The similar test may be done with future Hyperkamiokande in Japan and in Korea, The ideal energy lies at 21 GeV to test the disappearance or (for any tiny CPT violation) the partial anti ν_μ appearance. Such a tuned detection experiment may lead to a strong signature of τ or anti tau generation even within its neutral current noise background events. The tau appearance signal is above (or within) 10σ a year, even for one year a 1% OPERA-like experiment. Peculiar configurations for θ_{13} and the hierarchy neutrino mass test may also be better addressed by a DeepCore-PINGU array detector beaming ν_μ and observing ν_e at 6 GeV neutrino energy windows.

Content of the contribution

Both

Author: FARGION, Daniele (Rome University 1 Sapienza and INFN)

Presenter: FARGION, Daniele (Rome University 1 Sapienza and INFN)

Session Classification: Neutrino masses, mixing and discrete symmetries

Track Classification: [4] Neutrino masses, mixing and discrete symmetries