

# Heavy Neutral Leptons at Belle II?

Michele Tammaro

'Anomalies and Precision in the Belle II Era', 07/09/2021

Preliminary work with

S. Dreyer, T. Ferber & S. Longo (DESY), J. Zupan & P. Ilten (U. Cincinnati)



# The search for Long Lived Particles (LLP)




Figure from Codex-b EOI:1911.00481

# The search for Long Lived Particles (LLP)

A. Filimonova's talk





Figure from Codex-b EOI:1911.00481

# Heavy Neutral Leptons

Massive neutrinos + Standard Model  $\longrightarrow$  New Particles [Ma ('98)]

Heavy Neutral Leptons (HNL) aka Sterile Neutrinos

$$\mathcal{L}_{int} = \sum_{\alpha, I} c_{\alpha, I} (\bar{L}_{\alpha} H) N'_I \xrightarrow{\text{EWSB}} \nu'_{\alpha} \simeq \nu_{\alpha} + U_{\alpha} N$$



# Heavy Neutral Leptons

Massive neutrinos + Standard Model  $\longrightarrow$  New Particles [Ma ('98)]

For practical purposes, focus on one flavoured HNL


$$\mathcal{L}_{int} = c_\alpha (\bar{L}_\alpha H) N' \xrightarrow[\alpha = e, \mu, \tau]{\text{EWSB}} \nu'_\alpha \simeq \nu_\alpha + U_\alpha N$$

## PRODUCTION


[K. Bondarenko et al:1805.08567]

$$m_N < m_\tau$$

### B meson decay



### Tau lepton decay



# Heavy Neutral Leptons

Massive neutrinos + Standard Model  $\longrightarrow$  New Particles [Ma ('98)]


For practical purposes, focus on one flavoured HNL

$$\mathcal{L}_{int} = c_\alpha (\bar{L}_\alpha H) N' \xrightarrow[\alpha = e, \mu, \tau]{\text{EWSB}} \nu'_\alpha \simeq \nu_\alpha + U_\alpha N$$


DECAY

[K. Bondarenko et al:1805.08567]

Charged current




Neutral current




# Heavy Neutral Leptons

Muon-flavoured case



# Heavy Neutral Leptons

Tau-flavoured case, mostly unexplored



Can we explore more space with B mesons at Belle II?

# Simulating decays

Implementation of heavy meson decays

Pythia

(done but not embedded yet)



MadGraph

(no  $B \rightarrow D^*$  yet)

Simple but helpful codes for  
BR and widths  
(Python and Mathematica)



<https://gitlab.com/hnls>



# Simulating decays

Implementation of heavy meson decays

Pythia

(done but not embedded yet)



MadGraph

(no  $B \rightarrow D^*$  yet)

Simple but helpful codes for  
BR and widths  
(Python and Mathematica)



<https://gitlab.com/hnls>



# Simulating decays

Implementation of heavy meson decays

Pythia

(done but not embedded yet)



MadGraph

(no  $B \rightarrow D^*$  yet)

Simple but helpful codes for BR and widths  
(Python and Mathematica)



<https://gitlab.com/hnls>



# Expected Yield

50 ab<sup>-1</sup>

Require charged final states

$$N_N = N_{B\bar{B}} \times \mathcal{B}(B \rightarrow NX) \times \mathcal{B}(N \rightarrow \nu_\tau f^+ f^-) \times \mathcal{A}$$


N decays in tracking volume

$$0.5 < r_T < 80 \text{ cm}$$


$$0.3 < \theta_{lab} < 2.6 \text{ rad}$$

Cut on final state momenta

$$p_f > 0.1 \text{ GeV}$$



# Expected Yield



Still ideal case

# Take home message

- User-friendly simulations frameworks for meson decays are (almost) ready



<https://gitlab.com/hnls>

- Searches for HNLs at Belle 2 are still limited by statistics