Contribution ID: 53

Dark Energy Instabilities induced by Gravitational Waves

Monday 11 November 2019 16:10 (15 minutes)

In this talk I will discuss the decay of gravitational waves (GW) into dark energy fluctuations π in the context of the EFT of Dark Energy.

In such theories, the time-dependence of the Dark Energy (DE) field spontaneously breaks Lorentz invariance. Therefore as for light in a material, GW travelling in the cosmic medium are affected by dispersion phenomena and can decay into DE fluctuations. For cubic Horndeski and beyond Horndeski theories, the gravitational wave acts as a classical background for π and thus modifies its dynamics. In particular, for a sufficiently large amplitude of the wave, the kinetic term of π becomes pathological, featuring gradient and ghost instabilities. For smaller gravitational wave amplitude, π fluctuations are described by a Mathieu equation and feature instability bands that grow exponentially. The gravitational wave signal is affected by the π back-reaction and this provides very stringent bounds on cubic and quartic GLPV theories.

Author: TAMBALO, Giovanni (SISSA)

Co-authors: VERNIZZI, Filippo (CEA/IRFU,Centre d'etude de Saclay Gif-sur-Yvette (FR)); CREMINELLI, Paolo (Scuola Normale Superiore (SNS)); YINGCHAROENRAT, Vicharit (SISSA)

Presenter: TAMBALO, Giovanni (SISSA)

Session Classification: Short talks