Contribution ID: 58

Type: Lightning talk

Extreme Dark Matter Tests with Extreme Mass Ratio Inspirals

Tuesday 12 November 2019 12:15 (5 minutes)

Future space-based laser interferometry experiments such as LISA are expected to detect O(100 - 1000) stellar-mass compact objects (e.g., black holes, neutron stars) falling into massive black holes in the centers of galaxies, the so-called extreme-mass-ratio inspirals (EMRIs). If dark matter forms a "spike" due to the growth of the massive black hole, it will induce a gravitational drag on the inspiraling object, changing its orbit and gravitational-wave signal. We show that detection of even a single dark matter spike from the EMRIs will severely constrain several popular dark matter candidates, such as ultralight bosons, keV fermions, MeV-TeV self-annihilating dark matter, and sub-solar mass primordial black holes, as these candidates would flatten the spikes through various mechanisms. Future space gravitational wave experiments could thus have a significant impact on the particle identification of dark matter.

Authors: Dr HANNUKSELA, Otto (Nikhef); Dr NG, Kenny (Gravitation Astroparticle Physics Amsterdam (GRAPPA)); Prof. LI, Tjonnie (The Chinese University of Hong Kong)

Presenter: Dr HANNUKSELA, Otto (Nikhef)

Session Classification: Lightning talks