

# Primordial GW in Modified Gravity & Cosmology

**Anish Ghoshal**


Laboratorie Nazionale Frascati - INFN  
University Roma Tre

*anishghoshal1@gmail.com*

November, 2019 Amsterdam

Based on -  
arXiv: 1905.10410  
arXiv: 1911.xxxxx

# History of the Universe



## The idea: Naively

Propagation of Primordial GW generated during Inflation:

$$\ddot{h}_{ij} + 3H \dot{h}_{ij} + \frac{k^2}{a^2} h_{ij} = 16\pi G \Pi_{ij}^{TT}, \quad (1)$$

Solution:


$$h_{ij}(t, \vec{x}) = \sum_P \int \frac{d^3k}{(2\pi)^3} h^P(t, \vec{k}) \epsilon_{ij}^P(\vec{k}) e^{i\vec{k}\cdot\vec{x}}, \quad (2)$$

$$h_{\vec{k}}^P = h_{\vec{k},0}^P U(t, k), \quad (3)$$

$$\Pi_{ij} = \frac{T_{ij} - p g_{ij}}{a^2} \quad (4)$$

$$\Omega_{GW}(\eta, k) = \frac{1}{12 a^2(\eta) H^2(\eta)} \mathcal{P}_T(k) [U'(\eta, k)]^2 \quad (5)$$

## On the GW sensitivity Map




## Non-standard Cosmology

- History of the Universe before BBN is unknown.
- Non-standard cosmology predicts scalar field  $\phi$  and its energy density dominates in the Early Universe.
- Its Equation-of-state  $\omega_\phi$ .
- Modifies the Hubble expansion:  $H^2 \propto \rho_\phi$ .
- Parameter  $\xi = \frac{\rho_\phi}{\rho_R}$ .

# On the GW sensitivity Map


Radiation  
domination like  
scenario





# On the GW sensitivity Map

Modulus or matter  
domination like  
scenario


$$\omega_\phi = 0, \xi = 10^{-11}$$



$$T_{\text{dec}} = 10 \text{ MeV}$$



# On the GW sensitivity Map



## Modified Gravity

- Modify cosmological expansion (motivated from modified gravity theories)

$$H_{MC}(T) = A(T)H_{GR}(T), \quad (6)$$

- Strategy:

$$A(T) = 1 + \eta \left( \frac{T}{T_*} \right)^\nu, \quad (7)$$

where  $T_*$  is a parameter with dimensions of the temperature, and  $\{\eta, \nu\}$  are free parameters.

$$A(T) = \begin{cases} 1 + \eta \left( \frac{T}{T_*} \right)^\nu \tanh \frac{T - T_{re}}{T_{re}} & \text{for } T > T_{BBN} \\ 1 & \text{for } T \leq T_{BBN} \end{cases} \quad (8)$$




- $\nu$  labels cosmological models:

1.  $\nu = 2$  in Randall-Sundrum type II brane cosmology,  $\nu = 1$  in kination models,
2.  $\nu = 0$  in cosmologies with an overall boost of the Hubble expansion rate like in the case of a large number of additional relativistic degrees of freedom in the thermal plasma
3.  $\nu = -1$  in scalar-tensor cosmology,
4.  $\nu = 2/n - 2$  in  $f(x)$  cosmology, with  $f(x) = x + \alpha x^n$ , with  $x = R, \mathcal{T}$  where  $R$  and  $\mathcal{T}$  stand for the scalar curvature and the scalar torsion, respectively.







# On the GW sensitivity Map

Some Preliminary results only -

$\eta = 1, \nu = 0, T = 1 \text{ GeV}$



$\eta = 1, \nu = 1, T = 1 \text{ GeV}$      $\eta = 100, \nu = 1, T = 1 \text{ k GeV}$






CONCLUSION:

*Probe of Modified Cosmological Expansion History in  
Early Universe !*







# On the GW sensitivity Map

Some Preliminary results only -

$\eta = 1, \nu = 0, T = 1 \text{ GeV}$



$\eta = 1, \nu = 1, T = 1 \text{ GeV}$      $\eta = 100, \nu = 1, T = 1 \text{ k GeV}$



Thank You