


**Exploring dense QCD
through hamiltonian lattice simulations
in (1+1) dimensions**

**Yoshimasa Hidaka
(YITP)**

Based on

Hayata, YH, Nishimura, JHEP 07 (2024) 106, arXiv:2311.11643

QCD at finite density



- What is the equation of state for QCD at finite density?
- How does the quark distribution function change from baryonic matter to quark matter?
- What kind of phase is realized?
An inhomogeneous phase?

QCD₂

We study QCD₂ at density using density matrix renormalization group (DMRG) technique.

QCD₂

We study QCD₂ at density using density matrix renormalization group (DMRG) technique.

Properties of (1+1) dimensions

- Gauge fields are nondynamical
- Hilbert space is finite dimensional in Open Boundary Condition(OBC)

(dimensionless) QCD₂ Hamiltonian

$$J = \frac{ag_0}{2}, w = \frac{1}{2g_0a}, m = m_0/g_0 \quad \text{We use } g_0 = 1 \text{ unit}$$

$$H/g_0 = J \sum_{n=1}^{N-1} E_i^2(n) \quad \text{Electric field term}$$

$$+ w \sum_{n=1}^{N-1} \left(\chi^\dagger(n+1)U(n)\chi(n) + \chi^\dagger(n)U^\dagger(n)\chi(n+1) \right)$$

Hopping term

$$+ m \sum_{n=1}^N (-1)^n \chi^\dagger(n)\chi(n) \quad \text{Mass term}$$

Elimination of Link variables U

Sala, Shi, Kühn, Bañuls, Demler, Cirac, Phys. Rev. D 98, 034505 (2018)

Atas, Zhang, Lewis, Jahanpour, Haase, Muschik, Nature Commun. 12, 6499 (2021)

$$\Theta\chi(n)\Theta^\dagger := U(n-1)U(n-2)\cdots U(1)\chi(n)$$

Elimination of Link variables U

Sala, Shi, Kühn, Bañuls, Demler, Cirac, Phys. Rev. D 98, 034505 (2018)

Atas, Zhang, Lewis, Jahanpour, Haase, Muschik, Nature Commun. 12, 6499 (2021)

$$\Theta\chi(n)\Theta^\dagger := U(n-1)U(n-2)\cdots U(1)\chi(n)$$

$$\Theta H \Theta^\dagger = J \sum_{n=1}^{N-1} \left(\sum_{m=1}^n \chi^\dagger(m) T_i \chi(m) \right)^2 \quad \text{Electric fields term}$$

$$+ w \sum_{n=1}^{N-1} \left(\chi^\dagger(n+1)\chi(n) + \chi^\dagger(n)\chi(n+1) \right)$$

Hopping term

$$+ m \sum_{n=1}^N (-1)^n \chi^\dagger(n)\chi(n) \quad \text{mass term}$$

As a variational ansatz of wave function

- We employ a matrix product state

$$|\psi\rangle = \sum_{\{n_i\}} |n_1\rangle \cdots |n_N\rangle \text{tr} M_1^{n_1} \cdots M_N^{n_N}$$

$$[M_i^{n_i}]_{ij} : D \times D \text{ matrix}$$

- Optimize the wave function by density matrix renormalization group technique

$$E = \min_{\psi} \langle \psi | H | \psi \rangle$$

We employ iTensor



Numerical results

SU(2) QCD with $N_f = 1$

Color SU(2), 1 flavor, vacuum

single baryon state $\dim \mathcal{H} = 2^{300}$ $J = 1/20$ $w = 5$ volume $V = 15$

Baryon number density Quark distribution function




Baryon size ~ 1


Color SU(2), 1 flavor, vacuum

$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$

Pressure




Energy density




Color SU(2), 1 flavor, vacuum

$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$

Chemical potential




Sound velocity



Inhomogeneous phase (density wave)


$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$



Wave number dependence

$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$


Wave number dependence



Wave number dependence


$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$

Wave number dependence



Hadronic picture


If hadron interactions are repulsive



Wave number dependence


$$J = 1/8 \quad w = 2 \quad V = 40 \quad \dim \mathcal{H} = 2^{320}$$

Wave number dependence



Hadronic picture

If hadron interactions are repulsive




Quark picture

If interactions between quarks
Fermi surface is unstable

\Rightarrow density wave $k = 2p_F = 2\pi n_B$

Quark distribution function

$$J = 1/8 \quad w = 2 \quad V = 60 \quad \dim \mathcal{H} = 2^{480}$$



- **Low density**
No Fermi sea
- **High density**
Fermi-sea
+BCS like pairing
(density wave)


baryon quark transition around $n_B \sim 0.2$

SU(3) QCD with $N_f = 1$


Color SU(3), 1 flavor

$$J = 1/8 \quad w = 2 \quad V = 12 \quad \dim \mathcal{H} = 2^{144}$$

Pressure




Energy density




Color SU(3), 1 flavor, vacuum

$$J = 1/8 \quad w = 2 \quad V = 12 \quad \dim \mathcal{H} = 2^{144}$$

Chemical potential




Sound velocity




Color SU(3), 1 flavor

$$J = 1/8 \quad w = 2 \quad V = 12 \quad \dim \mathcal{H} = 2^{144}$$


density wave



Wave number dependence



Quark distribution



Baryon quark transition around $n_B = 0.3$?

Summary

- We study QCD_2 at finite density with one flavor for two and three colors.
- We employ Hamiltonian formalism and density matrix renormalization group techniques
- We find inhomogeneous phases both two and three colors.