XVIth Quark Confinement and the Hadron Spectrum



Contribution ID: 251

Type: Oral

Lightcone and quasi distribution amplitudes for light octet and decuplet baryons

Thursday 22 August 2024 17:10 (20 minutes)

We present a comprehensive investigation of leading-twist lightcone distribution amplitudes (LCDAs) and quasi distribution amplitudes (quasi-DAs) for light octet and decuplet baryons within large momentum effective theory (LaMET). In LaMET, LCDAs can be factorized in terms of a hard kernel and quasi-DAs that are defined as spatial correlators and calculable on Lattice QCD. To renormalize quasi-DAs and eliminate the singular terms \ln (left \mu^2 z_i^2\right)'s in them, which undermine the efficacy in perturbative expansion, we adopt a hybrid renormalization scheme that combines the self-renormalization and ratio scheme. Through self-renormalization, we eliminate UV divergences and linear divergences at large spatial separations in quasi distribution amplitudes without introducing additional nonperturbative effects. By taking a ratio with respect to the zero-momentum matrix element, we effectively remove UV divergences at small spatial separations. Under the hybrid renormalization scheme, we calculate the hard kernels up to one-loop accuracy. It turns out that only four different hard kernels are needed for all leading-twist LCDAs of octet and decuplet baryons. These results are crucial for the lattice-based exploration of the LCDAs of a light baryon from the first principle of QCD.

Author: Dr ZENG, Jun (Shanghai Jiao Tong University)Presenter: Dr ZENG, Jun (Shanghai Jiao Tong University)Session Classification: Light Quarks

Track Classification: B: Light Quarks