

Session B – Round Table (Wednesday, 21 August 2024 at 12:00-13:00):

Precision QCD: What we know what we want to know

Chair: Jianwei Qiu

Panel Members : Ian Cloët - Zein-Eddine Meziani - Wei Wang

Description in the Conference Program:

In recent years, rapid progress has been made on the determination of PDFs and others from Lattice QCD. Very promising results are obtained while there are still some open questions that need to be discussed. New perspectives are also required, and we hope through this round table discussion, new thoughts can be inspired.

Topics and Questions by the Panel:

- **Matching the partons to hadrons (Jianwei Qiu)**
- **Transverse momentum dependent PDFs (Wei Wang)**
- **Intrinsic properties of nucleons and nuclei (Zein-Eddine Meziani)**
- **Nuclear PDFs and Imaging ($A>1$) (Ian Cloët)**

Additional Topics, Questions and Thoughts from the Audience?

Matching the partons to hadrons:

□ What we know:

- Owing to the Confinement, NO quarks/gluons (the partons) can be seen in isolation in Detectors
- Any physical observable with identified hadron is NOT perturbatively calculable
- QCD factorization is an approximation – valid for many physical observables at the leading power

$$\sigma_{\{h_i\}}^{\text{phy}}(\{Q_i\}, 1/R_h \sim \Lambda_{\text{QCD}}) = \sum_{\{f_i\}} \hat{H}_{\{f_i\}}(\{x_i\}, \{Q_i\}, \mu) \prod_i \otimes \phi_{f_i/h_i}(x_i, \mu, 1/R_h) + \mathcal{O}(1/R_h Q)^n$$

Physical cross section

Calculable partonic “Probe”



Matching Parton to Hadron

Approximation “controllable?”

- QCD factorization works:

MSHT20

Data set	N_{pts}	NLO χ^2/N_{pts}	NNLO χ^2/N_{pts}
ATLAS 8 TeV s. diff $t\bar{t}$	25	1.56	0.98
CMS 8 TeV d. diff $t\bar{t}$	15	2.19	1.50
ATLAS 7 TeV W, Z	61	5.00	1.91
ATLAS 8 TeV W	22	3.85	2.61
ATLAS 8 TeV d. diff Z	59	2.67	1.45
ATLAS 8 TeV Z p_T	104	2.26	1.81
ATLAS 8 TeV W + jets	39	1.13	0.60
Total LHC data	1328	1.79	1.33
Total non-LHC data	3035	1.13	1.10
Total	4363	1.33	1.17




EIC's help on unpol. PDFs

Matching the partons to hadrons:

□ What we want to know:

- Hadron's 3D partonic structure – confined parton distributions and motions ?
- How to extract the x-dependence of GPDs (moments for GFFs)?

Cédric Lorcé, DIS2024



without breaking the hadron!

(xi, t) fixed by p and p'

$$i\mathcal{M} \propto \int_{-1}^1 dx \frac{F(x, \xi, t)}{x - \xi + i\epsilon} \equiv "F_0(\xi, t)"$$


GPDs + Shadow GPDs

$$\int_{-1}^1 \frac{dx S(x, \xi)}{x - \xi \pm i\epsilon} = 0$$

Need new processes:

$$i\mathcal{M} \propto \int_{-1}^1 dx \frac{F(x, \xi, t)}{x - x_p(\xi, q) + i\epsilon}$$

Have **x** entangled to the measured **q**!
DDVCS, SDHEP, ..., need more !



[Belitsky, Ji, Yuan, PRD69 (2004) 074014]
 [Meissner, Metz, Schlegel, JHEP08 (2009) 056]
 [CL, Pasquini, Vanderhaeghen, JHEP05 (2011) 041]
 [CL, Pasquini, JHEP09 (2013) 138]

→ $\Delta = p' - p = 0$
 → $\int dx$
 → $\int d^2 k_\perp$

Matching the partons to hadrons – the role of LQCD:

□ What we know:

See also Huey-Wen's talk

- LQCD can NOT calculate parton correlation functions (t-dependence of parton correlators) directly ?
- Ji's wonderful idea: to calculate the equal-time version – quasi-PDFs and let $P_z \rightarrow \infty$ to get PDFs
- The quasi-PDFs are NOT physical, need renormalization → pseudo-PDFs, ...
- Equal-time quark/gluon correlators are proved to be **multiplicative renormalizable** !
- Renormalized hadron matrix elements of the equal-time quark/gluon correlators **can be factorized** !

$$\langle h(P) | \mathcal{O}_{q,g}^R(z, \mu_R) | h(P) \rangle = \sum_f \int_{-1}^1 \frac{dx}{x} \hat{K}_f(xP \cdot z, z^2, \mu_R, \mu_F) \phi_{f/h}(x, \mu_F) + \mathcal{O}(z^2)^n$$



- Matching \hat{K}_f to NNLO! Quasi-PDFs: $\tilde{\phi}_{(q,g)/h}(\tilde{x}, \mu_R) \propto \int_{-\infty}^{\infty} \frac{d(P \cdot z)}{2\pi} e^{i\tilde{x}P \cdot z} \langle h(P) | \mathcal{O}_{q,g}^R(z, \mu_R) | h(P) \rangle$
- LQCD data in Global fits: more impactful for polarized PDFs, transversity, GPDs, ..., not yet for PDFs

□ What we want to know:


- Systematic uncertainties from the continue limit, connected diagrams, renormalization, ... ?
- LQCD calculations to generate data that are good for the extraction of x-dependent TMDs, ... ?

Transverse Momentum Dependent PDFs

Leading Quark TMDPDFs  Nucleon Spin  Quark Spin


		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \text{Unpolarized}$		$h_1^\perp = \text{Boer-Mulders}$
	L		$g_1 = \text{Helicity}$	$h_{1L}^\perp = \text{Worm-gear}$
	T	$f_{1T}^\perp = \text{Sivers}$	$g_{1T}^\perp = \text{Worm-gear}$	$h_1 = \text{Transversity}$ $h_{1T}^\perp = \text{Pretzelosity}$

$$F(x, b; \mu, Q^2) = \left(\frac{Q^2}{\zeta_\mu(b)} \right)^{-\mathcal{D}(b, \mu)} F(x, b)$$




Moos, Scimemi, Vladimirov, Zurita, 2305.07473

- ✓ A lot of data on SIDIS and DY are available.
- ✓ In addition to perturbative constraints, several global analyses of TMDPDFs are obtained.
- ✓ Soft function (intrinsic and CS kernel) has been calculated from several lattice groups (D.Lin's talk on 19/08).
- ✓ A calculation of quark unpolarized isovector TMDPDFs from lattice QCD ($a=0.12\text{fm}$) is available.




LPC, 2211.02340


Transverse Momentum Dependent PDFs



EIC Yellow Report: 2103.05419



- ✓ With more reliable data, to what extent can the global analysis obtain?
- ✓ Can one obtain sophisticated results on TMDPDFs from lattice QCD in a few years?
- ✓ Gluon TMDPDFs (+small x) and subleading twist?
- ✓ TMD fragmentation functions?



Zeng, Liu, Sun, Zhao, 2208.14620

Intrinsic properties of nucleons and nuclei

□ What do we know?



- Total mass, Total charge, Total magnetic moment, Total Spin
- Electromagnetic form factors (quarks driven)
 - Current and charge distribution of proton and neutron and nuclei

□ What do we want to know?

$$\langle N' | T_{q,g}^{\mu,\nu} | N \rangle = \bar{u}(N') \left(A_{g,q}(t) \gamma^{\{\mu P \nu\}} + B_{g,q}(t) \frac{i P^{\{\mu \sigma \nu\}} \rho \Delta_\rho}{2M} + C_{g,q}(t) \frac{\Delta^\mu \Delta^\nu - g^{\mu\nu} \Delta^2}{M} + \bar{C}_{g,q}(t) M g^{\mu\nu} \right) u(N)$$

- Energy-momentum tensor (gravitational) form factors of quarks and gluons
 - Mechanical properties of the proton/neutron **and nuclei**
 - Mass density distribution of gluons and quarks
 - Scalar energy density distribution of gluons
 - Pressure density distribution for quarks and gluons
 - **If the gluon scalar radius of the nucleon is larger than the charge radius, what is the consequence in nuclei?**

Example gluonic GFFs today & tomorrow: Can we improve on the accuracy?




B.Duran, et al., proton, *Nature* **615**, no.7954, 813-816 (2023)
 K. A. Mamo and I. Zahed, *Phys. Rev. D* **106**, no.8, 086004 (2022)
 D. A. Pefkou, D. C. Hackett and P. E. Shanahan, *Phys. Rev. D* **105** (2022) no.5, 054509

Different Mass decompositions

Proton Mass budget decompositions C. Lorcé (from 2022 INT workshop)


Trace decomposition

$\mu = 2 \text{ GeV}$




Relies on
virial theorem

Energy decomposition



Independent of
virial theorem

Ji's decomposition




Motivated by
virial theorem

The many fronts of experimental studies


Since 1998
Generalized Parton Distributions
(GPDs)

Exclusive reactions
In nucleons and
nuclei

Elastic form factors
Deep Virtual Compton
Scattering
Deep Virtual Meson
Production





Inclusive
Parton distributions
Sum rules and polarizabilities



Since 2002
Transverse Momentum Distributions
(TMDs)

Semi-Inclusive DIS
In nucleons and nuclei

Distributions and
Fragmentation functions





Electroweak
probe to hadronic systems

NUCLEAR PDFS & IMAGING

- Little is known about Nuclear PDFs
 - EMC Effect provides an enduring puzzle at the intersection of QCD and traditional nuclear structure approaches
- To gain deeper insight into QCD and Nuclei need to address several interesting questions, for example:
 - What is the quark flavor dependence of nuclear PDFs/EMC effect?
 - How are gluon PDFs modified by the nuclear medium?
 - Observation of the gluon transversely PDF in $J \geq 1$ nuclei (e.g. ${}^2\text{H}$ or ${}^6\text{Li}$) first direct evidence for non-nucleonic components in nuclei
 - Can the new PDFs that appear in $J \geq 1$ nuclei, such as $b_1(x)$, also provide inside into QCD effects in the NN potential?
- As a first step, important to obtain data on polarized nuclear PDFs (e.g. in ${}^6\text{Li}$, ${}^7\text{Li}$, and other light nuclei)
 - Will address the question: How is quark spin impacted by the nuclear medium?
 - For example, is quark spin converted to quark orbital angular momentum by the nuclear medium?


EMC Effect observed in unpolarized DIS




MOMENTUM TOMOGRAPHY

Ninomiya, Bentz and ICC, Phys. Rev. C **96**, no.4, 045206 (2017)

leading twist		quark operator		
		γ^+	$\gamma^+\gamma_5$	$\gamma^+\gamma^i\gamma_5$
target polarization	U	$f_1 = \text{unpolarized}$		$h_1^\perp = \text{Boer-Mulders}$
	L		$g_1 = \text{helicity}$	$h_{1L}^\perp = \text{worm gear 1}$
	T	$f_{1T}^\perp = \text{Sivers}$	$g_{1T} = \text{worm gear 2}$	$h_1 = \text{transversity}$ $h_{1T}^\perp = \text{pretzelosity}$
	ROSENFELDER	$\left. \begin{matrix} \theta_{LL}(x, \mathbf{k}_T^2) \\ \theta_{TT}(x, \mathbf{k}_T^2) \\ \theta_{LT}(x, \mathbf{k}_T^2) \end{matrix} \right\}$	$g_{1TT}(x, \mathbf{k}_T^2)$ $g_{1LT}(x, \mathbf{k}_T^2)$	$h_{1LL}^\perp(x, \mathbf{k}_T^2)$ h_{1TT}, h_{1TT}^\perp h_{1LT}, h_{1LT}^\perp







- TMDs enable the study of the EMC Effect as a function of transverse momentum
- Tensor polarization in spin-1 target gives 3 additional T -even and 7 additional T -odd quark TMDs
- These new TMDs are related to $b_1(x)$ PDF
 - Impossible to explain HERMES data with only bound nucleon degrees of freedom — need exotic QCD states, 6q bags, etc



SPATIAL TOMOGRAPHY

Images represent naïve expectations for ${}^4\text{He}$

- Spatial Imaging of nuclei via GPDs can provide rich insights into QCD
- Key physics questions include:
 - How does the NN interaction arise from QCD?
 - How do quark/gluon confinement length scales change in medium?
- Insights can be obtained by comparing quark and gluon distributions for slices in x and b_T^2
- Can also study quark and gluon gravitational form factors; mass/energy radii for quarks and gluons, etc.
- Nuclear PDFs & Imaging are essential for comprehensive understanding of QCD



Backup Slides


Unified View of Nucleon Structure

$W_p^u(x, k_T, r_T)$ Wigner distributions

Transverse Momentum Dist. (TMD)

Tomography

Generalized Parton Dist. (GPD)





Probing deeper for the fundamental
degrees of freedom, quarks and gluons

Quarks

Emergent Phenomenon

