

# Leading Fermionic Three-Loop Corrections to EWPOs

RADCOR-LoopFest 2021

Lisong Chen and Ayres Freitas

ArXiv:2002.05845[hep-ph](JHEP) ArXiv:2012.08605[hep-ph](JHEP)





PITT-PACC





1. Precision Test of EWPOs
2. Renormalization
3. Computing Observables
4. Technique Aspects
5. Numerical Results
6. Summary and Outlook



# Precision Test of Electroweak Precision Observables (EWPOs)

- The Standard Model can only be tested by considering higher-order corrections when confronting experimental high precision data.
- New physics unknown by experiments directly might be sensitive to quantum corrections.






J.Erler '19

□ A way of checking the inner consistency of the SM.

e.g. Constraints of  $m_H - m_t$  by various set of EWPOs.



# EWPOs Introduction (exp)


$M_W$  Measured via W-boson pair production

$\Gamma_Z$  By fitting the cross section of  $e^+e^- \rightarrow f\bar{f}$   $\Gamma_Z = \sum_f \Gamma_{f\bar{f}}$

$$\sigma_{hard} = \sigma_Z + \sigma_\gamma + \sigma_{\gamma Z} + \sigma_{box}$$

$$\sigma_Z = \sigma_{f\bar{f}}^{peak} \frac{s\Gamma_Z^2}{(s - m_Z^2)^2 + s^2 \frac{\Gamma_Z^2}{m_Z^2}},$$

where  $\sigma_{f\bar{f}}^{peak} = \frac{1}{\mathcal{R}_{QED}} \sigma_{f\bar{f}}^0$ , and  $\sigma_{f\bar{f}}^0 = \frac{12\pi}{m_Z^2} \frac{\Gamma_{ee}\Gamma_{f\bar{f}}}{\Gamma_Z^2}$




$\sigma[e^+e^- \rightarrow f\bar{f}] = \mathcal{R}_{QED}(s, s') \otimes \sigma_{hard}(s')$   
 $e^+e^- \rightarrow f\bar{f}$  at Z resonance peak.

$\sin^2 \theta_{eff}^l$  Extracted from the measured asymmetries, which are defined based on

$$A_{FB}^f = \frac{\sigma_f(\theta < \frac{\pi}{2}) - \sigma_f(\theta > \frac{\pi}{2})}{\sigma_f(\theta < \frac{\pi}{2}) + \sigma_f(\theta > \frac{\pi}{2})} \equiv \frac{3}{4} A_e A_f,$$

$$A_{LR}^f = \frac{\sigma_f(P_e < 0) - \sigma_f(P_e > 0)}{\sigma_f(P_e < 0) + \sigma_f(P_e > 0)} \equiv A_e |P_e|$$

$$A_f = 2 \frac{g_{Vf}/g_{Af}}{1 + (g_{Vf}/g_{Af})^2} = \frac{1 - 4|Q_f| \sin^2 \theta_{eff}^f}{1 - 4|Q_f| \sin^2 \theta_{eff}^f + 8(|Q_f| \sin^2 \theta_{eff}^f)^2} \quad (f = l, b, \dots)$$



## • How Far Have We Got?

- $M_W$
- **mixed QCD/EW 2-loop corrections** ✓. Djouai, Verzegnassi'87;Djouadi'88; Kniehl,Kühn, Stuar'99;Kniehl,Sirlin'93;Djouadi,Gambino'94
  - **complete EW 2-loop corrections** ✓. Freitas, Hollik, Walter, Weiglein'00;Awramik,Czakon '02;Onishchenko,Vertin '02
  - **improvements by 3-loop and 4-loop**  $\Delta\rho$  ✓. Avdeev et al.'94; Chetyrkin, Kühn, Steinhauser '95; v.d.Bij et al. '05; Schröder, Steinhauser '06; Faisst et al. '03; Boughezal, Tausk, v.d.Bij '05

$$\implies \Delta M_W \sim 4 \text{ MeV}$$

- $\Gamma_Z$
- **complete EW 1-loop and fermionic 2-loop** ✓. Freitas'13'14
  - **mixed QCD/EW 2-loop corrections** ✓. Djouai, Verzegnassi'87;Halzen Kniehl'91; Djouadi,Gambino'94; Chetyrkin, Kühn'96; Fleischer et al. '92
  - **improvements by 3-loop and 4-loop**  $\Delta\rho$  ✓. Avdee et al. '94'; v.d.Bij et al. '05;Schröder, Steinhauser'06;Faisst et al.'03;Boughezal, Tausk, v.d.Bij '05
  - **EW complete 2-loop corrections** . ✓.  $\mathcal{O}(\alpha_{bos}^2)$  Dubovyk, Freitas,Gluza, Riemann Usovitsch. '18




$\sin^2 \theta_{eff}^l$

$$\implies \Delta \Gamma_Z \sim 0.5 \text{ MeV}$$

- **mixed QCD/EW 2-loop and 3-loop**  $\Delta\rho$  **Corrections as for**  $M_W$
- **EW complete 2-loop corrections** ✓. Awramik, Czakon, Freitas, Weiglein '04; Hollik,Meier, Uccirati'05;Awramik,Czakon, Freitas '06

$$\implies \sin^2 \theta_{eff}^l \sim 4.5 \times 10^{-5}$$

- Current Status of Experimental Measurements



- Halcyon Time ?


- The SM shows good consistency by comparing measured EWPOs and theory predictions.
- The theoretical uncertainties is under well-control comparing to the known measurements. But...

	Current Theory	Main source	CEPC Exp	FCC-ee Exp	ILC Exp
$M_W$ [MeV]	4	$\alpha^3, \alpha^2\alpha_s$	1	1	2.5 – 5
$\Gamma_Z$ [MeV]	0.5	$\alpha^3, \alpha^2\alpha_s, \alpha\alpha_s^2$	0.5	0.1	0.8
$\sin^2 \theta_{eff}^l$	$4.3 \times 10^{-5}$	$\alpha^3, \alpha^2\alpha_s$	$2.3 \times 10^{-5}$	$0.6 \times 10^{-5}$	$10^{-5}$

- ❑ Due to the lack of knowledge of theory error estimation, we need  $|\Delta^{th}| \ll |\Delta^{obs}|$
- ❑ Current theoretical predictions are inadequate.
- ❑ The calculation of the next perturbative order  $\mathcal{O}(\alpha^3, \alpha^2\alpha_s)$  for the EWPOs will be necessary!!

### Why Leading Fermionic Corrections?

- ❑ Enhancement by power of Top Mass.
- ❑ Enhancement by power of flavor numbers  $N_f$



Considerably the leading numerical contribution!

# Renormalization

- ❖ Two schemes are considered.
  - ❑ On-Shell(OS) with **complex pole mass**  $\mathcal{O}(\alpha^3, \alpha^2\alpha_s)$
  - ❑ OS+  $\overline{MS}$  for top mass  $\mathcal{O}(\alpha^2\alpha_s)$
- Complex pole mass is a must for gauge-invariance.
- OS top mass closely connects to experiments, while suffers from renormalon issue and non-perturbative QCD.  $\overline{MS}$  top mass is preferable from theory point of view.
- Top masses calculated from two schemes related by a finite transformation.
- No asymptotic massive gauge boson, hence field renormalization of Z,W can be neglected.

**complex-pole**

$$s_0 \equiv \overline{M^2} - i\overline{M}\overline{\Gamma}$$

**The inverse dressed propagator (W/Z/H)**

$$D(p^2) = p^2 - \overline{M^2} - \delta Z(p^2 - \overline{M^2}) + \Sigma(p^2) - \delta\overline{M^2}$$

**yield mass counter term and widths**

$$\delta\overline{M^2} = \frac{\text{Re}\Sigma(\overline{M^2} - i\overline{M}\overline{\Gamma})}{Z} \quad \overline{\Gamma} = \frac{\text{Im}\Sigma(\overline{M^2} - i\overline{M}\overline{\Gamma})}{Z\overline{M}}$$

**mass ratio between two schemes**

$$\frac{M^{OS}}{M^{\overline{MS}}} = 1 + \alpha_s C_F \frac{3 \log M^{OS^2}/\mu^2 - 4}{4\pi} + \mathcal{O}(\alpha_s^2)$$

**Ward-Identity yields**

$$Z_e = (\sqrt{Z_{\gamma\gamma}} + \frac{\sin \theta_W}{\cos \theta_W} \sqrt{Z_{Z\gamma}})^{-1}$$

**Weak-Mixing Angle**

$$s_W + \delta s_W = \sqrt{1 - \frac{\overline{M_W^2} + \delta\overline{M_W^2}}{\overline{M_Z^2} + \delta\overline{M_Z^2}}}$$



- ❖ the self-energy function  $\Sigma(p^2)$  composed by 1-PI at desired order. Only transverse part contributes, longitudinal part cancels against unphysical amplitude. (Slavnov-Taylor Identity)

- ❖  $\gamma - Z$  mixing is included. Mixing counterterms are defined at poles

$$\hat{\Sigma}_{\gamma Z}(0) = 0 \quad \Re \hat{\Sigma}_{\gamma Z}(\overline{M_Z}^2 - i\overline{M_Z}\overline{\Gamma_Z}) = 0$$

- ❖ Different Breit-Wigner forms
- ❖ In Experiment

$$\sigma \sim \frac{1}{(s-M^2)^2 + s^2\Gamma^2/M^2}$$

$$\overline{M} = M/\sqrt{1 + \Gamma^2/M^2}$$

In Theory

$$\sigma \sim \frac{1}{(s-\overline{M}^2)^2 + \overline{\Gamma}^2\overline{M}^2}$$

$$\overline{\Gamma} = \Gamma/\sqrt{1 + \Gamma^2/M^2}$$

$$\Sigma_{V_1 V_2(3)} = \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3}$$

$$\Sigma_{V_1 V_2(\alpha_s \alpha^2)} = \text{Diagram 4} + \text{Diagram 5} + \text{Diagram 6} + \text{Diagram 7} + \dots$$

$$\begin{aligned} Z_\mu \Sigma_Z Z_\mu &= \\ & \text{Diagram 8} \\ & + \text{Diagram 9} \\ & + \text{Diagram 10} \\ & + \dots \end{aligned}$$

$$\Sigma_Z(p^2) = \Sigma_{ZZ}(p^2) - \frac{[\hat{\Sigma}_{\gamma Z}(p^2)]^2}{p^2 + \hat{\Sigma}_{\gamma\gamma}(p^2)},$$

$$\hat{\Sigma}_{\gamma Z}(p^2) = \Sigma_{\gamma Z}(p^2) + \frac{1}{2}\delta Z^{Z\gamma}(p^2 - \overline{M_Z}^2 - \delta\overline{M_Z}^2) + \frac{1}{2}\delta Z^{\gamma Z}p^2,$$

$$\hat{\Sigma}_{\gamma\gamma}(p^2) = \Sigma_{\gamma\gamma}(p^2) + \frac{1}{4}(\delta Z^{Z\gamma})^2(p^2 - \overline{M_Z}^2 - \delta\overline{M_Z}^2).$$

- ❖ Charge renormalization needs a special care. We need  $\alpha$  around  $q^2 \sim M_Z^2$ , while it's defined at Thomson limit ( $q^2 \sim 0$ ).
- ❖ light-quark masses are inherently ill-defined in EW Lagrangian due to non-perturbative feature at the given mass scale  $q^2 \sim m_{u,d,s}^2$ .
- ❖ Alternative methods needs to apply to carry out the contribution given by light quarks. **Dispersion relation** is the one frequently use. Other possible ways: Lattice QCD or Bhabha scattering.

## Charge Counterterms

### Pure EW

$$\delta Z_e^{(3)} = \frac{5}{2} \delta Z_e^{(1)}$$

### Mixed EW-QCD


$$\delta Z_e^{(3)} = 3\delta Z_{e(\alpha)} \delta Z_{e(\alpha_s\alpha)}$$

\*See Stefan Dittmaier's talk for more detail about charge renormalization.

$$\delta Z_e = -\frac{1}{2} \delta Z_{\gamma\gamma} = \frac{1}{2} \Sigma'_{\gamma\gamma}(0) \quad \text{at one-loop level}$$

$$\Sigma'_{\gamma\gamma}(0) \equiv \Pi(0) = \sum_f \frac{\alpha N_c Q_f^2}{3\pi} \left( \frac{2}{4-D} - \gamma_E - \log \frac{m_f^2}{4\pi\mu^2} \right)$$

$$\hat{\Pi}(s = M_Z^2) = \Pi(0) - \Re \Pi(M_Z^2) = \underbrace{\Pi^{lf}(0) - \Pi^{lf}(M_Z^2)}_{\Delta\alpha = \Delta\alpha_{lep} + \Delta\alpha_{had}} + \hat{\Pi}^{top}(M_Z^2)$$



$$\Delta_{had} = -\frac{\alpha}{3\pi} s \int_{4m_\pi^2}^{\infty} ds' \frac{R_{\gamma\gamma}(s')}{s'(s'-s-i\epsilon)} \Big|_{s=M_Z^2}$$

$$R_{\gamma\gamma}(s') = \frac{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow \mu^+\mu^-)}$$

- non-perturbative quantity, apply to ALL order.
- Good precision  $\sim 0.0001$

❖ **Mass counterterms:** By assuming  $\Gamma_{W,Z}/M_{W,Z} \sim \mathcal{O}(\alpha)$ , the imaginary part contributes to counterterms.

❑ **Pure EW corrections at 3-loop order**

$$\begin{aligned} \delta \overline{M}_{Z(3)}^2 = & \text{Re } \Sigma_{ZZ(3)}(\overline{M}_Z^2) + [\text{Im } \Sigma_{ZZ(2)}(\overline{M}_Z^2)] [\text{Im } \Sigma'_{ZZ(1)}(\overline{M}_Z^2)] \\ & + [\text{Im } \Sigma_{ZZ(1)}(\overline{M}_Z^2)] \left\{ \text{Im } \Sigma'_{ZZ(2)}(\overline{M}_Z^2) - [\text{Im } \Sigma'_{ZZ(1)}(\overline{M}_Z^2)] [\text{Re } \Sigma'_{ZZ(1)}(\overline{M}_Z^2)] \right. \\ & \quad \left. - \frac{1}{2} [\text{Im } \Sigma_{ZZ(1)}(\overline{M}_Z^2)] [\text{Re } \Sigma''_{ZZ(1)}(\overline{M}_Z^2)] \right. \\ & \quad \left. - \frac{\text{Im } \Sigma_{\gamma Z(1)}(\overline{M}_Z^2)}{M_Z^2} [2 \text{Re } \Sigma'_{\gamma Z(1)}(\overline{M}_Z^2) + \delta Z_{(1)}^{\gamma Z} + \delta Z_{(1)}^{Z\gamma}] \right\} \\ & + \frac{\text{Im } \Sigma_{\gamma Z(1)}(\overline{M}_Z^2)}{M_Z^2} \left\{ 2 \text{Im } \Sigma_{\gamma Z(2)}(\overline{M}_Z^2) - \frac{\text{Im } \Sigma_{\gamma Z(1)}(\overline{M}_Z^2)}{M_Z^2} [\text{Im } \Sigma_{\gamma\gamma(1)}(\overline{M}_Z^2)] \right\} \\ & + \frac{1}{2} \overline{M}_Z^2 \delta Z_{(1)}^{\gamma Z} \delta Z_{(2)}^{\gamma Z}. \end{aligned}$$

**Mixed EW-QCD corrections**

$$\begin{aligned} \delta \overline{M}_{Z(\alpha_s \alpha^2)}^2 = & \text{Re } \Sigma_{ZZ(\alpha_s \alpha^2)}(\overline{M}_Z^2) + [\text{Im } \Sigma_{ZZ(\alpha_s \alpha)}(\overline{M}_Z^2)] [\text{Im } \Sigma'_{ZZ(\alpha)}(\overline{M}_Z^2)] \\ & + [\text{Im } \Sigma_{ZZ(\alpha)}(\overline{M}_Z^2)] [\text{Im } \Sigma'_{ZZ(\alpha_s \alpha)}(\overline{M}_Z^2)] \\ & + \frac{2}{M_Z^2} [\text{Im } \Sigma_{\gamma Z(\alpha_s \alpha)}(\overline{M}_Z^2)] [\text{Im } \Sigma_{\gamma Z(\alpha)}(\overline{M}_Z^2)] + \frac{1}{2} \overline{M}_Z^2 \delta Z_{(\alpha)}^{\gamma Z} \delta Z_{(\alpha_s \alpha)}^{\gamma Z}. \end{aligned}$$


❑ **Total width of Z-boson at 3-loop order (Pure EW)**

$$\begin{aligned} \overline{\Gamma}_Z = & \frac{1}{M_Z} \left\{ \text{Im } \Sigma_{Z(1)} + \text{Im } \Sigma_{Z(2)} - (\text{Im } \Sigma_{Z(1)}) (\text{Re } \Sigma'_{Z(1)}) \right. \\ & + \text{Im } \Sigma_{Z(3)} - (\text{Im } \Sigma_{Z(2)}) (\text{Re } \Sigma'_{Z(1)}) \\ & + (\text{Im } \Sigma_{Z(1)}) [(\text{Re } \Sigma'_{Z(1)})^2 - \text{Re } \Sigma'_{Z(2)} - \frac{1}{4} (\delta Z_{(1)}^{\gamma Z})^2 - \frac{1}{2} (\text{Im } \Sigma_{Z(1)}) (\text{Im } \Sigma''_{Z(1)})] \\ & + \text{Im } \Sigma_{Z(4)} - (\text{Im } \Sigma_{Z(3)}) (\text{Re } \Sigma'_{Z(1)}) \\ & + (\text{Im } \Sigma_{Z(2)}) [(\text{Re } \Sigma'_{Z(1)})^2 - \text{Re } \Sigma'_{Z(2)} - \frac{1}{4} (\delta Z_{(1)}^{\gamma Z})^2 - (\text{Im } \Sigma_{Z(1)}) (\text{Im } \Sigma''_{Z(1)})] \\ & + (\text{Im } \Sigma_{Z(1)}) [ -(\text{Re } \Sigma'_{Z(1)})^3 + 2(\text{Re } \Sigma'_{Z(2)}) (\text{Re } \Sigma'_{Z(1)}) - \text{Re } \Sigma'_{Z(3)} \\ & \quad - \frac{1}{2} \delta Z_{(1)}^{\gamma Z} \delta Z_{(2)}^{\gamma Z} + \frac{1}{2} (\text{Re } \Sigma'_{Z(1)}) (\delta Z_{(1)}^{\gamma Z})^2 - \frac{1}{2} (\text{Im } \Sigma_{Z(1)}) (\text{Im } \Sigma''_{Z(2)}) \\ & \quad \left. + \frac{3}{2} (\text{Im } \Sigma_{Z(1)}) (\text{Re } \Sigma'_{Z(1)}) (\text{Im } \Sigma''_{Z(1)}) + \frac{1}{6} (\text{Im } \Sigma_{Z(1)})^2 (\text{Re } \Sigma'''_{Z(1)}) \right\}_{s=\overline{M}_Z^2}. \end{aligned}$$

❑ **Also one will obtain unstable particles' total widths by imposing on shell condition. (as a consequence of optical theorem)**

# Computing EWPOs


- ❖  $G_\mu$  is determined from measuring muon decay after subtracting QED corrections within 4-Fermi theory.
- ❖ Then move on to the SM,  $G_\mu$  receives corrections depicted on the right hand side. One can then use such a relation to predict W-boson mass.



$G_F$  from  $\mu$  decay in Fermi Model

$$\Gamma_\mu = \frac{G_F^2 m_\mu^5}{192\pi^3} F\left(\frac{m_e^2}{m_\mu^2}\right) (1 + \Delta q)$$

QED corrections (2-loop)



Ritbergen, Stuart '98  
Pak, Czarnecki '08


$G_F$  decay in Standard Model

$$\frac{G_F^2}{\sqrt{2}} = \frac{e^2}{8s_W^2 M_W^2} (1 + \Delta r)$$

electroweak corrections

Freitas '18talk

$$\Delta r(M_W, M_Z, M_H, \dots) =$$



One gets an implicit relation between W-boson mass and G-Fermi:

$$\overline{M_W}^2 = \overline{M_Z}^2 \left( \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\alpha\pi}{\sqrt{2}G_\mu \overline{M_Z}^2} (1 + \Delta r)} \right)$$

- ❖ We have seen parity-violating asymmetry can be determined by effective weak-mixing angle  $\sin^2 \theta_{eff}^f$ . It relates to the ratio between dressed vector and axial-vector coupling.
- ❖ Using the decay rate equation in terms of dressed vector and axial-vector couplings. We can derive the total and partial width of Z-boson.

### Using optical theorem

$$\Im \Sigma_Z = \frac{1}{3M_Z} \sum_f \sum_{spins} \int d\Phi (|g_V^f|^2 + |g_A^f|^2)$$

Plugging what we have from OS condition in complex pole scheme.

$$\bar{\Gamma}_Z = \frac{N_c^f}{12\pi M_Z} C_Z (\mathcal{R}_V^f |g_V^f|^2 + \mathcal{R}_A^f |g_A^f|^2)$$


where  $C_Z$  features all self-energy contributions, and  $\mathcal{R}_{V,A}^f$  feature final-state QCD and QED corrections. Here for closed fermionic loops we set them to 1.

$$\sin^2 \theta_{eff}^f = \frac{1}{4|Q_f|} \left(1 - \Re \frac{g_V^f}{g_A^f}\right)_{s=M_Z^2}$$

$$g_V^f = Z_e (v_f^Z - Q_f \sqrt{Z_{\gamma Z}}) - v_f^\gamma \frac{\hat{\Sigma}_{\gamma Z}}{s + \hat{\Sigma}_{\gamma\gamma}}$$

$$g_A^f = Z_e a_f^Z - a_f^\gamma \frac{\hat{\Sigma}_{\gamma Z}}{s + \hat{\Sigma}_{\gamma\gamma}}$$

### Decompstion of the effective Zff vertex




## Technical Aspects

- ❑ In pure EW case, All loop integrals can be written as 1-loop scalar master integrals and their derivatives up to second order.
- ❑ Exact agreement at 2-loop was found comparing to previous work (hep-ph:004091;0202131;0407317;13102256), except one missing term as the second term in the following:


$$\Re \Sigma'_{ZZ(2)}(s) - \frac{d}{ds} \left( \frac{\Im \Sigma_{\gamma Z(1)}^2(s)}{s} \right)$$

of which numerical impact shall be investigated.

- ❑ Unlike pure EW, mixed EW-QCD at 3-loop order features non-unique master integral (**2-loop**) basis. ( difficult to cross-check symbolically)
- ❑ Integral reduction is non-trivial. (IBP and technique from G.Weiglein,R.Scharf et.al.hep-ph:9310358 were adopted in this work in parallel)
- ❑ The **derivative** of 2-loop master integral is needed.
- ❑ Both cases have been carried out in two independent implementations.




One of the two-loop master integral basis we use in calculations.




$B_0 A_0$




$B_0^2$




$A_0^2$




$T_3$




$T_{3a}$



$T_{4a}$



$T_{4a1}$



$T_{5a}$

## The derivative of Two-Loop Master Integral

Define a general two-loop scalar integral as

$$I(\nu_1, \nu_2, \dots, m_1, m_2, \dots; p^2) \equiv \int \frac{d^D q_1 d^D q_2}{(q_1^2 - m_1^2)^{\nu_1} ((q_1 + p)^2 - m_2^2)^{\nu_2} ((q_2 - q_1)^2 - m_3^2)^{\nu_3} (q_2^2 - m_4^2)^{\nu_4} ((q_2 + p)^2 - m_5^2)^{\nu_5}}$$

For  $p^2 = 0$

$$\begin{aligned} \frac{\partial}{\partial p^2} I(\dots; p^2 = 0) &= \frac{1}{2D} \frac{\partial^2}{\partial p_\mu \partial p^\mu} I(\dots; p^2) \Big|_{p^2=0} \\ &= \frac{2}{D} \left[ \left(1 + \nu_2 + \nu_5 - \frac{D}{2}\right) (\nu_2 I(\nu_2 + 1) + \nu_5 I(\nu_5 + 1)) \right. \\ &\quad + m_2^2 \nu_2 (\nu_2 + 1) I(\nu_2 + 2) + m_5^2 \nu_5 (\nu_5 + 1) I(\nu_5 + 2) \\ &\quad \left. + \nu_2 \nu_5 ((m_2^2 - m_3^2 + m_5^2) I(\nu_2 + 1, \nu_5 + 1) - I(\nu_2 + 1, \nu_3 - 1, \nu_5 + 1)) \right] \Big|_{p^2=0} \end{aligned}$$

For  $p^2 \neq 0$

$$\begin{aligned} \frac{\partial}{\partial p^2} I(\dots; p^2 \neq 0) &= -\frac{1}{2p^2} p^\mu \frac{\partial}{\partial p^\mu} I(\dots; p^2) \\ &= -\frac{1}{2p^2} \left[ (\nu_2 + \nu_5) I - \nu_2 I(\nu_1 - 1, \nu_2 + 1) - \nu_5 I(\nu_4 - 1, \nu_5 + 1) \right. \\ &\quad \left. + \nu_2 (m_2^2 - m_1^2 + p^2) I(\nu_2 + 1) + \nu_5 (m_5^2 - m_4^2 + p^2) I(\nu_5 + 1) \right] \end{aligned}$$

New  $I(\dots; p^2)$  can be further reduced down to a linear combination of the chosen master integrals. Such a process can be carried out by using IBP technique.



# Numeric and Algebraic Cross-check

- ❑ For pure EW case, we can cross-check on calculations algebraically at any level.
- ❑ For Mixed EW-QCD, due to the ambiguity in the choice of master integrals, only the UV part can be checked algebraically via TVID2.1. The finite parts are carried out numerically in TVID2.
- ❑ Some  $\mathcal{O}(4 - D)$  coefficients from scalar one-loops have been computed

## Numerical Inputs

- ❑ We turn-off the CKM mixing due to its negligible numerical impact.
- ❑ For  $\overline{MS}$  scheme, we change out top mass into  $m_t(\mu = m_t) = 163.229 \text{ GeV}$
- ❑ Due to the internal relation between  $G_\mu$  and W-boson mass, one can treat either one as induced from another. (Usually W-boson mass is predicted from  $G_\mu$ )

$$\begin{array}{l} M_Z = 91.1876 \text{ GeV} \\ \Gamma_Z = 2.4952 \text{ GeV} \end{array} \left. \vphantom{\begin{array}{l} M_Z \\ \Gamma_Z \end{array}} \right\} \Rightarrow \overline{M}_Z = 91.1535 \text{ GeV}$$
$$\begin{array}{l} M_W = 80.358 \text{ GeV} \\ \Gamma_W = 2.089 \text{ GeV} \end{array} \left. \vphantom{\begin{array}{l} M_W \\ \Gamma_W \end{array}} \right\} \Rightarrow \overline{M}_W = 80.331 \text{ GeV}$$
$$M_t = 173.0 \text{ GeV}$$
$$M_{f \neq t} = 0$$
$$\alpha_s = 0.1179$$
$$\alpha = 1/137.035999084$$
$$\Delta\alpha = 0.05900$$
$$G_\mu = 1.1663787 \times 10^{-5} \text{ GeV}^{-2}$$

# Numerical Results

## ❖ On Shell Scheme

### □ On-shell in pure EW case

$$\Delta r_{(3)} = 2.5 \times 10^{-5}$$

### □ On-shell in mixed EW-QCD case

$$\Delta r_{(\alpha^2 \alpha_s)} = -0.000109$$

	$\Delta \bar{M}_W$ [MeV]	$\Delta \sin^2 \theta_{\text{eff}}^f$	$\Delta' \sin^2 \theta_{\text{eff}}^f$	$\Delta \bar{\Gamma}_{\text{tot}}$ [MeV]	$\Delta' \bar{\Gamma}_{\text{tot}}$ [MeV]
$\mathcal{O}(\alpha^3)$	-0.389	$1.34 \times 10^{-5}$	$2.09 \times 10^{-5}$	0.331	0.255
$\mathcal{O}(\alpha^2 \alpha_s)$	1.703	$1.31 \times 10^{-5}$	$-1.98 \times 10^{-5}$	-0.103	0.229
Sum	1.314	$2.65 \times 10^{-5}$	$0.11 \times 10^{-5}$	0.228	0.484

	CEPC	FCC-ee	ILC/GigaZ
$M_W$ [MeV]	1	1	2.5
$\Gamma_Z$ [MeV]	0.5	0.1	1.0
$\sin^2 \theta_{\text{eff}}^f$ [ $10^{-5}$ ]	2.3	0.6	1

- the parametric shift of  $G_\mu$  can go into W-boson mass.

$$\Delta \bar{M}_W(\alpha^2 \alpha_s) \approx \frac{\pi \alpha \bar{M}_Z^2}{2\sqrt{2} G_\mu \bar{M}_W (\bar{M}_Z^2 - 2\bar{M}_W^2)} \Delta r_{(\alpha^2 \alpha_s)}$$

- Similarly, one gets effective weak mixing angle and Z width with leading W-boson mass shift

$$\Delta' \sin^2 \theta_{\text{eff},(\alpha^2 \alpha_s)}^f = \Delta \sin^2 \theta_{\text{eff},(\alpha^2 \alpha_s)}^f - \frac{2\Delta \bar{M}_W(\alpha^2 \alpha_s) \bar{M}_W}{\bar{M}_Z^2}$$

$$\Delta' \bar{\Gamma}_{f,(\alpha^2 \alpha_s)} = \Delta \bar{\Gamma}_{f,(\alpha^2 \alpha_s)} - \frac{2\Delta \bar{M}_W(\alpha^2 \alpha_s) \bar{M}_W}{\bar{M}_Z} \times \frac{\alpha N_c^f}{6s_W^4 c_W^4} [(2s_W^2 - 1)(I_3^f)^2 + 2s_W^4 Q_f(Q_f - I_3^f)]$$

❖ On-shell +  $\overline{MS}$  in mixed EW-QCD case.

$\Delta r_{(\alpha^2\alpha_s)} [10^{-4}]$	$\Delta M_{W(\alpha^2\alpha_s)} [\text{MeV}]$
-0.50	0.78

$X$	$\Delta X_{(\alpha^2\alpha_s)}$	$\Delta' X_{(\alpha^2\alpha_s)}$
$\sin^2 \theta_{\text{eff}} [10^{-5}]$	0.75	-0.76
$\Gamma_\ell [\text{MeV}]$	-0.0003	0.0047
$\Gamma_\nu [\text{MeV}]$	0.0009	0.0086
$\Gamma_d [\text{MeV}]$	-0.0018	0.0223
$\Gamma_u [\text{MeV}]$	-0.0029	0.0183
$\Gamma_{\text{tot}} [\text{MeV}]$	-0.0093	0.143

	on-shell $M_t$		$\overline{MS}$ $m_t$	
	$\mathcal{O}(\alpha^2)$	$\mathcal{O}(\alpha^2\alpha_s)$	$\mathcal{O}(\alpha^2)$	$\mathcal{O}(\alpha^2\alpha_s)$
$\Delta r [10^{-4}]$	7.85	-1.09	7.56	-0.50
$\Delta \sin^2 \theta_{\text{eff}}^f [10^{-5}]$	30.98	1.31	31.18	0.75
$\Delta \overline{\Gamma}_\ell [\text{MeV}]$	0.2412	-0.0157	0.2284	-0.0003
$\Delta \overline{\Gamma}_\nu [\text{MeV}]$	0.4145	-0.0002	0.4152	0.0009
$\Delta \overline{\Gamma}_d [\text{MeV}]$	0.6666	-0.0049	0.6780	-0.0018
$\Delta \overline{\Gamma}_u [\text{MeV}]$	0.4964	-0.0203	0.4911	-0.0029
$\Delta \overline{\Gamma}_{\text{tot}} [\text{MeV}]$	4.951	-0.103	4.947	-0.0093

❖ Comparing between two schemes

- ❑  $\overline{MS}$  Top mass must be used at previous order  $\mathcal{O}(\alpha^2)$  when using  $\overline{MS}$  renormalization scheme for top mass.
- ❑ A better convergence behavior from  $\overline{MS}$  is observed. Also the numerical size of corrections at given order gets reduced comparing to on shell scheme.
- ❑ Numerical numbers given by two schemes at each order are partially compensate each other.

❖ Numerical impact given by the missing term from previous study

$$\Delta\Gamma_{f(2)}|_{\text{this work}} - \Delta\Gamma_{f(2)}|_{\text{Freitas,Hollik,Walter,Weiglein '00,'02}}$$

$$= -N_c^f (v_{f(0)}^2 + a_{f(0)}^2) \overline{M}_Z \frac{25\alpha^2(3 - 8s_w^2)^2}{3888\pi s_w^2 c_w^2}$$

$$= \begin{cases} -0.0028 \text{ MeV} & \text{for } f = \ell, \\ -0.0056 \text{ MeV} & \text{for } f = \nu, \\ -0.0126 \text{ MeV} & \text{for } f = d, \\ -0.0098 \text{ MeV} & \text{for } f = u, \\ -0.0830 \text{ MeV} & \text{for } f = \text{tot.} \end{cases}$$

# Summary and Outlook

- ❑ EWPOs measurements at future electron-positron colliders require higher order corrections beyond 2-loop level.
  - ❑ Closed fermionic loops gets numerical enhancement from power of top mass and large multiplicity of light fermion d.o.f.
  - ❑ We present the results for contributions with maximal closed fermionic loops at given order  $\mathcal{O}(\alpha^3)$ ,  $\mathcal{O}(\alpha^2\alpha_s)$ .
  - ❑ Various aspects in renormalization: gauge invariance, complex pole mass, photon-Z-boson mixing, etc...
  - ❑ For mixed EW-QCD corrections, two different renormalization schemes on top mass were performed.
  - ❑ All results are carried out in two independent calculations, with the help of computer-algebra tools.
  - ❑ An error was found in previous work, we corrected it and investigated its numerical impact (very small).
- 
- ❑ The new results do not significantly reduce the theoretical error. Other missing three-loop contributions are needed.
  - ❑ The difference of the sum  $\mathcal{O}(\alpha^2) + \mathcal{O}(\alpha^2\alpha_s)$  between two schemes could be used to estimate of the size of the unknown higher-order  $\mathcal{O}(\alpha^2\alpha_s^2)$

THANK YOU.

## Backup Slides

□ Partial Widths of Z-boson decay in Pure EW case  $\mathcal{O}(\alpha^3)$

$X$	$l^+l^-$	$\nu\bar{\nu}$	$U\bar{U}$	$D\bar{D}$	Total
$\Delta\Gamma_X$ MeV	0.019	0.026	0.041	0.035	0.331
$\Delta'\Gamma_X$ MeV	0.017	0.022	0.029	0.024	0.255

□ partial Widths of Z-boson decay in Mixed EW-QCD case  $\mathcal{O}(\alpha^2\alpha_s)$

$X$	$l^+l^-$	$\nu\bar{\nu}$	$U\bar{U}$	$D\bar{D}$	Total
$\Delta\Gamma_X$ MeV	-0.0157	-2.0E-4	-0.0049	-0.0203	-0.103
$\Delta'\Gamma_X$ MeV	-0.0049	0.0166	0.0475	0.0260	0.2296



## Theoretical uncertainty due to missing higher order

- ❑ Collect all common prefactors, such as couplings, Lie algebra number, particle multiplicities and mass ratios.
- ❑ Vary renormalization scale ( $\overline{MS}$  only!), this is frequently used in QCD.
- ❑ Compare results from two different schemes.
- ❑ Extrapolate to higher order by assuming geometric series behavior of perturbation theory.

prefactor method yields

$$\mathcal{O}(\alpha_{\text{bos}}) \sim \Gamma_Z \alpha^2 \approx 0.13 \text{ MeV},$$

$$\mathcal{O}(\alpha^3) - \mathcal{O}(\alpha_t^3) \sim \Gamma_Z \alpha \alpha_t^2 \approx 0.12 \text{ MeV},$$

$$\mathcal{O}(\alpha^2 \alpha_s) - \mathcal{O}(\alpha_t^2 \alpha_s) \sim \Gamma_Z \frac{\alpha \alpha_t n_q}{\pi} \alpha_s(m_t) \approx 0.23 \text{ MeV},$$

$$\mathcal{O}(\alpha \alpha_s^2) - \mathcal{O}(\alpha_t \alpha_s^2) \sim \Gamma_Z \frac{\alpha n_q}{\pi} \alpha_s^2(m_t) \approx 0.35 \text{ MeV},$$

$$\mathcal{O}(\alpha \alpha_s^3) - \mathcal{O}(\alpha_t \alpha_s^3) \sim \Gamma_Z \frac{\alpha n_q}{\pi} \alpha_s^3(m_t) \approx 0.04 \text{ MeV}.$$

geometric series extrapolation yields

$$\mathcal{O}(\alpha_{\text{bos}}) \sim [\mathcal{O}(\alpha_{\text{bos}})]^2 \approx 0.10 \text{ MeV},$$

$$\mathcal{O}(\alpha^3) - \mathcal{O}(\alpha_t^3) \sim \frac{\mathcal{O}(\alpha^2)}{\mathcal{O}(\alpha)} [\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)] \approx 0.26 \text{ MeV},$$

$$\mathcal{O}(\alpha^2 \alpha_s) - \mathcal{O}(\alpha_t^2 \alpha_s) \sim \frac{\mathcal{O}(\alpha \alpha_s)}{\mathcal{O}(\alpha)} [\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)] \approx 0.30 \text{ MeV},$$

$$\mathcal{O}(\alpha \alpha_s^2) - \mathcal{O}(\alpha_t \alpha_s^2) \sim \frac{\mathcal{O}(\alpha \alpha_s)}{\mathcal{O}(\alpha)} [\mathcal{O}(\alpha \alpha_s) - \mathcal{O}(\alpha_t \alpha_s)] \approx 0.23 \text{ MeV},$$

$$\mathcal{O}(\alpha \alpha_s^3) - \mathcal{O}(\alpha_t \alpha_s^3) \sim \frac{\mathcal{O}(\alpha \alpha_s^2)}{\mathcal{O}(\alpha)} [\mathcal{O}(\alpha \alpha_s) - \mathcal{O}(\alpha_t \alpha_s)] \approx 0.035 \text{ MeV}.$$

$$\delta_{th} \Gamma_Z \sim 0.5 \text{ MeV}$$