

A Proposed Australian Facility for Particle and Nuclear Physics at the MeV-scale

Thursday 8 December 2022 17:10 (20 minutes)

Determining the nature of New Physics extensions to the Standard Model is one of the most pressing issues for Particle Physics. Well-motivated theories employ New Physics to solve the strong CP, hierarchy or axion quality problems by introducing new pseudoscalar particles which are weakly coupled to the standard model. These axion-like particles can have MeV–GeV masses and predominantly decay to photons and leptons [1, 2]. Recently, the ATOMKI group found evidence [3–5] for a new fundamental boson, named the X17, observed via $p + {}^7\text{Li} \rightarrow {}^8\text{Be} + (\text{X17} \rightarrow (e^+e^-))$, $p + {}^3\text{H} \rightarrow {}^4\text{He} + (\text{X17} \rightarrow (e^+e^-))$, and $p + {}^{11}\text{B} \rightarrow {}^{12}\text{C} + (\text{X17} \rightarrow (e^+e^-))$ reactions with a mass of 17 MeV and high statistical significance. There are now numerous searches for weakly coupled bosons, including the X17, using particle physics experiments [6–8]. However, only the ATOMKI group have utilized nuclear reactions in a competitive way to date. We intend to employ the Pelletron accelerator in Melbourne to initiate nuclear reactions of the kind: $p + {}^Z\text{X} \rightarrow {}^{Z+1}\text{Y} + (e^+e^-)$ and to build a low mass, high precision, Time Projection Chamber (TPC) with a micropatterned readout and magnetic field. The invariant mass resolution of the TPC to the (e^+e^-) final state is expected to be 0.1 MeV. This provides a substantially more sensitive search for anomalous (e^+e^-) production than any other experiment and 200 times more sensitivity than ATOMKI. Accordingly, we will either observe the ATOMKI anomaly or exclude it at very high significance. Following this, we propose a program to search for anomalous (e^+e^-) production with world-leading sensitivity in the 5–25 MeV mass region. In addition, the very large acceptance, and excellent angular and energy resolution of the TPC enables more sensitive investigations of nuclear internal pair conversion decays, enabling a range of novel nuclear physics investigations. The presentation will describe the proposed TPC, its expected performance, together with its application and anticipated impact in particle and nuclear physics investigations.

- [1] D. Alves Phys. Rev. D103 055018 (2021)
- [2] M. Bauer et al. arXiv:2110.10698
- [3] A.J. Krasznahorkay, et al., Phys. Rev. Lett. 116, 042501 (2016)
- [4] A.J. Krasznahorkay, et al., Phys. Rev. C104, 044003 (2021)
- [5] A.J. Krasznahorkay, et al., arXiv:2209.10795
- [6] A. M. Baldini et al. (MEG II Collaboration), Eur. Phys. J. C 78, 380 (2018)
- [7] J. Balewski et al. (DarkLight), arXiv:1412.4717
- [8] C. Ahdida et al. (SHiP Collaboration), arXiv:2010.11057

Authors: Dr BIGNELL, Lindsey (Department of Nuclear Physics and Accelerator Applications, The Australian National University); Prof. SEVIOR, Martin (School of Physics, University of Melbourne); Prof. KIBEDI, Tibor (Department of Nuclear Physics and Accelerator Applications, The Australian National University); Prof. JAMIESON, David (School of Physics, University of Melbourne); Prof. STUCHBERY, Andrew (The Australian National University); Dr THAMM, Andrea (University of Melbourne (AU))

Presenter: Dr BIGNELL, Lindsey (Department of Nuclear Physics and Accelerator Applications, The Australian National University)

Session Classification: BSM physics

Track Classification: Physics beyond the Standard Model