International Joint Workshop on the Standard Model and Beyond 2024 & 3rd Gordon Godfrey Workshop on Astroparticle Physics

Contribution ID: 69

Type: Contributed Talk

Revisiting Metastable Cosmic String Breaking

Tuesday 10 December 2024 16:50 (20 minutes)

Metastable cosmic strings are gathering attention as potential progenitors of stochastic gravitational wave background. They result from a two-step symmetry breaking $G \to H \to 1$ with $\pi_1(G) = 0$ and $\pi_1(H) \neq 0$, and decay via internal monopole-antimonopole pair creation.

Conventionally, the breaking rate has been estimated by an infinitely thin string approximation, which requires a large hierarchy between the symmetry breaking scales.

We numerically constructed a tunneling path and thus obtained a robust lower limit on the tunneling factor e^{-S} even for mild scale hierarchy. In particular, it is relevant to the cosmic string interpretation of the gravitational wave signals recently reported by pulsar timing array experiments.

Authors: CHITOSE, Akifumi (Institute for Cosmic Ray Research, University of Tokyo); WATANABE, Keiichi; IBE, Masahiro (Institute for Cosmic Ray Research, University of Tokyo); SHIRAI, Satoshi (Kavli IPMU); Mr NAKAYAMA, Yuhei (The University of Tokyo)

Presenter: CHITOSE, Akifumi (Institute for Cosmic Ray Research, University of Tokyo)

Session Classification: Cosmology