

GRAVITATIONAL DARK MATTER PORTAL IN EXTRA DIMENSIONS

N. Rius

With **N. Bernal, A. Donini and M.G. Folgado,**

JHEP 01 (2020) 161, *JHEP* 09 (2020) 142, *Eur.Phys.J.C* 81 (2021) 3, 197

Dark Matters 2022, Brussels



Outline




1) Introduction

2) Freeze-out DM

3) Freeze-in DM

4) Summary and outlook

1) Introduction

- Only gravitational evidence of DM ...
- ... but maybe in extra dimensions there are possible observational signatures
- Extra Dimension theories were proposed at the end of the XX century in order to solve the hierarchy problem.
- Large Extra-Dimensions (LED)  extra dimensions flat
- Randall-Sundrum (RS)  warped extra dimensions
- Clockwork/Linear Dilaton (CW/LD) 

Giudice, McCullough, JHEP 02(2017)036

Mini-review of extra dimension scenarios

5-Dimensional Metric

$$ds^2 = e^{2\sigma(y)} (\eta_{\mu\nu} dx^\mu dx^\nu - e^{-2l\sigma(y)} r_c^2 dy^2)$$

Large Extra-Dimensions

- $\sigma(y) = 0$

$$\bar{M}_{pl}^2 = M_5^3 2\pi r_c$$



Volume factor

Randall-Sundrum

- $\sigma(y) = -kr_c|y|$

- $l = 1$

$$\bar{M}_{pl}^2 = \frac{M_5^3}{k} (1 - e^{-2k\pi r_c})$$

$$k, M_5 \sim \bar{M}_{pl}$$

Clockwork/Linear Dilaton

- $\sigma(y) = \frac{2}{3}kr_c|y|$

- $l = 0$


$$\bar{M}_{pl}^2 = \frac{M_5^3}{k} (e^{2k\pi r_c} - 1)$$



Warping factor


$$k, M_5 \ll \bar{M}_{pl}$$

Randall-Sundrum



$$ds^2 = e^{-2kr_c|y|} \eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 dy^2$$

Clockwork/LD



$$ds^2 = e^{\frac{4}{3}kr_c|y|} (\eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 dy^2)$$

R-S: $S = S_{\text{Bulk}} + S_{\text{IR}} + S_{\text{UV}} \quad y = r_c \phi$

$$S_{\text{Bulk}} = \frac{M_5^3}{2} \int d^4x \int_0^\pi r_c d\phi \sqrt{G} (\mathcal{R} - 2\Lambda_5)$$

$$S_{\text{IR}} = \int d^4x \int_0^\pi r_c d\phi \sqrt{-g_{\text{IR}}} (-V_{\text{IR}} + \mathcal{L}_{\text{IR}}) \delta(\phi - \pi)$$

$$S_{\text{UV}} = \int d^4x \int_0^\pi r_c d\phi \sqrt{-g_{\text{UV}}} (-V_{\text{UV}} + \mathcal{L}_{\text{UV}}) \delta(\phi)$$


- 5D Graviton field $g_{\mu\nu} = \eta_{\mu\nu} + \hat{h}_{\mu\nu}$
- KK decomposition: $\hat{h}_{\mu\nu}(x, y) = \sum_{n=0}^{\infty} h_{\mu\nu}^n(x) \frac{\chi_n(y)}{\sqrt{r_c}}$
- $(\eta^{\mu\nu} \partial_\mu \partial_\nu + m_n^2) h_{\mu\nu}^n(x) = 0$
- In R-S: $\chi^n(y) = \frac{e^{2\sigma(y)}}{N_n} [J_2(z_n) + \alpha_n Y_2(z_n)]$ $z_n(y) = m_n/k e^{\sigma(y)}$
- In CW/LD: $\chi_n(y) = N_n e^{-ky} [\sin(\beta_n y) + \omega_n \cos(\beta_n y)]$

5D massless graviton




4D tower massive KK gravitons

Randall-Sundrum



$$ds^2 = e^{-2kr_c|y|} \eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 dy^2$$




Mass spectra


$$m_n = kx_n e^{-k\pi r_c} \longleftrightarrow J_1(x_n) = 0$$

$$m_0 = 0$$

Clockwork/LD



$$ds^2 = e^{\frac{4}{3}kr_c|y|} (\eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 dy^2)$$



Mass spectra

$$m_n^2 = k^2 + \frac{n^2}{r_c^2}$$

$$m_0 = 0$$

$k = 0$
LED

Brane distance stabilization mechanism

The distance between the two 4D-branes is determined by r_c , to stabilise dynamically this distance a scalar field in the 5D-bulk is introduced (Radion).

Randall-Sundrum

m_r → **New Free parameter**

Goldberger-Wise mechanism:
typically $m_r < m_1$

Clockwork/Linear Dilaton

- Already present bulk dilaton field (Φ_n).
- The 5D dilaton field can be written as a KK tower:

$$m_r^2 = m_{\Phi_0}^2 = \frac{8}{9}k^2$$

$$m_{\Phi_n}^2 = k^2 + \frac{n^2}{r_c^2}$$

- Here, I focus on Randall–Sundrum scenario (dual to a 4D strongly interacting model)
- SM and DM in the IR brane
- CW/LD studied in A. Donini, M.G. Folgado, N. Rius, *JHEP* 04 (2020) 036, N. Bernal et al., *JHEP* 04 (2021) 061

- From the weak field expansion of the metric:

$$ds^2 = e^{-2kr_c y} e^{-2r} (\eta_{\mu\nu} + \hat{h}_{\mu\nu}) dx^\mu dx^\nu - (1 + 2r)^2 dy^2$$

- Graviton-matter interaction:

$$\mathcal{L} = -\frac{1}{M_5^{3/2}} T^{\mu\nu}(x) h_{\mu\nu}(x, y = \pi) = -\frac{1}{M_P} T^{\mu\nu}(x) h_{\mu\nu}^0(x) - \frac{1}{\Lambda} \sum_{n=1} T^{\mu\nu}(x) h_{\mu\nu}^n(x)$$


- Radion-matter interaction:

$$\mathcal{L}_r = \frac{1}{\sqrt{6}\Lambda} r T + \frac{\alpha_{\text{EM}} C_{\text{EM}}}{8\pi\sqrt{6}\Lambda} r F_{\mu\nu} F^{\mu\nu} + \frac{\alpha_S C_3}{8\pi\sqrt{6}\Lambda} r \sum_a F_{\mu\nu}^a F^{a\mu\nu}$$

$$\Lambda = \bar{M}_{pl} e^{-\pi k r_c}$$

- Gravitational field interactions (3rd order expansion)

2) Freeze-out DM



$$Y \equiv n(T)/s(T)$$

$$s(T) = \frac{2\pi^2}{45} g_{*s}(T) T^3$$


$$x \equiv m_{DM}/T$$

$$\frac{dY}{dx} = \frac{-x \langle \sigma v \rangle s}{H(m_{DM})} (Y^2(x) - Y_{eq}^2(x))$$

$$\begin{aligned} \langle \sigma v \rangle &\sim \int_{4m_{DM}}^{\infty} ds (s - 4m_{DM}) \sqrt{s} \sigma_{an}(s) K_1(\sqrt{s}/T) \\ &= 2.2 \times 10^{-26} \text{ cm}^3/\text{s} \end{aligned}$$

➤ Only DM-SM scattering (mediated by KK gravitons)



Rueter, Rizzo, Hewett, *JHEP* 10 (2017) 094






- SM + DM in the IR brane
- 3rd generation quarks in the IR brane, fermions near the UV brane, gauge fields in the bulk (avoid LHC strong bounds)
- All SM in the bulk, with different BLKTs (to explain fermion masses and CKM mixing matrix)


➤ DM annihilation into KK gravitons Min Lee, Park, Sanz
Eur.Phys.J.C 74 (2014) 2715, *JHEP* 05 (2014) 063

- DM and Higgs in the IR brane, SM in the UV brane
- Only DM in the IR brane





	Scalar	Fermion	Vector
Graviton Virtual Exchange	v^4 (d)	v^2 (p)	v^0 (s)
Radion Virtual Exchange	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Gravitons	v^0 (s)	v^0 (s)	v^0 (s)
Annihilation into Radions	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Radion + Graviton	v^0 (s)	v^0 (s)	v^0 (s)






	Scalar	Fermion	Vector
Graviton Virtual Exchange	v^4 (d)	v^2 (p)	v^0 (s)
Radion Virtual Exchange	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Gravitons	v^0 (s)	v^0 (s)	v^0 (s)
Annihilation into Radions	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Radion + Graviton	v^0 (s)	v^0 (s)	v^0 (s)






	Scalar	Fermion	Vector
Graviton Virtual Exchange	v^4 (d)	v^2 (p)	v^0 (s)
Radion Virtual Exchange	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Gravitons	v^0 (s)	v^0 (s)	v^0 (s)
Annihilation into Radions	v^0 (s)	v^2 (p)	v^0 (s)
Annihilation into Radion + Graviton	v^0 (s)	v^0 (s)	v^0 (s)



Gravitational portal to DM in Xdims

Anomalous enhancement due to the sum over massive KK graviton polarizations



$$m_r = 100 \text{ GeV}, M_{G_1} = 1 \text{ TeV}, \Lambda = 10 \text{ TeV}$$


$$\sigma(\text{DM}, \text{DM} \rightarrow G_n G_n) \propto \frac{s}{\Lambda^2} \left(\frac{s}{m_n^2} \right)^k$$

fermion: $k = 2$
 scalar, vector: $k = 4$

Scalar DM


- It is needed to sum over the infinite tower of KK gravitons in the s channel

Giorgi, Vogl, *JHEP* 04 (2021) 143;
JHEP 11 (2021) 036



Subtle cancellations due to sum rules of Bessel functions (wave functions of KK gravitons)

Radion contribution is also essential



Gravitational portal to DM in X_{dims} $m_r = 100 \text{ GeV}, M_{G_1} = 1 \text{ TeV}, \Lambda = 20 \text{ TeV}$


Final result, as expected : $\sigma(\text{DM}, \text{DM} \rightarrow G_n G_n) \propto \frac{s}{\Lambda^2}$

Analogous result for


$$\sigma(\text{DM}, \text{DM} \rightarrow G_n r) \propto \frac{s}{\Lambda^2} \left(\frac{s}{m_n^2} \right)^2$$

- Vector and fermion DM, as well as CW/LD: work in progress with A. Donini, G. Landini, A. Muñoz

Bounds from resonance searches at LHC



ATLAS: Search for new phenomena in high-mass **dilepton** final states using 37 fb^{-1}
(1707.02424)




ATLAS: Search for new phenomena in high-mass **diphoton** final states using 37 fb^{-1}
(1707.04147)


Scalar

fermion

vector



3) Freeze-in DM




DM is a Feebly Interacting Massive Particle (FIMP)
It never reaches thermal equilibrium

Scalar DM

Direct Freeze In


- DM production via virtual graviton and radion exchange



Relevant when G, ϕ are in thermal equilibrium or $T < m_r, m_1$

Sequential Freeze In

- DM production via graviton and radion decay.



- Sequential freeze-in: Boltzmann equations for DM, G and ϕ
- SM-G_n G_m scattering suffers from unphysical divergence when $m_1 \ll s \sim T^2$ (work in progress)
- Approx. solutions:
 - 1) Direct freeze-in: $T < m_r$

$$\frac{dY}{dT} \simeq \frac{\gamma_{\text{DM} \rightarrow \text{SM}}}{H s T} \left[\left(\frac{Y}{Y^{\text{eq}}} \right)^2 - 1 \right] \simeq -\frac{\gamma_{\text{DM} \rightarrow \text{SM}}}{H s T}$$

$$\sigma_{\text{DM} \rightarrow \text{SM}}(s) \simeq \frac{49}{1440\pi} \frac{s^3}{\Lambda^4} \left| \sum_{n=1}^{\infty} \frac{1}{s - m_n^2 + i m_n \Gamma_n} \right|^2 + \frac{s^3}{288\pi\Lambda^4} \frac{1}{(s - m_r^2)^2 + m_r^2 \Gamma_r^2}$$

$$Y_0 \simeq \frac{3 \times 10^{-1}}{g_{*s}} \sqrt{\frac{10}{g_*}} \left(\frac{M_P}{m_r^4 \Lambda^4} \right)$$

2) Sequential freeze in via inverse decays:

$$\begin{aligned} \frac{dY}{dT} &\simeq \frac{\gamma_{\text{KK} \rightarrow \text{SM}}^d}{H s T} \left[\frac{Y_K}{Y_K^{\text{eq}}} - 1 \right] \text{BR}(\text{KK} \rightarrow \text{DM}) + \frac{\gamma_{\text{r} \rightarrow \text{SM}}^d}{H s T} \left[\frac{Y_r}{Y_r^{\text{eq}}} - 1 \right] \text{BR}(\text{r} \rightarrow \text{DM}) \\ &\simeq -\frac{1}{H s T} \left[\gamma_{\text{KK} \rightarrow \text{SM}}^d \text{BR}(\text{KK} \rightarrow \text{DM}) + \gamma_{\text{r} \rightarrow \text{SM}}^d \text{BR}(\text{r} \rightarrow \text{DM}) \right] \end{aligned}$$

Summing over G_n with $m_n < T_{\text{rh}}$:


$$\begin{aligned} Y_0 &\simeq \frac{2.2 \times 10^{-4}}{g_{*s}} \sqrt{\frac{10}{g_s} \frac{M_P T_{\text{rh}}^2}{m_1 \Lambda^2}} + \frac{3.5 \times 10^{-2}}{g_{*s}} \sqrt{\frac{10}{g_s} \frac{M_P m_r}{\Lambda^2}} \left(\frac{z}{z + 37} \right) \\ z_n &\equiv \left(1 - 4 \frac{m_\chi^2}{m_n^2} \right)^{5/2}, \\ z &\equiv \sqrt{1 - 4 \frac{m_\chi^2}{m_r^2}} \left(1 + 2 \frac{m_\chi^2}{m_r^2} \right)^2 \end{aligned}$$


3) Sequential freeze in via annihilations

$$\frac{dY}{dT} \simeq -\frac{1}{H s T} [\gamma_{\text{KK} \rightarrow \text{SM}} \text{BR}(\text{KK} \rightarrow \text{DM}) + \gamma_{\text{r} \rightarrow \text{SM}} \text{BR}(\text{r} \rightarrow \text{DM})]$$

$$\gamma_{\text{KK} \rightarrow \text{SM}}(T) \simeq 4.8 \times 10^4 \frac{T^{16}}{\Lambda^4 m_n^8} \quad (\text{to be corrected})$$


$$\gamma_{\text{r} \rightarrow \text{SM}}(T) \simeq 2.2 \times 10^{-4} \frac{T^8}{\Lambda^4}$$





Fermionic DM

- Only sequential freeze-in via inverse decays
- Radion contribution suppressed, since $\propto m_{\text{DM}}^2$
- Constraints from velocity distribution of DM: too fast due to KK graviton late decays



Giorgi, Vogl, arXiv:2208.03153

Thermalization
 BBN ($\tau_1 > 10s$)
 Warm DM bound
 meshed: KK no LL
 Strong dependence
 on T_{rh}
 Constant T_{rh}

4) Summary and outlook

- Importance of summing over all KK graviton modes to recover unitarity
- WIMP DM freeze-out strongly constrained by LHC (unless SM is not confined on the IR brane)
- Giving up the hierarchy problem, plenty of room for FIMP DM freeze-in
- To do:
 - Sum rules for fermion and vector scattering to G_n , G_m , and G_n in RS scenario (Donini, Landini, Muñoz)
 - Sum rules for other Xdim models: CW/LD, Flat
 - Gravitational portal to DM in $Xdim$ models (Bernal, Cosme, Donini)