



Contribution ID: 41

Type: **not specified**

## Plasma Wave Recognition by Multipoint Filtering Techniques

*Thursday 29 April 2021 19:05 (15 minutes)*

Turbulent wave fluctuations of plasma parameters obtained from the spacecraft measurements have mixed manifestations of spatial and temporal scales due to the dynamics of the considered environment. Decomposing of such plasma perturbations into frequency and spatial range have allowed us to analyse power distribution in branches of dispersion equations. We have demonstrated efficiency of each specific method in resolving  $\omega - k$  distribution in magnetospheric events. These techniques such as k-filtering, MSR (Multi-point Signal Resonator), phase difference methods yield the power spectrum  $P = P(\omega, k)$  in four-dimensional  $(\omega, k)$  space, and thereafter for a fixed value of the frequency  $\omega_0$  it is possible to study the power distribution of fluctuation in spatial dimensions.

The work was conducted in the frame plasma physics program of the National Academy of Sciences of Ukraine, the grant Az. 97 742 from the Volkswagen Foundation ("VWStiftung") and International Institution of Space Research (ISSI-BJ).

**Author:** Mr PETRENKO, Bohdan (Taras Shevchenko National University of Kyiv, Kyiv, Ukraine)

**Co-authors:** LUI, A. (Johns Hopkins University Applied Physics Laboratory, Laurel MD, USA); GRIGORENKO, E. (Space Research Institute, RAS, Russia); KRONBERG, E. (Max Planck Institute for Solar System Research, Göttingen, Germany); KOZAK, L. (Kyiv Taras Shevchenko University of Kyiv, Kyiv, Ukraine; Space Research Institute National Academy of Sciences of Ukraine and State Space Agency of Ukraine, Kyiv, Ukraine)

**Presenter:** Mr PETRENKO, Bohdan (Taras Shevchenko National University of Kyiv, Kyiv, Ukraine)

**Session Classification:** Solar physics and heliosphere; Atmospheric studies and space geophysics