Polylogs

Def Let $s = (s_1, \dots, s_e)$ positive index

$t \in \mathbb{C}$ $|t| < 1$

$$Li_s(t) = \sum_{n_1 > n_2 > \dots > n_e \geq 1} \frac{t^{n_1}}{n_1^{s_1} \dots n_e^{s_e}}$$

(multiple) polylog funct° of 1 variable)

Similarly, $$Li_s(t_1, \dots, t_e) = \sum_{n_1 > n_2 > \dots > n_e \geq 1} \frac{t_1^{n_1} t_2^{n_2} \dots t_e^{n_e}}{n_1^{s_1} \dots n_e^{s_e}}$$

with $|t_1| < 1$ $\quad |t_i| \leq 1 \quad \forall i \geq 2$.

Prop: cplx analysis : if $s_1 \geq 2$, $Li_s(t)$ is holomorphic on the open unit disc. It and can be extended continuously to the closed disc $|t| \leq 1$ and satisfies

$$Li_s(1) = \zeta(s)$$

$$Li_2(1) = \sum_{n_1 \geq 1} \frac{1}{n^2}$$

$\int_0^t$ int along straight path

Sweet relations $$Li_1(t) = \sum_{n_1 \geq 1} \frac{t^{n_1}}{n_1} = -\log(1-t) = \int_0^t \frac{dt_1}{1-t_1}$$

$$Li_2(t) = \sum_{n_1 \geq 1} \frac{t^{n_1}}{n_1^2} = -\int_0^t \log(1-t_1) \frac{dt_1}{t_1}$$

primitive of $\frac{Li_1(t)}{t}$ vanishing at $t=0$

Prop functional equations

- $$\int_0^t Li_{s_1, \dots, s_e}(t_1) \frac{dt_1}{t_1} = Li_{s_1+1, \dots, s_e}(t)$$
- $$\int_0^t Li_{s_1, \dots, s_e}(t_1) \frac{dt_1}{1-t_1} = Li_{1, s_1, \dots, s_e}(t)$$

proof

$$\int_0^t \sum_{n_1 > n_2 \cdots > n_e \geq 1} \frac{t_1^{n_1 - 1}}{n_1^{s_1} \cdots n_e^{s_e}} \, dt_1$$

$$= \sum_{n_1 > \cdots > n_e \geq 1} \frac{t_1^{n_1}}{n_1^{s_1 + 1} \cdots n_e^{s_e}}$$

$$\int_0^t \sum_{n_1 > \cdots > n_e \geq 1} \frac{t_1^{n_1}}{n_1^{s_1} \cdots n_e^{s_e}} \frac{dt_1}{1 - t_1}$$

$$\frac{t_1^{n_1 + k - 1}}{n_1^{s_1} \cdots n_e^{s_e}}$$

$$\rightsquigarrow \quad \frac{t_1^{n_1 + k} \, n_0}{(n_1 + k) n_1^{s_1} \cdots}$$

n_0

Thm If $0 < t < 1$ real number

$$Li_s(t) = \int_{\Delta^{wt(s)}(t)} \omega_s$$

proof by induction on $wt(s)$ using functional eq.

$$\Delta^n(t) = \{ t \leq t_n \leq \cdots \leq t_1 \leq 0 \}$$

Appear in Feynman ~~integral~~ amplitudes computation

(not all of them but huge class of them)

SRS
SwissMAP Research
Station in Les Diablerets

Feynman amplitudes

"scattering amplitudes"

We represent interactions between particles using graphs



two particles scattering (ie interacting) here exchanging a photon then scattering apart.



same, but photon spontaneously becomes a positron/e pair that later annihilates & forms a photon again.

- one has to consider all possible interactions (∞, yay!) (more and more loops/orders)
- have to associate $G \longmapsto I_G$ s.th we can work with

I_G amplitude given by an integral

function of masses & momenta of particles.

= Q answer to "how likely is this scatter"?

Predictions are obtained by summing over all graphs.

$\sum_G I_G$ diverges?

We'll work in scalar Euclidean space-time $\mathbb{R}^D$, $D>0$ even

norm $x^2 = \sum x_i^2$

SRS
SwissMAP Research Station in Les Diablerets

Feynmann graphs

following Brown

- $G = (V_G, E_G^{int}, E_G^{ext})$
  $N_G = \#E_G^{int}$
  G connected for us today
  together with the data
- masses $m_e \in \mathbb{R}_{\geq 0}$ $\forall e \in E_G^{int}$
- momenta $q_i \in \mathbb{R}^d$ $\forall i \in E_G^{ext}$ subject to momentum conservation

convention: all arrows incoming $q_i < 0$ means outgoing particle

$$\sum_{i \in E_G^{ext}} q_i = 0$$

- internal edges "classic edges" $E_G^{int} \to Sym^2 V_G$
- external edges particles linked to a single vertex.



To each internal edge we assign a momentum variable $k_e \in \mathbb{R}^D$ (virtual particle)

⇒ integration variable since we don't know the internal momenta (def by dig struct + cons. laws)

we integrate over all poss

$$p_e = \begin{cases} k_e \text{ if } e \in E^{int} \\ q_e \text{ if } e \in E^{ext} \end{cases}$$

& restricts the domain of int to where momenta is conserved

→ Dirac

$$I_G(q,m) = \int_{\mathbb{R}^{D \times N_G}} \prod_{e \in E_G^{int}} \frac{d^D k_e}{k_e^2 + m_e^2} \cdot \prod_{v \in V_G} \delta\Big(\sum_{\substack{e \text{ edges} \\ \text{which meet} \\ \text{at } v}} p_e\Big)$$

momentum cons at every vertex

momentum space FI

- may diverge
- looks terrifying.

⇒ use parametric representation to benefit from iterated integral setting

Rq: No coupling constants — simplified setting
Panzer, Brown

How to make it palatable ?

① Schwinger's trick

$$\frac{1}{\alpha} = \int_0^\infty e^{-\alpha x}\, dx \qquad x > 0$$

Idea: we introduce a new variable (= Schwinger parameter) for every internal edge and replace $\frac{1}{k_e^2 + m_e^2}$ with $\int_0^\infty e^{-(k_e^2+m_e^2)\alpha_e} d\alpha_e$

$$\rightsquigarrow \int_0^\infty \int_{\mathbb{R}^{D \cdot N_G}} \exp\left(-\sum_e (k_e^2+m_e^2)\alpha_e\right) d \cdot \prod \delta(\Sigma_{pe}) \prod_{e \in E^{int}_G} d\alpha_e \; (d\alpha_1 \cdots d\alpha_{N_G})$$

② Gauss $\int_{\mathbb{R}} e^{-\pi x^2} dx = 1$ $\quad x, b \in \mathbb{R}^D$

$$\int_{\mathbb{R}^D} e^{Q(x)} dx = \frac{\pi^{D/2}}{\sqrt{\det A}} e^{b^T A^{-1} b}$$

$Q(x) = -x^T A x + 2 b^T x$, A sym + def pos $D \times D$

$\rightsquigarrow$ D-dimensional equivalent

Example



(dirac fund.) $k_1 + k_2 = q$ (or int domain)

$$-\sum_e (k_e^2+m_e^2)\alpha_e$$
$$= \alpha_1(k_1^2+m_1^2) + \alpha_2(k_2^2+m_2^2)$$
$$= \alpha_1 k_1^2 + \alpha_2 (q-k_1)^2 + \alpha_1 m_1^2 + \alpha_2 m_2^2$$

complete the square

$$= (\alpha_1+\alpha_2)\left(k_1 - q\frac{\alpha_2}{\alpha_1+\alpha_2}\right)^2 + q^2 \frac{\alpha_1\alpha_2}{\alpha_1+\alpha_2} + \alpha_1 m_1^2 + \alpha_2 m_2^2$$

variable shift $k_1 = k_1 - q\frac{\alpha_2}{\alpha_1+\alpha_2}$

$$= (\alpha_1+\alpha_2)k^2 + \text{——————}$$

Integral is translation inv.

Deduce that $I_G(q,m) = \pi^{D/2} \int_0^\infty \frac{d\alpha_1 d\alpha_2}{(\alpha_1+\alpha_2)^{D/2}} \exp\left(\frac{-\alpha_1\alpha_2}{\alpha_1+\alpha_2} q^2 - m\chi^2 - m_3^2 \alpha_3\right)$

SRS
SwissMAP Research
Station in Les Diablerets

Mark's ingral (complete the square, etc)
one gets the determinant of a certain matrix that are then expressed in terms of graph theoretical data.

$$I_G(q,m) = \text{trivial factors} \times \frac{\prod_{h_G/2}}{(2\pi)^{D h_G}} \int_0^\infty \frac{\prod_{e\in E^{int}_G} d\alpha_e}{\psi_G^{D/2}} \exp\left(-\frac{\phi_G(q)}{\psi_G} - \sum_{e\in E^{int}_G} m_e^2 \alpha_e\right)$$

where ψ_G and $\phi_G(q)$ are called **Symanzik polyn**

$\psi_G \in \mathbb{Z}[\alpha_e, e \in E^{int}_G]$
- doesn't depend on external momenta nor masses

$$\psi_G = \sum_{\substack{T\subseteq G \\ \text{spanning tree}}} \prod_{e\notin T} \alpha_e$$

T spanning tree of G
$\hookrightarrow V_T = V_G$
T tree, subset of E^{int}_G

eg:



$\psi_G = \alpha_2 + \alpha_1$



$\psi_G = \alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_1\alpha_3$

$\phi_G(q) \in \mathbb{Z}[\alpha_e, e\in E^{int}_G, q_i\cdot q_j]$ ← Euclidean product
scalar prod of ext. momenta

$$\phi_G(q) = -\sum_{\substack{T=T_1\cup T_2 \\ \text{spanning 2-trees}}} q^{T_1}\cdot q^{T_2} \prod_{e\notin T} \alpha_e$$

↳ forest with 2 components, simply connected, spans
q^{T_i} = total momentum entering T_i



$\rightarrow q$ $\leftarrow -q$ $\quad -(q\cdot -q)\alpha_1\alpha_2$
$= +q^2\alpha_1\alpha_2$







$$-q_1(q_2+q_3)\alpha_2\alpha_3 - \dots -$$
$$= q_1^2\alpha_2\alpha_3 + q_2^2\alpha_2\alpha_3 + q_3^2\alpha_1\alpha_2$$

Rq: Ψ_G is homogeneous of degree h_G = #loops of G

$\Phi_G(q)$ hom. in α_e of degree h_G+1

Now we just want to get rid of the exp factor

- ~~Dim regularized~~ ~~since~~ this homogeneity means we can rescale. $\alpha_e \to \lambda\alpha_e$

change of var $\alpha_e = \lambda\beta_e$ $\sum\beta_e = 1$ $\lambda\in\mathbb{R}^{>0}$

$$\prod d\alpha_e = \lambda^{N_G-1}\, d\lambda \wedge \Omega_G$$

Up to now: • started from product of propagators $\frac{1}{p_e^2+m_e^2}$ integrated over internal momenta p_e

• Schwinger parameters, get exponentials

• Gaussian integrals ⇝ collapses int. into det.

• det are related to trees

→ parametric form depending only on graph. consts + external data.

• To get rid of the exp factor, one more step / integration to get a projective integral

↳ $\Omega_G = \sum_i (-1)^i \beta_i\, d\beta_1\wedge\dots\wedge\widehat{d\beta_i}\wedge\dots d\beta_n$

$$= \int_{\substack{\beta_e>0\\ \sum\beta_e=1}} \frac{\Omega_G}{\Psi_G^{D/2}} \int_0^\infty e^{-\lambda}\,(\text{usual stuff})\,\lambda^{N_G - D/2\,h_G}\,\frac{d\lambda}{\lambda}$$

(usual stuff) computable

$$= \Gamma(-N_{\mathrm{d}}(G))\left(\frac{\Psi_G}{\Phi_G(q)|_{\sum m_e\alpha_e|\Psi_G}}\right)^{-s_{\mathrm{d}}(G)}$$

Γ value s_{d}

SRS
SwissMAP Research
Station in Les Diablerets

Where $sdG = -N_G + D/2\, h_G$ "superficial degree of div of the graph"

>0 div
≤ 0 cv

logarithmic div when $= 0$
(as special case in a reg)

$$\Rightarrow I_G = \underbrace{\Gamma(N_G - D/2\, h_G)}_{\text{hello poss. poles}} \int \frac{\Omega_G}{\Psi_G^{D/2}} \cdot \left(\frac{\Psi_G}{\phi_G(a) + (\sum m_e^2 x_e)\Psi} \right.$$

$\sigma \subseteq \mathbb{P}^{N_G - 1}(\mathbb{R})$ — sd(

$\sigma = \{(B_1 : \dots : B_{N_G}) \in \mathbb{P}^{N_G-1}(\mathbb{R}) \mid B_i \geq 0\}$

'nam of deg 0

proj. m

If pole $\rightarrow$ Dimensional reg

replace D with $D - \varepsilon$, perform Taylor exp in ε

compute coeffs of ε^i

SRS
SwissMAP Research
Station in Les Diablerets

Panorama

- The general 1-loop diagram can be expressed using

$$\left.\begin{array}{l} Li_2(x) = \sum_{n\geq 1}^{\infty} \frac{x^n}{n^2} \quad |x|\leq 1 \\ \log(x) \end{array}\right\} = \int_{0\leq t_1\leq t_2\leq x} \frac{dt_1}{1-t_1}\frac{dt_2}{t_2}$$

with x some function of masses & momenta

- General 2-loop : not known.

ex Sunrise



solved for all m, q.
not !

iterated elliptic integrals

- A huge class of amplitudes are known & expressible using multiple polylogs

$$Li_{n_1,\dots,n_r}(x_1,\dots,x_r) = \sum_{1\leq k_1\leq\dots\leq k_r} \frac{x_1^{k_1}\cdots x_r^{k_r}}{k_1^{n_1}\cdots k_r^{n_r}}$$

$|x_i|<1$, extends to a multivalued function on

$$\left(\mathbb{C}\setminus\{0,1\}\right)^r \setminus \bigcup_{1\leq i\leq j\leq r} \{x_i x_{i+1}\cdots x_j = 1\}$$

$$\cong \mathcal{M}_{0,n+3}(\mathbb{C})$$

Single scale processes : our nice stuff

$I_G(q,m)$ is a function of a single variable

(e.g. all masses are 0 or only 1 momentum)

$\rightsquigarrow$ I_G boils down to a number.

Residues in massless ϕ^4 $\quad D=4$ all $m_e=0$

G in ϕ^4 $\leftrightarrow$ all vertices of G are of degree 4



$N_G = 2h_G \Rightarrow \Gamma\left(N_G - \frac{D}{2}h_G\right)$ has a pole

Residue

$$I_G = \int_\sigma \frac{\Omega_G}{\Psi_G^2}$$

This thing will cv iff G is "primitive"

i.e. all of its proper subgraphs verify $N_\gamma > 2h_\gamma$

Ex : ∘ 

$$I_G = \int_\sigma \frac{\alpha_1 d\alpha_2 - \alpha_2 d\alpha_1}{(\alpha_1+\alpha_2)^2}$$

$$= \int_{\alpha_2=0}^{\infty} \frac{d\alpha_2}{(1+\alpha_2)^2} = 1$$

∘ Next interesting graphs

$I_G = 6\zeta(3)$

WS3 detailed fun

$I_G = 20\zeta(5)$ $I_G = 36\zeta(3)^2$

next with 6 loops → lin combs of $\zeta(5,3)$, $\zeta(5)\cdot\zeta(3)$, $\zeta(8)$ | ham of weight 8

Encl

Cool stuff we actually know

- infinite class of such graphs s.t. residue is a MZV
- for a while there was hope for all of them, but nope (8-loop 9-loop planar)
- not every MZV seems to occur, only a specific subspace → to investigate
- All odd zeta values do appear!

Thm (Brown-Schnetz 12) conj by Broadhurst Kreimer

Zig zag graphs

Z_5

$$I_{Z_n} = \frac{4(2n-2)!}{n!(n-1)!}\left(1-\frac{1-(-1)^n}{2^{2n-3}}\right)\zeta(2n-3)$$

(here I is a residue again)

But no other family of graphs for which amplitudes are known for all #s of loops in this theory ... no conjectures

Conj (B-Schnetz)

Let G be a prim Γ in ϕ^4 ($N_G = 2h_G$) with h_G loops. Then $0 < I_G \leq I_{Z_{h_G}}$

~~work in progress~~

UNIVERSITÉ DE GENÈVE ETH zürich swissmaprs.ch Swiss National Science Foundation SwissMAP The Mathematics of Physics National Centre of Competence in Research