

GEOLOGICAL NOTE

From Cosmic Explosions to Terrestrial Fires?

Adrian L. Melott^{1,} and Brian C. Thomas²*

1. Department of Physics and Astronomy, University of Kansas, Lawrence, Kansas 66045, USA;

2. Department of Physics and Astronomy, Washburn University, Topeka, Kansas 66621, USA

ABSTRACT

Multiple lines of evidence point to one or more moderately nearby supernovae, with the strongest signal at ~2.6 Ma. We build on previous work to argue for the likelihood of cosmic ray ionization of the atmosphere and electron cascades leading to more frequent lightning and therefore an increase in nitrate deposition and wildfires. The potential exists for a large increase in the prehuman nitrate flux onto the surface, which has previously been argued to lead to CO₂ drawdown and cooling of the climate. Evidence for increased wildfires exists in an increase in soot and carbon deposits over the relevant period. The wildfires would have contributed to the transition from forest to savanna in northeast Africa, long argued to have been a factor in the evolution of hominin bipedalism.

Introduction

Astrophysical ionizing radiation events affect the earth episodically. Effects come from the sun through flares and solar proton events as well as interstellar supernovae (SNs) and gamma ray bursts (reviewed in Melott and Thomas 2011). Much of the interest regarding terrestrial effects has centered on major mass extinctions (e.g., Melott et al. 2004; Melott and Thomas 2009). Extreme events may eliminate all higher life on Earthlike planets but are unlikely to cause complete global sterilization (Sloan et al. 2017). However, less intense effects are also possible (e.g., Knipp et al. 2018), more frequent (and therefore more likely to have extant evidence), and still interesting.

There is a great deal of accumulated evidence for SNs moderately near the earth (Knie et al. 2004; Binns et al. 2016; Breitschwerdt et al. 2016; Fimiani et al. 2016; Fry et al. 2016; Ludwig et al. 2016; Mamajek 2016; Melott 2016; Wallner et al. 2016; Erlykin et al. 2017). Although the strongest signal is of an event at about 2.6 Ma, which coincides within the uncertainties to the Pliocene-Pleistocene (PP) boundary, it may well be a series beginning as long as 7 My ago related to the evacuation of the Local Bubble, a large

cavity in the interstellar medium (Breitschwerdt et al. 2016).

The possible terrestrial effect of a nearby SN has long been of interest. More recently, new information on prompt and early emissions from SNs as well as the timing and distance of relatively recent proximate SNs were used as input in a series of computations (Thomas et al. 2016; Melott et al. 2017). Although early work focused on ozone depletion from photon and cosmic ray (CR) irradiation, we have found additional increased importance for the extended arrival of CRs.

We will not repeat the description of those computations here except for a brief summary. We refer readers to Melott et al. (2019) for a summary or to Melott et al. (2017) for more detail.

Base Study Methods and Conclusions

We assumed that an SN of the most common type (IIP) took place at about 2.6 Ma at a distance of 50 pc. This is taken to be representative and possibly a member of a series. The Local Bubble had already been formed, and the magnetic field is assumed to have been largely expelled and residing within the walls of this structure, which would reflect CRs back into the interior, extending the irradiation time for the earth.

Manuscript received November 16, 2018; accepted March 3, 2019; electronically published May 28, 2019.

* Author for correspondence; email: melott@ku.edu.

[The Journal of Geology, 2019 volume 127, p. 000–000] © 2019 by The University of Chicago.
All rights reserved. 0022-1376/2019/12704-00XX\$15.00. DOI: 10.1086/703418

CR propagation in the turbulent field leads to a time-dependent flux at the earth. Here we use the CR flux for case B as computed in Melott et al. (2017) at 100 y after the arrival of the first photons and assume that magnetic reflection off the walls will later keep a similar flux inside the cavity of the Local Bubble. The flux of CRs is greatly increased beyond normal levels. For many decades, the primary effect considered for terrestrial life from a nearby SN was ozone depletion (Gehrels et al. 2003). We computed this effect based on a series of studies beginning with Thomas et al. (2005). In Melott et al. (2017), it was estimated that these events would produce a ~50% increase in ultraviolet B radiation at the surface, with a range of potential biological impacts (Thomas 2018).

Muons are highly penetrating elementary particles that penetrate up to 1 km in the ocean but, by virtue of their large numbers, still constitute the primary component of irradiation from CRs at the earth's surface. The large increase in CRs following a nearby SN leads to a similar large increase in muon flux at the earth's surface (Melott et al. 2017) and in the top kilometer of the oceans (Melott et al. 2019). Our previous work argued that a megafaunal extinction around the PP boundary (Pimienta et al. 2017) may have been induced by increased muon flux in the top few hundred meters of ocean (Melott et al. 2019).

In this short article, we discuss additional substantial effects that could also have indirectly had

a major effect on the biota, including the evolution of our species. Atmospheric ionization and dissociation from CRs and their subsequent showers of elementary particles led to the ozone depletion mentioned earlier. The ionization may also have had other effects.

Computation of Atmospheric Ionization

Atmospheric ionization was computed using tables from Atri et al. (2010), originally generated by CORSIKA (Cosmic Ray Simulations for KASCADE). CORSIKA is a package combining high- and low-energy interaction models with a specialized scheme for transport in air. The version used for atmospheric ionization was EPOS 1.61, UrQMD 1.3. This gives ionization rate (ions $\text{cm}^{-2} \text{s}^{-1}$) as a function of altitude for various primary proton energies. The tables in Atri et al. (2010) give ionization for primaries with energies from 300 MeV to 1 PeV (10^{15} eV) from the ground to 90 km. Here we used primary proton energies between 1 GeV and 1 PeV. The ionization rate as a function of altitude was created by convolving the CR flux from Melott et al. (2017) with tables from Atri et al. (2010).

In figure 1, we show the energy spectrum of galactic CRs at the present time and at 100 y after the receipt of the first photons from an SN at 50 pc (Melott et al. 2017). Although the computations had open boundary conditions, we expect that the environment




Figure 1. Approximate flux of galactic cosmic rays (GCR; dashed line) and the cosmic ray flux computed in Melott et al. (2017) at 100 y after the arrival of the photons from a supernova at 50-pc distance (SNCR; solid line). The units are such that equal areas under the lines correspond to equal total energy flux at the earth. There is an enhancement of more than two orders of magnitude at the upper range of energies.

would be reflective due to the compression of the magnetic field in the walls of the Local Bubble. Given the size of the structure, a long duration of the flux at the 100-y level would be expected. The units are such that equal areas in this plot imply equal amounts of energy deposition at the earth. Normally, high-energy CRs are too low in flux to contribute much to atmospheric ionization. Thus, besides an overall increase in CR flux, there is a unique predominance of high-energy primaries in the CR flux, up to about 1 PeV, which are normally not very important for atmospheric ionization. It is important to understand the consequences of this change.

Figure 2 shows results of our analysis of the ionization effects of CR primaries. We show the ionization flux (ions generated per area per time) per GeV of total incident CR energy as a function of altitude and primary energy. It is important to notice that the ionization rate is plotted per GeV unit of primary energy and not per primary. Thus, the expected simple increase in ionization due to more energy per primary is factored out. But an increase is still seen, particularly in the troposphere. This increase is due to the greater penetration of the high-energy CRs in the atmosphere. This will have consequences for activity in the troposphere; our primary interest here is lightning, especially cloud-to-ground lightning. An energy of 1 GeV is representative of the majority of usual galactic CR primary protons, while the 10 GeV and 10^6 GeV energies bracket the range most affected by the SN. In the lowest 2-km bin in our model, ionization caused by the higher-energy primary protons (the range associated with SN CRs) is about two orders of magnitude higher than that caused by the lower energy associated with galactic CR primaries. There is a great potential for major changes of weather near the surface.

In figure 3, we show the altitude profile of ionization from galactic CRs under normal conditions and at our fiducial time. (Energetic protons from the sun mostly do not penetrate down to the troposphere and are not relevant for the issues considered here, as discussed in Usoskin et al. [2011].) The ionization SNCR in figure 3 is computed by convolving the 100-y, case-B CR flux as seen in figure 1 with the ionization as a function of altitude and CR energy as seen in figure 2. It can be seen that there is a major difference, with ionization in the troposphere about 50 times greater than normal. For comparison, we also show our computed ionization profile from the 1956 solar superflare, the hardest and most energetic in the modern era (Usoskin et al. 2011) and therefore amenable to quantitative measurement. Consequently, the great increase in TeV to PeV CR flux on the earth from an even moderately near SN

should have strong consequences for atmospheric physics near the surface.

Initiation of Cloud-to-Ground Lightning

It has been proposed that CRs set off electron avalanches in the atmosphere that are the main initiator of lightning (Gurevich et al. 1999, 2008, 2009; Erlykin and Wolfendale 2010; Chilingarian et al. 2017a; Kumar et al. 2018). Until recently, this was an eminently reasonable idea with only circumstantial evidence in its favor. One of the obstacles has been the difficulty of making any in situ measurements of the processes initiating lightning. However, Chilingarian et al. (2017b) reported observations on Aragats Mountain, Armenia, where the cloud layer is quite low. They found a number of electron avalanches of duration less than a microsecond that were terminated by nearby lightning flashes. This is a smoking gun that makes this a compelling theory, which we will take as our working model.

We note here that it has been proposed that cloud cover is related to CR flux (Svensmark et al. 2017 and references therein). Since this proposal is highly controversial, we shall only mention it here.

A 50-fold increase in atmospheric ionization in the troposphere would clearly make the breakdown and electron cascade much easier, and one could expect a great increase in lightning (e.g., Erlykin and Wolfendale 2010). Furthermore, the originally isotropic distribution of CRs would result in showers that are much more preferentially vertical due to the variation in atmospheric column density with angle. Thus, not only would lightning be enhanced, but cloud-to-ground lightning should be preferentially enhanced. The theory of lightning initiation is not well developed, and we cannot say that a 50-fold increase in ionization would lead to a 50-fold increase in the number of lightning events. However, the potential is there for a large increase. We note that in the SN-enhanced regime, there would be high-energy primaries on the order of hundreds per square kilometer per second, so that ionization would be intermediate between continuous and pulsed.

Effects of Increased Lightning Frequency

One examined effect of increased atmospheric ionization from SN CRs (A. L. Melott, B. C. Thomas, and B. D. Fields, unpublished manuscript; see <https://arxiv.org/abs/1810.01995>) was fixation of nitrogen. Normally, the nitrogen in the atmosphere is largely unavailable to the biota due to the strength of the triple bond in N_2 . For this reason, nitrogen must be con-




Figure 2. Ionization from the high-energy cosmic rays (CRs) incident from a moderately nearby supernova ionize the atmosphere at much lower altitudes than incident CRs from any other usual source for the earth. *A*, Ionization rate (flux) per GeV of total incident CR energy as a function of altitude and primary energy. We have adjusted to constant energy input units to bring out the altitude effect due to the energy of the primary. Normal CR flux peaks near the left boundary of this plot, but moderately nearby supernovae contribute all across it. *B*, Ionization rate (flux) in the same units at 1 km altitude, the lowest we resolve. This is most relevant for cloud-to-ground lightning, which should be greatly increased by the additional ionization by high-energy CRs.

verted into other forms; there is a large literature on the synthesis and deposition of nitrate as a side effect of atmospheric ionization and dissociation (e.g., Thomas et al. 2005). Our computations suggested that the increase in available nitrogen would be relatively

small, of order 10% from the CR ionization. However, a large increase in the rate of lightning could provide a strong enhancement in the deposition of nitrate. Lightning is the dominant source of nitrogen fixation in the preindustrial world (Schlesinger and Bernhardt




Figure 3. Ionization as a function of altitude for present-day typical cosmic rays (GCR; dashed line), and the 1956 solar event (SPE; dot-dashed line), as compared with the spectrum of incident cosmic rays 100 y after the arrival of photons from a supernova at 50 pc (SNCR; solid line) as computed in Melott et al. (2017). It can be seen that ionization in the lowest levels of the atmosphere is increased by a factor of approximately 50.

2013, table 12.3). The computational methods of Thomas et al. (2005) include results from nitrate deposition, mostly in the form of rain or snow. We estimate that it would not be unreasonable for a large increase in the preindustrial, prehuman nitrogen flux from lightning to result in a similarly large, perhaps severalfold nitrate enhancement. This might lead, as discussed in A. L. Melott et al. (unpublished manuscript), to a CO₂ drawdown and cooling of the climate as was observed at the onset of the Pleistocene. We stress that this all depends on there being a large increase in the lightning frequency, which we are unable to estimate.

Lightning is the main initiator of wildfires if one excludes humans (Pausas and Keeley 2009; Veraverbeke et al. 2017). A large increase in lightning would again be expected to cause a large increase in the number of wildfires started. There has been a coincident worldwide increase in the number of wildfires since about 7 My ago (Zhou et al. 2014; Bond 2015), which is probably responsible for the conversion of forest to savanna worldwide (Karp et al. 2018). There has been no good explanation for this increase

in fires across various continents and climatic zones (Bond 2015). We suggest that a global increase in lightning might provide such an explanation.

The evidence for the increase in fires after impact events can be found in soot and other carbon-related sediments (Wolbach et al. 1985, 2018). Such evidence might be found here and is in fact cited in Zhou et al. (2014) and Bond (2015). While impact-related fires would show a sudden spike, we would expect the SN-CR wildfires to take place over a much more extended period of time: 10⁴–10⁵ y for each SN.

The conversion from woodland to savanna has long been held to be a central factor in the evolution of hominins to bipedalism, although more recent thinking tends to view it as a contributing factor (Senut et al. 2018). Thus, it is possible that nearby SNs played a role in the evolution of humans. This should be borne in mind as more research is done, particularly in the initiation of lightning.

The obvious question to many is the probability of such an event in our future. The nearest apparent type II SN progenitor is Betelgeuse, which is likely to go off sometime in the next 1 My (Meynet et al.

2013). However, at a distance of ~ 200 pc, it is not likely to cause serious consequences. We could be affected by a gamma ray burst from an unseen nearby progenitor with only a few hours' warning (Medvedev and Loeb 2013), but this is a very low-probability event. The analysis of such events is most useful as a clue to understanding the geological past. For technological humans, solar events are a more reason-

able basis for immediate concern (Melott and Thomas 2012; Knipp et al. 2018).

ACKNOWLEDGMENTS

We thank B. Lieberman and an anonymous reviewer for helpful suggestions about the presentation of this article.

REFERENCES CITED

- Atri, D.; Melott, A. L.; and Thomas, B. C. 2010. Lookup tables to compute high energy cosmic ray induced atmospheric ionization and changes in atmospheric chemistry. *J. Cosmol. Astropart. Phys.* 2010:008. doi:10.1088/1475-7516/2010/05/008.
- Binns, W. R.; Israel, M. H.; Christian, E. R.; Cummings, A. C.; de Nolfo, G. A.; Lave, K. A.; Leske, R. A.; et al. 2016. Observation of the ^{60}Fe nucleosynthesis-clock isotope in galactic cosmic rays. *Science* 352:677–680.
- Bond, W. J. 2015. Fires in the Cenozoic: a late flowering of flammable ecosystems. *Front. Plant Sci.* 5:749. doi:10.3389/fpls.2014.00749.
- Breitschwerdt, D.; Feige, J.; Schulreich, M. M.; de Avillez, M. A.; Dettbarn, C.; and Fuchs, B. 2016. The locations of recent supernovae near the sun from modelling ^{60}Fe transport. *Nature* 532:73–76.
- Chilingarian, A.; Chilingaryan, S.; Karapetyan, T.; Kozliner, L.; Khanikyants, Y.; Hovsepyan, G.; Pokhsrayan, D.; and Soghomonian, S. 2017a. On the initiation of lightning in thunderclouds. *Sci. Rep.* 7:1371. doi:10.1038/s41598-017-01288-0.
- Chilingarian, A.; Hovsepyan, G.; and Mailyan, B. 2017b. In situ measurements of the runaway breakdown (RB) on Aragats Mountain. *Nucl. Inst. Method. Phys. Res. A* 874:19–27.
- Erlykin, A. D.; Machavariani, S. K.; and Wolfendale, A. W. 2017. The Local Bubble in the interstellar medium and the origin of the low energy cosmic rays. *Adv. Space Res.* 59:748–750.
- Erlykin, A. D., and Wolfendale, A. W. 2010. Long term time variability of cosmic rays and possible relevance to the development of life on Earth. *Surv. Geophys.* 31:383–398. doi:10.1007/s10712-010-9097-8.
- Fimiani, L.; Cook, D. L.; Faestermann, T.; Gómez-Guzmán, J. M.; Hain, K.; Herzog, G.; Knie, K.; et al. 2016. Interstellar ^{60}Fe on the surface of the moon. *Phys. Rev. Lett.* 116:151104.
- Fry, B. J.; Fields, B. D.; and Ellis, J. R. 2016. Radioactive iron rain: transporting ^{60}Fe in supernova dust to the ocean floor. *Astrophys. J.* 827:48.
- Gehrels, N.; Laird, C. M.; Jackman, C. H.; and Chen, W. 2003. Ozone depletion from nearby supernovae. *Astrophys. J.* 585:1169–1176.
- Gurevich, A. V.; Karashtin, A. N.; Ryabov, V. A.; Chubenko, A. P.; and Shepetov, A. L. 2009. Non-linear phenomena in ionosphere plasma: the influence of cosmic rays and the runaway electron breakdown on the thunderstorm discharges. *Phys.-Uspekhi* 52:735–745 (in Russian).
- Gurevich, A. V.; Zybin, K. P.; and Medvedev, Y. V. 2008. Runaway breakdown in strong electric field as a source of terrestrial gamma flashes and gamma bursts in lightning leader breakdown and cosmic ray showers. *Phys. Lett. A* 254:79–87.
- Gurevich, A. V.; Zybin, K. P.; and Roussel-Dupre, R. A. 1999. Lightning initiation by simultaneous effect of runaway breakdown and cosmic-ray showers. *Phys. Lett. A* 254:79–87.
- Karp, A. T.; Behrensmeier, A. K.; and Freeman, K. H. 2018. Grassland fire ecology has roots in the late Miocene. *Proc. Natl. Acad. Sci. USA* 115:12,130–12,135.
- Knie, K.; Korschinek, G.; Faestermann, T.; Dorfi, E. A.; Rugel, G.; and Wallner, A. 2004. ^{60}Fe anomaly in a deep-sea manganese crust and implications for a nearby supernova source. *Phys. Rev. Lett.* 93:171103.
- Knipp, D. J.; Fraser, B. J.; Shea, M. A.; and Smart, D. F. 2018. On the little-known consequences of the 4 August 1972 ultra-fast coronal mass ejecta: facts, commentary, and call to action. *Space Weather* 16:1635–1653. doi:10.1029/2018SW002024.
- Kumar, S.; Singh, D.; Singh, R. P.; Singh, A. K.; and Kamra, A. K. 2018. Lightning discharges, cosmic rays, and climate. *Surv. Geophys.* 39:861–899.
- Ludwig, P.; Bishop, S.; Egli, R.; Chernenko, V.; Deneva, B.; Faestermann, T.; Famulok, N.; et al. 2016. Time-resolved 2-million-year-old supernova activity discovered in Earth's microfossil record. *Proc. Natl. Acad. Sci. USA* 113:9232–9237. doi:10.1073/pnas.1601040113.
- Mamajek, E. E. 2016. A pre-Gaia census of nearby stellar groups. *In* Kastner, J. H.; Stelzer, B.; and Metchev, S. A., eds. *Young stars and planets near the sun: proceedings of the International Astronomical Union Symposium* 314, p. 21–26. Cambridge, Cambridge University Press.
- Medvedev, M. V., and Loeb, A. 2013. On Poynting-flux-driven bubbles and shocks around merging neutron star binaries. *Mon. Not. R. Astron. Soc.* 431:2737–2744. doi:10.1093/mnras/stt366.

- Melott, A. L. 2016. Supernovae in the neighbourhood. *Nature* 532:40–41.
- Melott, A. L.; Lieberman, B. S.; Laird, C. M.; Martin, M. V.; Thomas, B. C.; Cannizzo, J. K.; Gehrels, N.; and Jackman, C. H. 2004. Did a gamma-ray burst initiate the late Ordovician mass extinction? *Int. J. Astrobiol.* 3:55–61.
- Melott, A. L.; Marinho, F.; and Paulucci, L. 2019. Muon radiation dose and marine megafaunal extinction at the end-Pliocene supernova. *Astrobiology* 19. doi:10.1089/ast.2018.1902.
- Melott, A. L., and Thomas, B. C. 2009. Late Ordovician geographic patterns of extinction compared with simulations of astrophysical ionizing radiation damage. *Paleobiology* 35:311–320. doi:10.1666/0094-8373-35.3.311.
- . 2011. Astrophysical ionizing radiation and the earth: a brief review and census of intermittent intense sources. *Astrobiology* 11:343–361. doi:10.1089/ast.2010.0603.
- . 2012. Causes of an AD 774–775 ¹⁴C increase. *Nature* 491:E1. doi:10.1038/nature11695.
- Melott, A. L.; Thomas, B. C.; Kachelrieß, M.; Semikoz, D. V.; and Overholt, A. C. 2017. A supernova at 50 pc: effects on the earth's atmosphere and biota. *Astrophys. J.* 840:105. doi:10.3847/1538-4357/aa6c57.
- Meynet, G.; Haemmerle, L.; Ekstrom, S.; and Geogy, C. 2013. The past and future evolution of a star like Betelgeuse. *Eur. Astron. Soc. Publ. Ser.* 60:17–28. doi:10.1051/eas/1360002.
- Pausas, J. G., and Keeley, J. E. 2009. A burning story: the role of fire in the history of life. *BioScience* 59:593–601. doi:10.1525/bio.2009.59.7.10.
- Pimiento, C.; Griffin, J. N.; Clements, C. F.; Silvestro, D.; Varela, S.; Uhen, M. D.; and Jaramillo, C. 2017. The Pliocene marine megafauna extinction and its impact on functional diversity. *Nat. Ecol. Evol.* 1:1100–1106.
- Schlesinger, W. H., and Bernhardt, E. S. 2013. *Biogeochemistry*. Waltham, MA, Academic Press.
- Senut, B.; Pickford, M.; Gommery, D.; and Segalen, L. 2018. Palaeoenvironments and the origin of hominid bipedalism. *Hist. Biol.* 30:284–296. doi:10.1080/08912963.2017.1286337.
- Sloan, D.; Batista, R. A.; and Loeb, A. 2017. The resilience of life to astrophysical events. *Sci. Rep.* 7:5419.
- Svensmark, H.; Enghoff, M. B.; Shaviv, N. J.; and Svensmark, J. 2017. Increased ionization supports growth of aerosols into cloud condensation nuclei. *Nat. Commun.* 8:2199.
- Thomas, B. C. 2018. Photobiological effects at Earth's surface following a 50 pc supernova. *Astrobiology* 18:481–490.
- Thomas, B. C.; Engler, E. E.; Kachelrieß, M.; Melott, A. L.; Overholt, A. C.; and Semikoz, D. V. 2016. Terrestrial effects of nearby supernovae in the early Pleistocene. *Astrophys. J. Lett.* 826:L3.
- Thomas, B. C.; Melott, A. L.; Jackman, C. H.; Laird, C. M.; Medvedev, M. V.; Stolarski, R. S.; Gehrels, N.; Cannizzo, J. K.; Hogan, D. P.; and Ejzak, L. M. 2005. Gamma-ray bursts and the earth: exploration of atmospheric, biological, climatic, and biogeochemical effects. *Astrophys. J.* 634:509–533.
- Usoskin, I. G.; Kovaltsov, G. H.; Mironova, I. A.; Tylka, A. J.; and Dietrich, W. F. 2011. Ionization effect of solar particle GLE events in low and middle atmosphere. *Atmos. Chem. Phys.* 11:1979–1988. doi:10.5194/acp-11-1979-2011.
- Veraverbeke, S.; Rogers, B. M.; Goulden, M. L.; Jandt, R. R.; Miller, C. E.; Wiggins, E. B.; and Randerson, J. T. 2017. Lightning as a major driver of recent large fire years in North American boreal forests. *Nat. Clim. Change* 7:529–534.
- Wallner, A.; Feige, J.; Kinoshita, N.; Paul, M.; Fifield, L. K.; Golser, R.; Honda, M.; et al. 2016. Recent near-Earth supernovae probed by global deposition of interstellar radioactive ⁶⁰Fe. *Nature* 532:69–72.
- Wolbach, W. S.; Ballard, J. P.; Mayewski, P. A.; Parnell, A. C.; Cahill, N.; Adedeji, V.; Bunch, T. E.; et al. 2018. Extraordinary biomass-burning episode and impact winter triggered by the Younger Dryas cosmic impact ~12,800 years ago. II. Lake, marine, and terrestrial sediments. *J. Geol.* 126:185–205.
- Wolbach, W. S.; Lewis, R. S.; and Anders, E. 1985. Cretaceous extinctions: evidence for wildfires and search for meteoritic material. *Science* 230:167–170. doi:10.1126/science.230.4722.167.
- Zhou, B.; Shen, C.; Sun, W.; Bird, M.; Ma, W.; Taylor, D.; Liu, W.; et al. 2014. Late Pliocene-Pleistocene expansion of C₄ vegetation in semiarid East Asia linked to increased burning. *Geology* 42:1067–1070. doi:10.1130/G36110.1.