




UNIVERSITAT POLITÈCNICA  
DE CATALUNYA  
BARCELONATECH



# New High Resolution Measurement of $^{56}\text{Fe}$ (n, $\gamma$ ) cross-section at n\_TOF

A. B. Allannavar<sup>1</sup>, A. Casanovas<sup>1</sup>, C. Domingo-Pardo<sup>2</sup>, F. Calviño<sup>1</sup>, G. Cortès<sup>1</sup>, C. Paradela<sup>6</sup>, M. Boromiza<sup>7</sup>,  
J. Lerendegui-Marco<sup>2</sup>, J. Balibrea-Correa<sup>2</sup>, B. Gameiro<sup>2</sup>, G. de la Fuente<sup>2</sup>, A. Tarifeño-Saldivia<sup>2</sup>, V. Alcayne<sup>3</sup>, D. Cano-Ott<sup>3</sup>, E.  
Mendoza<sup>3</sup>, C. Guerrero<sup>4</sup>, P. Pérez-Maroto<sup>4</sup>, F. García-Infantes<sup>5</sup>


On behalf of the n\_TOF Collaboration.

- <sup>1</sup> Universitat Politècnica de Catalunya (UPC), Spain
- <sup>2</sup> Instituto de Física Corpuscular (CSIC-Universitat de Valencia), Spain
- <sup>3</sup> CIEMAT, Spain
- <sup>4</sup> Universidad de Sevilla, Spain
- <sup>5</sup> Universidad de Granada, Spain
- <sup>6</sup> Joint Research Centre, Geel
- <sup>7</sup> Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest



# Motivation

- Fe – an important element for both:
  1. Structural materials in nuclear technologies
  2. Shielding
- $^{56}\text{Fe}$  – 91.754% of natural Fe
- Presence as structural material in both primary and secondary circuits.
- Treated by CIELO evaluation as one of the “Major isotopes for reactor application” among  $^1\text{H}$ ,  $^{16}\text{O}$ ,  $^{56}\text{Fe}$ ,  $^{235}\text{U}$ ,  $^{239}\text{Pu}$ ...
- Also taken up by the INDEN<sup>2</sup> project.
- The EXFOR data is from measurements from the past decades - scope for higher resolution measurements in the current times.
- FeCrAl alloy cladding explored for their high corrosion resistance.





# Motivation

- **Astrophysics** – stellar nucleosynthesis

1. Seed of the s-process
2. 6<sup>th</sup> most abundant isotope in the universe
3. Determines the neutron to seed ratio

Significance as seed nuclei motivates efforts to lower the Maxwellian Averaged Cross-section (MACS) uncertainty for He-shell (MACS at 30 keV) and C-shell (MACS at 90 keV) weak s-process nucleosynthesis

 Neutron capture  
 beta- decay

<sup>60</sup> Zn 2.38 m β <sup>+</sup>	<sup>61</sup> Zn 1.48 m β <sup>+</sup>	<sup>62</sup> Zn 9.19 h β <sup>+</sup>	<sup>63</sup> Zn 38.47 m β <sup>+</sup>	<sup>64</sup> Zn 48.63% 59 mb
<sup>59</sup> Cu 1.36 m β <sup>+</sup>	<sup>60</sup> Cu 23.70 m β <sup>+</sup>	<sup>61</sup> Cu 3.33 h β <sup>+</sup>	<sup>62</sup> Cu 9.67 m β <sup>+</sup>	<sup>63</sup> Cu 69.17% 94 mb
<sup>58</sup> Ni 68.077% 38.7 mb	<sup>59</sup> Ni 75.99 ka 87 mb, β <sup>+</sup>	<sup>60</sup> Ni 26.223% 30 mb	<sup>61</sup> Ni 1.14% 82 mb	<sup>62</sup> Ni 3.634% 22.3 mb
<sup>57</sup> Co 271.76 d β <sup>+</sup>	<sup>58</sup> Co 70.86 d β <sup>+</sup>	<sup>59</sup> Co 100% 38 mb	<sup>60</sup> Co 5.27 a β <sup>-</sup>	<sup>61</sup> Co 1.65 h β <sup>-</sup>
<sup>56</sup> Fe 91.754% 11.7 mb	<sup>57</sup> Fe 2.119% 40 mb	<sup>58</sup> Fe 0.282% 12.1 mb	<sup>59</sup> Fe 44.50 d β <sup>-</sup>	<sup>60</sup> Fe 1.50 Ma β <sup>-</sup>

# EXFOR entries

Major wide energy range capture measurement data (EXFOR):

Wang <sup>1</sup>	(2010)	: 10.0 keV – 90 keV	(Tokyo, 30 cm, 6Li-glass)
Käppeler <sup>2</sup>	(1983)	: 22.80 keV – 113 keV	(GELINA, 60 cm, C6D6)
Corvi <sup>3</sup>	(1982)	: 1.15 keV – 296 keV	(GELINA, 58 m, C6D6)
Allen <sup>4</sup>	(1976)	: 2.50 keV – 870 keV	(ORELA, 40 m, C6F6)
Macklin <sup>5</sup>	(1964)	: 14.20 keV – 59.4 keV	(ORNL, 8 cm, C6F6)

The recent evaluations are based on relatively old data ( ~4 decades old).

Reacode	nReacstr	MF	MT	Target	Reaction	Quant	nPts	enMin	enMax	year1	author1
26-FE-56(N,G)26-FE-57,,SIG,,SPA	1	3	102	Fe-56	N,G	CS	5	15000	90000	2010	T.Wang+
26-FE-56(N,G)26-FE-57,,SIG	1	3	102	Fe-56	N,G	CS	136	731	1410	1977	O.A.Shcherbakov+
26-FE-56(N,G)26-FE-57,,SIG,,AV	1	3	102	Fe-56	N,G	CS	27	1000	850000	1976	B.J.Allen+
26-FE-56(N,G)26-FE-57,,SIG,,MXW,DERIV	1	3	102	Fe-56	N,G	CS	15	5000	75000	1975	H.Beer+
26-FE-56(N,G)26-FE-57,,SIG,,TTA	1	3	102	Fe-56	N,G	CS	464	5900	42900	1970	A.Ernst+
26-FE-56(N,G)26-FE-57,,SIG,,TTA	1	3	102	Fe-56	N,G	CS	370	19100	219000	1970	A.Ernst+
26-FE-56(N,G)26-FE-57,,SIG	1	3	102	Fe-56	N,G	CS	33	14200	59400	1964	R.L.Macklin+

1. [T. Wang et. al., Nuclear Instruments and Methods in Physics Research Section B, 2.2010](#)
2. [F. Kaeppeler et. al., Nuclear Science and Engineering, 1983](#)
3. [F. Corvi et. al., ND, 1982](#)
4. [B. J. Allen et. Al., Nuclear Physics A, 1976](#)
5. [R. L. Macklin et. al., Physical Review, 1964](#)

# Objective

To disseminate to the nuclear data community a new measurement with:



1. Broad energy range (upto 500 keV) : n\_TOF white neutron source
2. High energy resolution : time-of-flight measurements with 184 m flight path
3. Good statistics : 4  $C_6D_6$  Detectors

The experimental database used in the evaluation is summarized in Table IX of the primary reference (NDS 148,241(2018)). In doing this selection we favored: (i) broad energy range covered by the experiment, (ii) high energy resolution, and (iii) high (but reasonable) accuracy of the experiment. Thus the measurements providing possibly wide and reliable picture were given precedence over single energy-point experiments, especially within the energy range where strong fluctuations were observed. On the other hand, many single- or a few-point

[A. Herman et. Al., Nuclear Data Sheets, 2018](#)



# Aspects to consider

- Challenges in resolving the 27.7keV s-wave resonance
  - Mostly dedicated to multiple scattering and prompt background - Thinner sample, Monte Carlo studies and reduced background in EAR1 phase4 target.




# Aspects to consider

- 24 keV transmission
  - Fe<sup>56</sup> transparent to neutrons of this energy - significant for shielding calculations.
  - Overlap with 27.7 keV capture resonance - call for efforts to measure this resonance more accurately.





# n\_TOF ( neutron time of flight) collaboration at CERN

- High intensity spallation neutron source produced by 20 GeV/c protons from the PS at the CERN accelerator complex.
- Lead spallation target - conversion of PS proton beam (20 GeV/c) to a white neutron (and other particles) beam source.
- Beam “cleaned” off of charged particle with a sweeping magnet.
- Spallation gammas form gamma-flash.




# Experimental areas at n\_TOF

- Experimental ARea 1 (EAR1)
  - 186 m flight path
  - Borated moderator
- Experimental ARea 2 (EAR2)
  - ~19.5m flight path
  - Non-borated moderator
  - x10 flux compared to EAR1
- NEAR experimental area
  - 5m from the lead target




# Previous Fe<sup>56</sup> measurements at n\_TOF

- 2009: Measurement of Fe<sup>56</sup> at n\_TOF
  - non-borated water moderator (phase2)
  - 2 C<sub>6</sub>D<sub>6</sub> detectors
- Successful analysis of <sup>54</sup>Fe, challenges with <sup>56</sup>Fe



Experimental setup for <sup>56</sup>Fe (2009)  
Water moderator


Experimental set-up of the Fe-56 measurement with K6D6 (n\_TOF-Ph2)



From G. Giubrone talks at n\_TOF meetings, 2010-2012

# Previous Fe<sup>56</sup> measurements at n\_TOF


- Previous <sup>56</sup>Fe measurement:
  - Significant in-beam gamma effect
  - Short distance between the sample and the detector
  - ~20% gain drift
- Deduced and implemented new measurement standards:
  - Reduced in-beam gamma background with borated moderator
  - Distance between the sample and the detectors increased to 9 cm (deduced from Cr capture studies at n\_TOF)
  - Frequent energy calibrations programmed



(Guerrero et al., EPJA, 2013)



## New measurement: July 2025 campaign

- Sample:
  - ~31.4g high-purity  $\text{Fe}^{56}$  (>99%),
  - Metallic
  - thick sample (1mm/ $8.8 \times 10^{-3}$  atoms/barn)
  - 70mm diameter: BIF = 1
- $^{\text{nat}}\text{Fe}$ , thin sample (250  $\mu\text{m}$ / $2.12 \times 10^{-3}$  atoms/barn) (multiple scattering – 27.7keV resonance.)
- $^{\text{nat}}\text{Pb}$  and  $^{\text{nat}}\text{C}$  for background determination.



Ø 70 mm x 1mm high purity  $^{56}\text{Fe}$ .  
Courtesy of C. Paradela, GELINA and A. Negret/M. Boromiza, IFIN (Romania)

## Detector set-up



- 4  $C_6D_6$  detectors at  $125^\circ$  to the beam line – low efficiency for neutron detection
- Carbon fiber shell - reduction in scattered-neutron capture in Al body
- Frequent calibration with the full range of gamma sources (4 times -  $Ba^{133}$ ,  $Cs^{137}$ ,  $Y^{88}$ , AmBe, CmC)
- Immediate measurement of  $Au^{197}$  following calibration

# The campaign: protons and the beam


- Month-long beam-time at EAR1
- Thinner <sup>nat</sup>Fe sample: Multiple scattering effects
- Au<sup>197</sup>: normalization
- Empty: Beam on background
- Measurements with high energy neutron filters
- Stable beam, frequent calibrations.
- Measurement of <sup>nat</sup>Pb, <sup>nat</sup>C for background effects

Thanks to high-intensity beam from the PS, the campaign surpassed the planned number of protons.

<b>Capture</b>			
	<b>Planned</b>	<b>Measured</b>	<b>%</b>
56Fe	1.40E+18	1.56E+18	111.77%
natFe	6.00E+17	5.18E+17	86.26%
Au	1.00E+17	8.13E+16	81.29%
Ag	5.00E+16	2.01E+16	40.22%
Empty	2.00E+17	1.814E+17	90.69%
56Fe Al filters	1.50E+17	1.50E+17	99.83%
56Fe S filter	1.50E+17	1.37E+17	91.60%
56Fe Bi filter	1.50E+17	1.65E+17	109.68%
Empty Filter S	5.00E+16	1.20E+17	240.88%
Empty Filter Bi	5.00E+16	6.87E+16	137.43%
Empty Al filters	5.00E+16	8.35E+16	166.92%
natC	1.00E+17	9.95E+16	99.52%
natPb	1.00E+17	8.57E+16	85.67%
<b>Total</b>	<b>3.15E+18</b>	<b>3.27E+18</b>	<b>103.96%</b>
<b>Total proposal</b>	<b>3.00E+18</b>		

# First look at the capture data: Comparing structures to INDEN

Comparing the experimental structures with INDEN




- Non-relativistic approximation for tof-energy conversion.
- Deposited energy threshold - 140 keV
- Calibration with sources and Geant4 simulation of energy deposition in the detectors.
- Fe<sup>56</sup> structures clearly recognizable.
- Experimental empty background scaled to 1.8x
- Limiting to comparison of structures for the preliminary look.

*Preliminary calculations*

# First look at the capture data: the 1.15 keV resonance

- Experimental background added to the SAMMY counting rate estimate of INDEN data for a first look at the structures.
- First resonance at 1.15 keV clearly marked and in agreement with the evaluation.

Comparing the experimental structures with INDEN




*Preliminary calculations*

# First look at the capture data: Around the 27.7 keV s-wave

- Resonances between 10 keV and 50 keV.
- s-wave and other nearby resonance positions in agreement with INDEN.

Comparing the experimental structures with INDEN




Preliminary calculations

# First look at the capture data: Around the 27.7 keV s-wave

- Resonances between 50 keV and 100 keV.

Comparing the experimental structures with INDEN




Preliminary calculations

# First look at the capture data: Around the 27.7 keV s-wave

- Resonances between 100 keV and 500 keV.
- A first look shows a general good agreement.

Comparing the experimental structures with INDEN




Preliminary calculations

# First look at the capture data: Other resonances (upto 500keV)

- Resonances in 300 keV - 850 keV.
- Plan to explore higher energy deposition threshold to obtain some average cross-section above 500 keV.

Comparing the experimental structures with INDEN



Preliminary calculations

# Conclusions

- Measurement of Fe56 (n, gamma) reaction at CERN's n\_tof with high a purity sample.
- Analysis to produce resolved resonances upto 500 keV; average cross-sections beyond 500 keV.
- Initial investigation into the structures of resonances in counts reveal good agreement with the latest evaluated resonance structures.





# Thank you for your attention



Grant PID2022-138297NB-C22  
funded by  
MCIN/AEI/10.13039/50110001  
1033 and by "ERDF A way  
of making Europe"

Funded by the European Union  
NextGenerationEU



- $^{197}\text{Au}$  measurements distributed along the campaign
- Det #1



- $^{197}\text{Au}$  measurements distributed along the campaign
- Det #2



- $^{197}\text{Au}$  measurements distributed along the campaign
- Det #3



- $^{197}\text{Au}$  measurements distributed along the campaign
- Det #4

