

Direct Detection of Dark Matter: Status and Prospects








PASCOS 2026 in Sheffield
22nd June 2026

Dr Sally Shaw
sally.shaw@ed.ac.uk



Outline:

- ▶ **Big Questions**
- ▶ **DM Candidates**
- ▶ **Delve Deep**
- ▶ **Search Wide**
- ▶ **Conclusion**



The Universe



Dark Matter Big Questions

What is the nature of dark matter?

... that's kind of it.


But, in more detail:

- ▶ Is it a particle?
- ▶ What are its properties?
- ▶ How does it interact?
- ▶ Is there one particle or many (a dark sector)?
- ▶ How does it explain astrophysical observations?




Particle Dark Matter

- ▶ Leading cosmology model is Λ CDM.
- ▶ CDM = cold dark matter, has to be **non-baryonic, neutral, stable** (or $\tau \gg$ age of universe).
- ▶ Theoretically well-motivated is the Weakly Interacting Massive Particle (WIMP).
- ▶ **WIMP miracle**: measured relic abundance of DM \rightarrow annihilation cross section $O(\text{weak interaction})$
- ▶ WIMPs expected to couple to the SM: *WIMP-nucleon scattering*
- ▶ Lightest supersymmetric particle (LSP) from Supersymmetric models is an ideal candidate (e.g. neutralino)




3 ways to detect dark matter particles

Enabling discovery with a multi-faceted approach





3 ways to detect dark matter particles

Enabling discovery with a multi-faceted approach




Direct Detection

Dark matter exists in a **halo** extending far beyond the visible boundaries of the galaxy



Dark matter follows a **velocity distribution** within the halo (in the Galactic frame) with a maximum velocity of v_{esc}

In Earth's frame, see an **annual modulation**. DM has a higher relative v in June than Dec
 → more likely to be above threshold
 → peak rate in June, minimum in Dec




Direct Detection

detector
exposure


particle + nuclear
physics

astrophysics

$$\frac{dN}{dE_R} = \epsilon \frac{\rho}{2m_\chi \mu_{\chi N}} \sigma_0 F^2(E_R) \int_{v_{min}} \frac{f(\vec{v})}{v} d^3v$$



DM ν translated to recoil energy in DM-scattering, get an **exponentially falling rate**
 → threshold very important



WIMP Dark Matter

An undergrad student I taught:

*“Given this paper ‘Supersymmetric Dark Matter’ by Jungman, Kamionkowski and Griest is **older than me**, I’m wondering if the current state of the field is one that still favours WIMPs as one of the main focuses of detection efforts.”*

WIMP Dark Matter

FORBES > INNOVATION > SCIENCE

The 'WIMP Miracle' Hope For Dark Matter Is Dead

Ethan Siegel Senior Contributor
Starts With A Bang Contributor Group ⓘ

Follow

Feb 22, 2019, 02:00am EST

*“Given the
Kamionka
the current*


*ungman,
dering if
WIMPs as*

WIMP Dark Matter


Simplest models ruled out but WIMP models survive through more complex diagrams & effective field theories

Dark Matter candidates circa 2026: the WIMPs and beyond the WIMPs
Marco Cirelli - IDM 2026


SM weak scale SI interactions




tree level,
vector




tree level,
scalar



one loop



M. Cirelli, A. Strumia, J. Zupan 'Dark Matter', 2406.01705



“Most of the surviving CMSSM parameter space features a **Higgsino-like dark matter particle** with a mass **~1000–1100 GeV**, which could best be probed with future direct searches for dark matter scattering.”


The CMSSM Survives Planck, the LHC, LUX-ZEPLIN, Fermi-LAT, H.E.S.S. and IceCube
Ellis, Olive, Spanos, Stamou
arXiv:2210.16337

WIMP Dark Matter


Simplest models ruled out but WIMP models survive through more complex

Dark Matter candidates circ
Mar


SM weak sc



tree level,
vector




tree level,
scalar



one loop




M. Cirelli, A. Strumia, J. Zupan 'Dark Matter', 2406.01705



CMSSM parameter space
like dark matter particle
100 GeV, which could best
the direct searches for dark
scattering.”

LHC, LUX-ZEPLIN, Fermi-LAT, H.E.S.S.
and IceCube
e, Spanos, Stamou
arXiv:2210.16337

WIMP-Nucleon Cross Section Evolution




R.J.Gaitskell, "Direct Detection of Dark Matter", Annu. Rev. Nucl. and Part. Sci. 54 (2004) 315-359 (updated by R.J. Gaitskell et al., Private Communication, 2025)


Dark Matter Candidates

- ▶ Many, many candidates beyond “vanilla WIMPs”...
- ▶ Experiments to detect DM have an enormous range of masses to cover
- ▶ Current experiments are **not sensitive** to many candidates at either ends of the mass range
- ▶ Need to “*delve deep and search wide*”

Report of the Topical Group on Particle Dark Matter
for Snowmass 2021
[arXiv:2209.07426](https://arxiv.org/abs/2209.07426)




Delve Deep & Search Wide




Delve Deep & Search Wide

Low Threshold
Phonon detectors,
SPCs, CCDs,
optimized TPCs






More Exposure
Larger targets,
longer running
(LXe, LAr)

Delve Deep




LZ

- ▶ Dual-phase LXe TPC, 7T target at Sanford Underground Research Facility (SURF) in South Dakota
- ▶ 17T GdLS Outer Detector & 2T instrumented LXe skin
- ▶ Operating since 2021, 2025 results used 4.2 tonne-year exposure
- ▶ World-leading WIMP limits above 9 GeV






Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment
J. Aalbers et al, Phys. Rev. Lett. 135, 011802 (2025)



LZ

- ▶ New results focus on low energy region of interest 1-6 keVee
- ▶ First evidence (4.5σ) of coherent neutrino-nuclear scattering (CE ν NS) on Xe from Boron-8 solar neutrinos
- ▶ New low mass constraints between 3 - 9 GeV
- ▶ LZ continues to operate and take data towards a 1000 day dataset by end of 2027




Searches for Light Dark Matter and Evidence of Coherent Elastic Neutrino-Nucleus Scattering of Solar Neutrinos with the LZ Experiment
 D.S. Akerib et al, *aeXiv:2512.08065*



XENONnT

- ▶ Dual-phase LXe TPC, 5.9 T active located at LNGS
- ▶ Gd-water neutron veto
- ▶ Recent results include an ionisation-only search (“S2-only”) & constraints on light DM
- ▶ Machine learning techniques used to remove backgrounds from delayed electrons, accidental electrons and Pb-210 on cathode



Light Dark Matter Search with 7.8 Tonne-Year of Ionization-Only Data in XENONnT
 Aprile, E. et al, arXiv:2601.11296



PandaX

- ▶ PandaX-4T has completed its full physics program
 - ▶ ~700 live days collected
 - ▶ Calibration techniques tested (DD/DT pulsed neutrons, Ar37)
- ▶ PandaX-20T fully funded, all Xe acquired
 - ▶ Staged plan, eventually 47-tonne Xe with 40-tonne sensitive volume
 - ▶ Cold LS veto outside Cu vessel
- ▶ Timeline:
 - ▶ 2026: 4T decommissioning
 - ▶ 2027: 20T installation & commissioning
 - ▶ 2028: 20T physics operation

PandaX-20T: the next generation xenon detector for dark matter and neutrino searches
Xiang Xiao, IDM 2026





DarkSide-20K

Status of the DarkSide-20k experiment

Vicente Pesudo, IDM 2026

- ▶ Dual-phase underground Argon (UAr) TPC
- ▶ UAr inner veto, atm. Ar outer veto
- ▶ Argon lighter than LXe - **lower threshold**, pulse shape discrimination for ER/NR in addition to S1/S2
- ▶ **DarkSide-20k** construction at Gran Sasso Laboratory (LNGS) advanced
- ▶ Cryostat and internal crane installed
- ▶ Underground Ar cryogenics system to be upgraded by the end of the summer
- ▶ Detector will be ready for filling end of 2028




XLZD

XENON, DARWIN and LZ have joined forces to build a 40T-80T LXe TPC.

Science drivers:



- ▶ The **ultimate probe of WIMPs** down to the neutrino fog
- ▶ A **competitive and economic** search for neutrinoless double-beta decay
- ▶ Measurements of multiple astrophysical neutrino signals



XLZD

- ▶ Despite P5 recommendation in 2023, in Dec 2025 DOE HEP announces deferring of G3DM until early 2030's due to funding constraints
- ▶ International partners working towards CDR for early 2027
- ▶ 11 Canadian institutions recently joined XLZD
- ▶ Technical evaluation of 4 sites to complete in the coming months: **Boulby** (UK), **LNGS** (Italy), **SURF** (USA), **SNOLAB** (Canada)
- ▶ XLZD planned to be the definitive search for medium to high WIMP masses - 3σ discovery potential at SI cross section of $3 \times 10^{-49} \text{ cm}^2$ at 40 GeV

The XLZD Design Book: Towards the Next-Generation Liquid Xenon Observatory for Dark Matter and Neutrino Physics
 Aalbers, J. et al, Eur. Phys. J. C (2025) 85: 1192



Annual Modulation



DAMA/LIBRA claim 13.7σ annual modulation signal in 2-6 keV window in NaI(Tl) target.

Suite of experiments running to test this in the same and different targets, N/S hemisphere, different vetos etc.

COSINE-100 recent results with same target, NaI(Tl):

- ▶ 6-year data modulation analysis ruled out DAMA/LIBRA's claim with 3.25σ .
- ▶ Combined data with ANAIS-112 also disfavors the DAMA/LIBRA signal with 4.7σ .
- ▶ 3-year data event selection lowered threshold to 0.7 keV; SI limit excludes the DAMA/LIBRA contours


Further results from DAMA/LIBRA-phase2 and perspectives
R. Bernabei et al., Nucl. Phys. At. Energy 22, 329 (2021)



Status of the dark matter search in the COSINE-100 experiment
Jinyoung Kim- IDM 2026

Search Wide


Low Threshold
Phonon detectors,
SPCs, CCDs,
optimized TPCs



Low Mass DM


“The landscape of low-threshold dark matter direct detection in the next decade”
 Snowmass2021 Cosmic Frontier
 arXiv:2203.08297

- ▶ Light (< 1 GeV) DM invokes **technological challenges** for detection
- ▶ Interaction is mix of DM-nucleus, scattering DM-e scattering and both depending on model
- ▶ Need **different technology** - lots of small scale experiments currently probing the low mass regime & proposals for upgrades for larger experiments e.g. H-doping in LZ (HydroX)



This is by no means an exhaustive list of light DM candidates....

Candidate	Light WIMPs	Solar Axion	ALPs	Sterile ν	Hidden Photons
Mass	sub-GeV	$\mu\text{eV} - \text{meV}$	$10^{-11} - 10^3$ eV	keV	1 keV - 10 GeV
Detection Methods	ionisation-only searches, phonons, Migdal effect, doping of LXe with light elements	Axio-electric effect (ER spectrum), conversion to γ in microwave cavities	Axio-electric effect (mono energetic ER), conversion to γ in microwave cavities	Decay to ν through active/sterile mixing	Dark photo-electric effect, decay products



Cryogenic Detectors

▶ SNOLAB: SuperCDMS



- ▶ Target materials: Si (0.6 kg), Ge (1.4 kg)
- ▶ Two types of detector: Interleaved Z-sensitive Ionization and Phonon (iZIP) & High Voltage
- ▶ First pulses Jan 15, 2026 !

▶ LNGS: CRESST-III, BULLKID-DM, COSINUS


- ▶ CREST-III: Targets - CaWO₄, Si, LiAlO₂, Al₂O₃ (sapphire) and Si-On-Sapphire (SOS)
- ▶ BULLKID: 800g Si
- ▶ COSINUS: 34g NaI, cold DAMA check, taking data 2026-2027

▶ LSM: TESSERACT

- ▶ Ge/Si and superfluid He targets, TES-based sensors (QETs). Operations 2029-2031




“Direct searches for light dark matter with cryogenic experiments and the low-energy excess”
 Belina Von Krosigk, IDM 2026





Spherical Proportional Counters

NEWS-G: Sub-GeV DM detection with Spherical Proportional Counters

- ▶ Detects ionisation electrons drifted to an anode
- ▶ Easy to swap target gas - check different nuclei
- ▶ NEWS-G @ LSM recently set strongest constraints on spin-dependent WIMP-proton (directly on H) cross-section in the 0.18 – 1.2 GeV range
- ▶ SNOLAB run with Ne-CH₄, CH₄ and He-CH₄ - analysis underway
- ▶ Next: DarkSphere - 300cm, potentially at Boulby




"Searching for light dark matter with DarkSPHERE"
L. Milligan, IDM 2026




Directional Detection

Direct DM experiments ARE now detecting CE ν NS from astrophysics neutrinos - will limit DM sensitivity



**The neutrino fog is defined to be the regime for which $n > 2$, with the neutrino “floor” being the largest cross section for each mass where this transition occurs*



Directionality allows clear discrimination of neutrinos (position tracks the sun) from DM (always from Cygnus)

*Direct Detection Limits
Ciaran O'Hare*

CYGNUS


- ▶ Long term CYGNUS Vision: **Multi-site Galactic Recoil Observatory** with directional sensitivity to WIMPs and neutrinos using low-density gas TPCs. Targeting low-mass WIMP region (~ 10 keV) with directionality
- ▶ Prototyping in various stages across the world

e.g. CYGNO-04 at LNGS - 0.4m³ demonstrator

PHASE 1:
O(1) m³ demonstrator

2024/26
LNF/LNGS

CYGNO_04




- background
- material tests, gas purification
- scalability


PHASE 2:
30 m³ experiment


2027 ...
LNGS

CYGNO_30




- Physics research





Commissioning and design of the CYGNO-04 detector at LNGS

S. Piacentini, IDM 2026




Honourable Mentions

Widening the search through novel ideas!


Death by dark matter

Death and Serious Injury from Dark Matter
 J. S. Sidhu, R. Scherrer and G. Starkman
 Phys. Let. B, **803**, 135300, (2020)




DNA detectors

New Dark Matter Detectors using DNA or RNA for Nanometer Tracking
 A. Drukier et al
 arXiv:1206.6809




Paleodetectors

Projected Sensitivity of Paleo-Detectors to Dark Matter Effective Interactions with Nuclei
 D. P. Theodosopoulos, et al
 Phys. Rev. D **113**, 123044 (2026)




Conclusions



R.J.Gaitskell, "Direct Detection of Dark Matter", Annu. Rev. Nucl. and Part. Sci. 54 (2004) 315-359 (updated by R.J. Gaitskell et al., Private Communication, 2025)


Conclusions



R.J.Gaitskell, "Direct Detection of Dark Matter", Annu. Rev. Nucl. and Part. Sci. 54 (2004) 315-359 (updated by R.J. Gaitskell et al., Private Communication, 2025)

Conclusions

- ▶ The search for dark matter continues, and grows broader...
- ▶ Interesting era is beginning - we have started to measure the **neutrino fog**
 - ▶ This eventually limits standard direct detection techniques
- ▶ Dual focus on increasing exposure and increasing low mass sensitivity - **delve deep and search wide**
 - ▶ Multiple targets in Direct searches can probe a **diverse set of DM models** and **wide range of DM properties**
- ▶ Still chance for **DM discovery** with running and planned experiments - stay tuned!



Conclusions

- ▶ The search for dark matter continues, and grows broader...
- ▶ Interesting era is beginning - we have started to measure the **neutrino fog**
 - ▶ This eventually limits standard direct detection techniques
- ▶ Dual focus on increasing exposure and increasing low mass sensitivity - **delve deep and search wide**
 - ▶ Multiple targets in Direct searches can probe a **diverse set of DM models** and **wide range of DM properties**
- ▶ Still chance for **DM discovery** with running and planned experiments - stay tuned!

