

# State-Driven Fragmentation of All-Heavy and Hidden-Heavy Tetraquarks

**Francesco Giovanni Celiberto, UAH Madrid**



NSTAR 2026

 Sevilla - 08/09/2026 

[Phys. Rev. D Letters 111 (2025) 11, L111501 [arXiv:2504.03949]] 

[Phys. Rev. D 113 (2026) 5, 054014 [arXiv:2511.18984]] 

[Phys. Rev. D 113 (2026) 11, 114037 [arXiv:2604.11646]] 

+ Work in Progress [TQ4Qhf1.0]





QCD Hydrogen Atoms

# Why studying (Heavy) Exotics ?




JETHAD

v0.5.5

Precision & Exploration







# ¿ What is an Exotic Hadron ?

-  Baryons & Mesons → composite states →  $|q_1 q_2 q_3\rangle$  &  $|q_1 \bar{q}_2\rangle$
-  Ordinary Hadrons ← simplest valence-quark color-singlet states
-  QCD color neutrality → TetraQuarks, PentaQuarks, Hybrids allowed







# ¿ What is an Exotic Hadron ?

-  **Baryons** & **Mesons** → composite states →  $|q_1 q_2 q_3\rangle$  &  $|q_1 \bar{q}_2\rangle$
-  **Ordinary Hadrons** ← simplest valence-quark color-singlet states
-  QCD color neutrality → **TetraQuarks**, **PentaQuarks**, **Hybrids** allowed
-  **Exotic Hadrons** ← states beyond  $|q_1 q_2 q_3\rangle$  or  $|q_1 \bar{q}_2\rangle$







# ¿ What is an Exotic Hadron ?

-  Baryons & Mesons → composite states →  $|q_1 q_2 q_3\rangle$  &  $|q_1 \bar{q}_2\rangle$
-  Ordinary Hadrons ← simplest valence-quark color-singlet states
-  QCD color neutrality → TetraQuarks, PentaQuarks, Hybrids allowed
-  Exotic Hadrons ← states beyond  $|q_1 q_2 q_3\rangle$  or  $|q_1 \bar{q}_2\rangle$
-  GENUINE EXOTICS → forbidden  $J^{PC}$  or manifest multiquarks and/or valence gluons








# ¿ What is an Exotic Hadron ?

-  Baryons & Mesons → composite states →  $|q_1 q_2 q_3\rangle$  &  $|q_1 \bar{q}_2\rangle$
-  Ordinary Hadrons ← simplest valence-quark color-singlet states
-  QCD color neutrality → TetraQuarks, PentaQuarks, Hybrids allowed
-  Exotic Hadrons ← states beyond  $|q_1 q_2 q_3\rangle$  or  $|q_1 \bar{q}_2\rangle$ 
  - GENUINE EXOTICS → forbidden  $J^{PC}$  or manifest multiquarks and/or valence gluons
  - CRYPTO-EXOTICS → ordinary quantum numbers, but exotic internal structure



# ¿ What is an Exotic Hadron ?

-  Baryons & Mesons → composite states →  $|q_1 q_2 q_3\rangle$  &  $|q_1 \bar{q}_2\rangle$
-  Ordinary Hadrons ← simplest valence-quark color-singlet states
-  QCD color neutrality → TetraQuarks, PentaQuarks, Hybrids allowed
-  Exotic Hadrons ← states beyond  $|q_1 q_2 q_3\rangle$  or  $|q_1 \bar{q}_2\rangle$ 
  -  GENUINE EXOTICS → forbidden  $J^{PC}$  or manifest multiquarks and/or valence gluons
  -  CRYPTO-EXOTICS → ordinary quantum numbers, but exotic internal structure
-  First discovery of a (crypto-)exotic at Belle (🇯🇵) in 2003 →  $X(3872)$



# Interpretation of Exotic States

## Compact MultiQuark





# Interpretation of Exotic States

Compact MultiQuark



Hybrid



# Interpretation of Exotic States

diQuark + AntidiQuark

Compact MultiQuark



Hybrid



# Interpretation of Exotic States

diQuark + AntidiQuark

Compact MultiQuark



Hybrid



Molecule of Mesons



# Interpretation of Exotic States


diQuark + AntidiQuark

HadroQuarkonium

Compact MultiQuark



Hybrid



Molecule of Mesons



# All-Charm TetraQuarks: Experimental Highlights

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CMS-BPH-24-002



CERN-EP-2025-118  
2025/06/10

## Determination of the spin and parity of all-charm tetraquarks

The CMS Collaboration\*

### Abstract

The traditional quark model accounts for the existence of baryons, such as protons and neutrons, which consist of three quarks, as well as mesons, composed of a quark-antiquark pair. Only recently has substantial evidence started to accumulate for exotic states composed of four or five quarks and antiquarks. The exact nature of their internal structure remains uncertain. This paper reports the first measurement of quantum numbers of the recently discovered family of three all-charm tetraquarks, using data collected by the CMS experiment at the Large Hadron Collider from 2016 to 2018. The angular analysis techniques developed for the discovery and characterization of the Higgs boson have been applied to the new exotic states. The quantum numbers for parity  $P$  and charge conjugation  $C$  symmetries are found to be  $+1$ . The spin  $J$  of these exotic states is consistent with  $2\hbar$ , while  $0\hbar$  and  $1\hbar$  are excluded at 95% and 99% confidence level, respectively. The  $J^{PC} = 2^{++}$  assignment implies particular configurations of constituent spins and orbital angular momenta, which constrain the possible internal structure of these tetraquarks.

Submitted to Nature





# All-Charm TetraQuarks: Experimental Highlights

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CMS-BPH-24-002



CERN-EP-2025-118  
2025/06/10



## Determination of the spin and parity of all-charm tetraquarks

The CMS Collaboration\*

### Abstract

The traditional quark model accounts for the existence of baryons, such as protons and neutrons, which consist of three quarks, as well as mesons, composed of a quark-antiquark pair. Only recently has substantial evidence started to accumulate for exotic states composed of four or five quarks and antiquarks. The exact nature of their internal structure remains uncertain. This paper reports the first measurement of quantum numbers of the recently discovered family of three all-charm tetraquarks, using data collected by the CMS experiment at the Large Hadron Collider from 2016 to 2018. The angular analysis techniques developed for the discovery and characterization of the Higgs boson have been applied to the new exotic states. The quantum numbers for parity  $P$  and charge conjugation  $C$  symmetries are found to be  $+1$ . The spin  $J$  of these exotic states is consistent with  $2\hbar$ , while  $0\hbar$  and  $1\hbar$  are excluded at 95% and 99% confidence level, respectively. The  $J^{PC} = 2^{++}$  assignment implies particular configurations of constituent spins and orbital angular momenta, which constrain the possible internal structure of these tetraquarks.

Submitted to Nature





# All-Charm TetraQuarks: Experimental Highlights

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CMS-BPH-24-002



CERN-EP-2025-118  
2025/06/10




## Determination of the spin and parity of all-charm tetraquarks

The CMS Collaboration\*

### Abstract

The traditional quark model accounts for the existence of baryons, such as protons and neutrons, which consist of three quarks, as well as mesons, composed of a quark-antiquark pair. Only recently has substantial evidence started to accumulate for exotic states composed of four or five quarks and antiquarks. The exact nature of their internal structure remains uncertain. This paper reports the first measurement of quantum numbers of the recently discovered family of three all-charm tetraquarks, using data collected by the CMS experiment at the Large Hadron Collider from 2016 to 2018. The angular analysis techniques developed for the discovery and characterization of the Higgs boson have been applied to the new exotic states. The quantum numbers for parity  $P$  and charge conjugation  $C$  symmetries are found to be  $+1$ . The spin  $J$  of these exotic states is consistent with  $2\hbar$ , while  $0\hbar$  and  $1\hbar$  are excluded at 95% and 99% confidence level, respectively. The  $J^{PC} = 2^{++}$  assignment implies particular configurations of constituent spins and orbital angular momenta, which constrain the possible internal structure of these tetraquarks.

Submitted to Nature





## Article

# Determination of the spin and parity of all-charm tetraquarks

<https://doi.org/10.1038/s41586-025-09711-7>

Received: 9 June 2025

Accepted: 6 October 2025

Published online: 3 December 2025

Open access

 Check for updates

The CMS Collaboration\*✉

The traditional quark model<sup>1,2</sup> accounts for the existence of baryons, such as protons and neutrons, which consist of three quarks, as well as mesons, composed of a quark–antiquark pair. Only recently has substantial evidence started to accumulate for exotic states composed of four or five quarks and antiquarks<sup>3</sup>. The exact nature of their internal structure remains uncertain<sup>4–29</sup>. Here we report the first measurement of quantum numbers of the recently discovered family of three all-charm tetraquarks<sup>30–32</sup>, using data collected by the CMS experiment at the Large Hadron Collider from 2016 to 2018 (refs. 33,34). The angular analysis techniques developed for the discovery and characterization of the Higgs boson<sup>35–37</sup> have been applied to the new exotic states. Here we show that the quantum numbers for parity  $P$  and charge conjugation  $C$  symmetries are found to be  $+1$ . The spin  $J$  of these exotic states is determined to be consistent with  $2\hbar$ , while  $0\hbar$  and  $1\hbar$  are excluded at 95% and 99% confidence levels, respectively. The  $J^{PC} = 2^{++}$  assignment implies particular configurations of constituent spins and orbital angular momenta, which constrain the possible internal structure of these tetraquarks.





# Why Fragmentation is so relevant ?


JETHAD

v0.5.5



# Fragmentation in QCD (with Heavy Flavors)

PRECISION  
CONTROL





DISCOVERY  
POTENTIAL



# Fragmentation in QCD (with Heavy Flavors)

PRECISION  
CONTROL



DISCOVERY  
POTENTIAL



# Fragmentation in QCD (with Heavy Flavors)

PRECISION  
CONTROL

## Light hadron collinear fragmentation

Partonic  
hard factor

DGLAP A0evo  
All-Order evolution (numeric)

Initial-scale input

NP FF

! PRECISION QCD  
+ HADRON STRUCTURE  
INTERPLAY NEEDED !

SDCs  
NP FF

$$\mathcal{D}_i^Q(z, [n]) \otimes \mathcal{D}_i^{\text{NP}}(z, Q_0)$$

Energy

Evolution-ready scale  $Q_0 \equiv M_H$

Time

DISCOVERY  
POTENTIAL



# All-Heavy-TetraQ NonRelativistic Fragmentation

JETHAD  
v0.5.5

2



# Heavy-Flavor Non-Relativistic evolution

**HADRONIC STRUCTURE**

**QUARKONIUM THEORY**



# Heavy-Flavor Non-Relativistic evolution

**HADRONIC STRUCTURE**

**QUARKONIUM THEORY**

**PRECISION QCD**

**COLLINEAR FACTORIZATION**



# Heavy-Flavor Non-Relativistic evolution

**HADRONIC STRUCTURE**  
**QUARKONIUM THEORY**

**PRECISION QCD**  
**COLLINEAR FACTORIZATION**

Guiding principle  $\Rightarrow$  ; Use the Best of the Two Worlds as much as we can !

**Interpretation**



# Heavy-Flavor Non-Relativistic evolution

**HADRONIC STRUCTURE**  
**QUARKONIUM THEORY**

**PRECISION QCD**  
**COLLINEAR FACTORIZATION**

Guiding principle  $\Rightarrow$  ; Use the Best of the Two Worlds as much as we can !

**Interpretation**

**Evolution**



# Heavy-Flavor Non-Relativistic evolution

**HADRONIC STRUCTURE**  
**QUARKONIUM THEORY**

**PRECISION QCD**  
**COLLINEAR FACTORIZATION**

Guiding principle  $\Rightarrow$  ; Use the [Best of the Two Worlds](#) as much as we can !

**Interpretation**

**Evolution**


**Uncertainties**



# TetraQuark Fragmentation from NRQCD & symJETHAD

LO

Gluon




 [F. Feng, Y. Huang et al.,  
Phys. Rev. D 106 (2022) 11, 114029]



# TetraQuark Fragmentation from NRQCD & symJETHAD


LO

Gluon



[F. Feng, Y. Huang et al.,  
Phys. Rev. D 106 (2022) 11, 114029]

Constituent  
heavy quark




[X.-W. Bai, F. Feng et al.,  
JHEP 09 (2024) 002]



# TetraQuark Fragmentation from NRQCD & symJETHAD


LO

Gluon




[F. Feng, Y. Huang et al.,  
Phys. Rev. D 106 (2022) 11, 114029]

Constituent  
heavy quark



[X.-W. Bai, F. Feng et al.,  
JHEP 09 (2024) 002]

Non-constituent  
heavy or light quark




[X.-W. Bai, Y. Huang et al.,  
Phys. Rev. D 111 (2025) 5, 054006]



# TetraQuark Fragmentation from NRQCD & symJETHAD


LO

Gluon




[F. Feng, Y. Huang et al.,  
Phys. Rev. D 106 (2022) 11, 114029]

Constituent  
heavy quark



[X.-W. Bai, F. Feng et al.,  
JHEP 09 (2024) 002]

Non-constituent  
heavy or light quark



[X.-W. Bai, Y. Huang et al.,  
Phys. Rev. D 111 (2025) 5, 054006]

$$J^{PC} = 0^{++}$$

$$J^{PC} = 1^{+-}$$

$$J^{PC} = 2^{++}$$




# TetraQuark Fragmentation from NRQCD & symJETHAD

LO

2024


TQ4Q1.0

Gluon




[F. Feng, Y. Huang et al., Phys. Rev. D 106 (2022) 11, 114029]

Constituent heavy quark



[X.-W. Bai, F. Feng et al., JHEP 09 (2024) 002]

Non-constituent heavy or light quark



[X.-W. Bai, Y. Huang et al., Phys. Rev. D 111 (2025) 5, 054006]

$J^{PC} = 0^{++}$

TQ4Q1.0+ ✓

! Suzuki [TQ4Q1.0]

! Not implemented

$J^{PC} = 1^{+-}$

Landau-Yang  
LO: evolution only ✗

! Not implemented

C-parity  
LO: evolution only ✗

$J^{PC} = 2^{++}$

TQ4Q1.0+ ✓

! Suzuki [TQ4Q1.0]

! Not implemented



# TetraQuark Fragmentation from NRQCD & symJETHAD

LO

2024




2025

Gluon


Constituent heavy quark

Non-constituent heavy or light quark


TQ4Q1.1



[F. Feng, Y. Huang et al., Phys. Rev. D 106 (2022) 11, 114029]



[X.-W. Bai, F. Feng et al., JHEP 09 (2024) 002]



[X.-W. Bai, Y. Huang et al., Phys. Rev. D 111 (2025) 5, 054006]

$J^{PC} = 0^{++}$

TQ4Q1.0+ ✓

TQ4Q1.1+ ✓

! Not implemented

$J^{PC} = 1^{+-}$

Landau-Yang  
LO: evolution only ✗

TQ4Q1.1+ ✓

C-parity  
LO: evolution only ✗

$J^{PC} = 2^{++}$

TQ4Q1.0+ ✓

TQ4Q1.1+ ✓

! Not implemented



# TetraQuark Fragmentation from NRQCD & symJETHAD

LO

2024



2025



2026


Gluon




Constituent heavy quark

Non-constituent heavy or light quark


TQ4Q2.0



[F. Feng, Y. Huang et al., Phys. Rev. D 106 (2022) 11, 114029]



[X.-W. Bai, F. Feng et al., JHEP 09 (2024) 002]



[X.-W. Bai, Y. Huang et al., Phys. Rev. D 111 (2025) 5, 054006]

$J^{PC} = 0^{++}$

TQ4Q1.0+



TQ4Q1.1+



TQ4Q2.0+



$J^{PC} = 1^{+-}$

Landau-Yang  
LO: evolution only



TQ4Q1.1+



C-parity  
LO: evolution only



$J^{PC} = 2^{++}$

TQ4Q1.0+



TQ4Q1.1+






TQ4Q2.0+





# HF-NRevo: Initial-Scale NRCQD Inputs




Scalar  $[J^{PC} = 0^{++}]$

Tensor  $[J^{PC} = 2^{++}]$


HF-NRevo for  $T_{4c}(0^{++}, 2^{++})$  collinear fragmentation

Partonic  
hard factor



DGLAP A0evo

All-Order evolution (numeric)



DGLAP EDevo


Expanded Decoupled evolution



NRQCD

Color-composite

SDCs LDMEs



$$\mathcal{D}_i^Q(z, [n]) \otimes \langle \mathcal{O}^Q([n]) \rangle$$

Energy

Evolution-ready scale

$$Q_0 \equiv m_b + 4m_c$$




$$5m_c$$

$$m_{\bar{q}} + 4m_c$$

Time



# HF-NRevo: Initial-Scale NRCQD Inputs




Scalar  $[J^{PC} = 0^{++}]$

Tensor  $[J^{PC} = 2^{++}]$


## HF-NRevo for $T_{4c}(0^{++}, 2^{++})$ collinear fragmentation

Partonic  
hard factor



DGLAP A0evo

All-Order evolution (numeric)




DGLAP EDevo

Expanded Decoupled evolution



NRQCD

Color-composite  
SDCs LDMEs



$$\mathcal{D}_i^Q(z, [n]) \otimes \langle \mathcal{O}^Q([n]) \rangle$$

Energy

Evolution-ready scale

$$Q_0 \equiv m_b + 4m_c$$




$$5m_c$$

$$m_{\tilde{q}} + 4m_c$$

Time



# HF-NRevo: Initial-Scale NRCQD Inputs




Scalar [ $J^{PC} = 0^{++}$ ]

Tensor [ $J^{PC} = 2^{++}$ ]


## HF-NRevo for $T_{4c}(0^{++}, 2^{++})$ collinear fragmentation

Partonic  
hard factor




DGLAP A0evo

All-Order evolution (numeric)




DGLAP EDevo

Expanded Decoupled evolution



NRQCD

Color-composite  
SDCs LDMEs



$$\mathcal{D}_i^Q(z, [n]) \otimes \langle \mathcal{O}^Q([n]) \rangle$$

Energy

Evolution-ready scale

$$Q_0 \equiv m_b + 4m_c$$

$$5m_c$$


$$m_{\tilde{q}} + 4m_c$$

Time



# TQ4Q2.0: F-MHOU Replica Analysis

$T_{4Q}$







# TQ4Q2.0: F-MHOU Replica Analysis

$T_{4Q}$

Evolution-ready renormalization scale



Evolution-ready factorization scale





TQ4Q2.0\_cs\_n1o


JETHAD

v0.5.5



2




# TQ4Q2.0: Charm Fragmentation to Charmoniumlike



$[c \rightarrow T_{4c}(1^{+-})]$  fragmentation channel







$[c \rightarrow T_{4c}(2^{++})]$  fragmentation channel









# TQ4Q2.0: Gluon Fragmentation to Charmoniumlike






# TQ4Q2.0: Bottom Fragmentation to Charmoniumlike








# TQ4Q2.0: Light-Quark Fragmentation to Charmoniumlike



$[u \rightarrow T_{4c}(1^{+-})]$  fragmentation channel



$[u \rightarrow T_{4c}(2^{++})]$  fragmentation channel





A Multimodal Perspective

R

# Final Remarks

JETHAD

v0.5.5

A Portal to Exotic Matter



# Exotic Matter State-Driven MultiModal Fragmentation

## Quantum Superposition



Compact

$$+ e^{-i\vartheta_1\Delta t}$$




Diquark

$$+ e^{-i\vartheta_2\Delta t}$$



Molecule

$$+ e^{-i\vartheta_3\Delta t}$$





Hadron-onium



# Exotic Matter State-Driven MultiModal Fragmentation


## Quantum Superposition






# Exotic Matter State-Driven MultiModal Fragmentation

## Quantum Superposition





# Heavy-Flavor Fragmentation from HF-NRevo





# Heavy-Flavor Fragmentation from HF-NRevo

 **Collinear\_FF**s Public

main

1 Branch

0 Tags

Go to file

FGCeliberto

Update Readme.md

NRFF10

OMG3Q10

OMG3Q11

PQ5Q10

PQ5Q11

TQ4Q10

TQ4Q11

TQ4Q20

TQHL10

TQHL11

LICENSE

README.md

README

CC0-1.0 license

## Collinear\_FF

Collections of collinear fragmentation functions (FFs) given as LHPDF grids.

Available FF families to date:

- NRFFx.y (NonRelativistic Fragmentation Functions)  
NRFF1.0 (NLO, 2 sets, central value only, LDME uncertainties)
- OMG3Qx.y (OMeGa baryons with 3 heavy Quarks VFNS FFs)  
OMG3Q1.0 (NLO, 2 sets, central value only, diquark-like)  
OMG3Q1.1 (NLO, 2 sets, F-MHOU + F-NPWF uncertainties, replica 0->8, diquark-like)
- TQHLx.y (TetraQuarks with Heavy and Light flavors)  
TQHL1.0 (NLO, 4 sets, central value only)  
TQHL1.1 (NLO, 4 sets, central value only)
- TQ4Qx.y (TetraQuarks with 4 heavy Quarks)  
TQ4Q1.0 (NLO, 4 sets, central value only)  
TQ4Q1.1 (NLO, 6 main + 4 supplementary sets, replica-like MHOU uncertainties in main sets)  
TQ4Q2.0 (NLO, 6 sets, replica-like MHOU uncertainties)
- PQ5Qx.y (PentaQuarks with 5 heavy Quarks)  
PQ5Q1.0 (NLO, 2 sets, replica 0 = direct multiquark, replica 1 = diquark-like)  
PQ5Q1.1 (NLO, 1 set, F-MHOU + F-NPWF uncertainties, replica 0->8 = direct multicharm, replica 9->17 = diquark-like)



# Heavy-Flavor Fragmentation from HF-NRevo

 **Collinear\_FF's** Public

main

1 Branch

0 Tags

Go to file

 **FGCeliberto** Update Readme.md

NRFF10

OMG3Q10

OMG3Q11

PQ5Q10

PQ5Q11

TQ4Q10

TQ4Q11

TQ4Q20

TQHL10

TQHL11

LICENSE

README.md

README

CC0-1.0 license

## Collinear\_FF's

Collections of collinear fragmentation functions (FFs) given as LHPDF grids.

Available FF families to date:

- NRFFx.y (NonRelativistic Fragmentation Functions)  
NRFF1.0 (NLO, 2 sets, central value only, LDME uncertainties)
- OMG3Qx.y (OMeGa baryons with 3 heavy Quarks VFNS FFs)  
OMG3Q1.0 (NLO, 2 sets, central value only, diquark-like)  
OMG3Q1.1 (NLO, 2 sets, F-MHOU + F-NPWF uncertainties, replica 0->8, diquark-like)
- TQHLx.y (TetraQuarks with Heavy and Light flavors)  
TQHL1.0 (NLO, 4 sets, central value only)  
TQHL1.1 (NLO, 4 sets, central value only)
- TQ4Qx.y (TetraQuarks with 4 heavy Quarks)  
TQ4Q1.0 (NLO, 4 sets, central value only)  
TQ4Q1.1 (NLO, 6 main + 4 supplementary sets, replica-like MHOU uncertainties in main sets)  
TQ4Q2.0 (NLO, 6 sets, replica-like MHOU uncertainties)
- PQ5Qx.y (PentaQuarks with 5 heavy Quarks)  
PQ5Q1.0 (NLO, 2 sets, replica 0 = direct multiquark, replica 1 = diquark-like)  
PQ5Q1.1 (NLO, 1 set, F-MHOU + F-NPWF uncertainties, replica 0->8 = direct multicharm, replica 9->17 = diquark-like)

NEW

TQ4Qhf1.0 jto be released soon!


[https://github.com/FGCeliberto/Collinear\\_FF's/](https://github.com/FGCeliberto/Collinear_FF's/)



# Toward ExoFF




# Toward ExoFF




Exotic hadrons  $\Rightarrow$  precision QCD & production mechanisms



# Toward ExoFF




Exotic hadrons  $\Rightarrow$  precision QCD & production mechanisms




**HF-NRevo**  $\Rightarrow$  consistent evolution, uncertainty-aware FFs




# Toward ExoFF



Exotic hadrons  $\Rightarrow$  precision QCD & production mechanisms




**HF-NRevo**  $\Rightarrow$  consistent evolution, uncertainty-aware FFs




AI-driven extraction & global fits of heavy-hadron FFs




# Toward ExoFF




Exotic hadrons  $\Rightarrow$  precision QCD & production mechanisms



**HF-NRevo**  $\Rightarrow$  consistent evolution, uncertainty-aware FFs




AI-driven extraction & global fits of heavy-hadron FFs




From theory-driven benchmarks to state-oriented fragmentation




# Toward ExoFF




Exotic hadrons  $\Rightarrow$  precision QCD & production mechanisms




**HF-NRevo**  $\Rightarrow$  consistent evolution, uncertainty-aware FFs



AI-driven extraction & global fits of heavy-hadron FFs



From theory-driven benchmarks to state-oriented fragmentation



Road to a MULTIMODAL **ExoFF** framework



**EXTRAS**



# TetraQuark-Jet Systems with Resummation: Event Yields

| Expected events yields [13 TeV Run 2] |          |                      |                      |              |
|---------------------------------------|----------|----------------------|----------------------|--------------|
| $T_{4Q}$                              | $J^{PC}$ | TQ4Q2.0              | TQ4Q1.1              | $\Delta$ (%) |
| $T_{4c}$                              | $0^{++}$ | $2016031 \pm 399168$ | $1768764 \pm 309078$ | +13.98%      |
| $T_{4c}$                              | $1^{+-}$ | $8619 \pm 1770$      | $8619 \pm 1612$      | +0%          |
| $T_{4c}$                              | $2^{++}$ | $3071882 \pm 602910$ | $2721362 \pm 480942$ | +12.88%      |
| $T_{4b}$                              | $0^{++}$ | $6046 \pm 1168$      | $5151 \pm 945$       | +17.38%      |
| $T_{4b}$                              | $1^{+-}$ | $17 \pm 3$           | $17 \pm 2$           | +0%          |
| $T_{4b}$                              | $2^{++}$ | $8095 \pm 1552$      | $7108 \pm 828$       | +13.88%      |

TABLE IV: Expected event yields for all-heavy tetraquark production at  $\sqrt{s} = 13$  TeV, obtained by integrating the NLL-resummed  $\Delta Y$  distributions over the rapidity range  $|\Delta Y| < 6.5$ . The yields assume an integrated luminosity of  $\mathcal{L}^{(\text{CMS})} = 138.6 \text{ fb}^{-1}$ , corresponding to the total dataset collected by CMS during Run 2 [308, 309]. The last column shows the relative variation  $\Delta = (R - 1) \times 100$ , with  $R = \text{TQ4Q2.0}/\text{TQ4Q1.1}$ . Uncertainties are propagated from the  $\Delta Y$  distributions and reflect the combined effect of scale variations, FF evolution, and LDME inputs.

| Expected events yields [100 TeV FCC] |          |                        |                        |              |
|--------------------------------------|----------|------------------------|------------------------|--------------|
| $T_{4Q}$                             | $J^{PC}$ | TQ4Q2.0                | TQ4Q1.1                | $\Delta$ (%) |
| $T_{4c}$                             | $0^{++}$ | $22140238 \pm 4433814$ | $19465674 \pm 3373524$ | +13.74%      |
| $T_{4c}$                             | $1^{+-}$ | $23463 \pm 1094$       | $23463 \pm 1053$       | +0%          |
| $T_{4c}$                             | $2^{++}$ | $32784967 \pm 6421338$ | $29144313 \pm 5467493$ | +12.49%      |
| $T_{4b}$                             | $0^{++}$ | $23424 \pm 1982$       | $20237 \pm 1552$       | +15.75%      |
| $T_{4b}$                             | $1^{+-}$ | $50 \pm 2$             | $50 \pm 3$             | +0%          |
| $T_{4b}$                             | $2^{++}$ | $27894 \pm 1972$       | $24493 \pm 1323$       | +13.89%      |

TABLE V: Expected event yields for all-heavy tetraquark production at  $\sqrt{s} = 100$  TeV, obtained by integrating the NLL-resummed  $\Delta Y$  distributions over the rapidity range  $|\Delta Y| < 6.5$ . The yields assume an integrated luminosity of  $\mathcal{L}^{(\text{CMS})} = 138.6 \text{ fb}^{-1}$ , corresponding to the full Run 2 CMS dataset [308, 309]. This value is conservatively adopted at  $\sqrt{s} = 100$  TeV to enable a baseline comparison of energy scaling, independently of specific FCC projections. The last column shows the relative variation  $\Delta = (R - 1) \times 100$ , with  $R = \text{TQ4Q2.0}/\text{TQ4Q1.1}$ . Uncertainties are propagated from the  $\Delta Y$  distributions and reflect the combined effect of scale variations, FF evolution, and LDME inputs.



Pile-up effects can degrade  
Experimental discrimination



# TetraQuark-Jet Systems with Resummation: Event Yields

| Expected events yields [13 TeV Run 2] |          |                      |                      |              |
|---------------------------------------|----------|----------------------|----------------------|--------------|
| $T_{4Q}$                              | $J^{PC}$ | TQ4Q2.0              | TQ4Q1.1              | $\Delta$ (%) |
| $T_{4c}$                              | $0^{++}$ | $2016031 \pm 399168$ | $1768764 \pm 309078$ | +13.98%      |
| $T_{4c}$                              | $1^{+-}$ | $8619 \pm 1770$      | $8619 \pm 1612$      | +0%          |
| $T_{4c}$                              | $2^{++}$ | $3071882 \pm 602910$ | $2721362 \pm 480942$ | +12.88%      |
| $T_{4b}$                              | $0^{++}$ | $6046 \pm 1168$      | $5151 \pm 945$       | +17.38%      |
| $T_{4b}$                              | $1^{+-}$ | $17 \pm 3$           | $17 \pm 2$           | +0%          |
| $T_{4b}$                              | $2^{++}$ | $8095 \pm 1552$      | $7108 \pm 828$       | +13.88%      |

TABLE IV: Expected event yields for all-hadron tetraquark production at  $\sqrt{s} = 13$  TeV, obtained by integrating the NLL-resummed  $\Delta Y$  distributions over the rapidity range  $|\Delta Y| < 6.5$ . The yields assume an integrated luminosity of  $\mathcal{L}^{(\text{CMS})} = 138.6 \text{ fb}^{-1}$ , corresponding to the total dataset collected by CMS during Run 2 [308, 309]. The last column shows the relative variation  $\Delta = (R - 1) \times 100$ , with  $R = \text{TQ4Q2.0}/\text{TQ4Q1.1}$ . Uncertainties are propagated from the  $\Delta Y$  distributions and reflect the combined effect of scale variations, FF evolution, and LDME inputs.

| Expected events yields [100 TeV FCC] |          |                        |                        |              |
|--------------------------------------|----------|------------------------|------------------------|--------------|
| $T_{4Q}$                             | $J^{PC}$ | TQ4Q2.0                | TQ4Q1.1                | $\Delta$ (%) |
| $T_{4c}$                             | $0^{++}$ | $22140238 \pm 4433814$ | $19465674 \pm 3373524$ | +13.74%      |
| $T_{4c}$                             | $1^{+-}$ | $23463 \pm 1094$       | $23463 \pm 1053$       | +0%          |
| $T_{4c}$                             | $2^{++}$ | $32784967 \pm 6421338$ | $29144313 \pm 5467493$ | +12.49%      |
| $T_{4b}$                             | $0^{++}$ | $23424 \pm 1982$       | $20237 \pm 1552$       | +15.75%      |
| $T_{4b}$                             | $1^{+-}$ | $50 \pm 2$             | $50 \pm 3$             | +0%          |
| $T_{4b}$                             | $2^{++}$ | $27894 \pm 1972$       | $24493 \pm 1323$       | +13.89%      |

TABLE V: Expected event yields for all-hadron tetraquark production at  $\sqrt{s} = 100$  TeV, obtained by integrating the NLL-resummed  $\Delta Y$  distributions over the rapidity range  $|\Delta Y| < 6.5$ . The yields assume an integrated luminosity of  $\mathcal{L}^{(\text{CMS})} = 138.6 \text{ fb}^{-1}$ , corresponding to the full Run 2 CMS dataset [308, 309]. This value is conservatively adopted at  $\sqrt{s} = 100$  TeV to enable a baseline comparison of energy scaling, independently of specific FCC projections. The last column shows the relative variation  $\Delta = (R - 1) \times 100$ , with  $R = \text{TQ4Q2.0}/\text{TQ4Q1.1}$ . Uncertainties are propagated from the  $\Delta Y$  distributions and reflect the combined effect of scale variations, FF evolution, and LDME inputs.




Pile-up effects can degrade  
Experimental discrimination




# All-Charm TetraQuarks & Intrinsic Charm

## Quakonium Studies: Challenges



Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity



Low  $x$  and low  $Q^2$


- [G. Altarelli et al. (1998)]
- [A. Candido et al. (2020)]
- [J. Collins et al. (2022)]
- [A. Candido et al. (2023)]



# All-Charm TetraQuarks & Intrinsic Charm

## Quakonium Studies: Challenges

Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity




Low  $x$  and low  $Q^2$

- [G. Altarelli et al. (1998)]
- [A. Candido et al. (2020)]
- [J. Collins et al. (2022)]
- [A. Candido et al. (2023)]

$\gamma + g \rightarrow J/\psi + c \Rightarrow$  Valence Intrinsic Charm @EIC

Large  $x$   
Moderate to large  $p_T$




( $J/\psi$  @EIC) [C. Flore, J.-P. Lansberg, H.-S. Shao, Y. Yedelkina (2020)]  
(Intrinsic Charm + valence) studies [NNPDF Collaboration (2022, 2023)]



# All-Charm TetraQuarks & Intrinsic Charm

## Quakonium Studies: Challenges

Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity




Low  $x$  and low  $Q^2$

- [G. Altarelli et al. (1998)]
- [A. Candido et al. (2020)]
- [J. Collins et al. (2022)]
- [A. Candido et al. (2023)]

$\gamma + g \rightarrow J/\psi + c \Rightarrow$  Valence Intrinsic Charm @EIC

Large  $x$   
Moderate to large  $p_T$




( $J/\psi$  @EIC) [C. Flore, J.-P. Lansberg, H.-S. Shao, Y. Yedelkina (2020)]  
(Intrinsic Charm + valence) studies [NNPDF Collaboration (2022, 2023)]

## TetraQuark Explorations: Assets

$\{T_{4c}\} \ni X(6900) \Rightarrow$  Charm PDF @HL-LHC, FCC

Forward  $0^{++}$ ,  $1^{+-}$ ,  $2^{++}$  within the CGC

[F.G. C, A.V. Giannini, V.P. Gonçalves, Y.N. Lima (2025)]




$$\sigma(pp \rightarrow T_{4c}X) \propto g(x_1, Q^2) \otimes \mathcal{N}_A(x_2) \otimes D_{g/T_{4c}} + c(x_1, Q^2) \otimes \mathcal{N}_F(x_2) \otimes D_{c/T_{4c}}$$



# All-Charm TetraQuarks & Intrinsic Charm

## Quakonium Studies: Challenges

Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity




Low x and low  $Q^2$

- $\circ$  [G. Altarelli et al. (1998)]
- $\circ$  [A. Candido et al. (2020)]
- $\circ$  [J. Collins et al. (2022)]
- $\circ$  [A. Candido et al. (2023)]

$\gamma + g \rightarrow J/\psi + c \Rightarrow$  Valence Intrinsic Charm @EIC

Large x  
Moderate to large  $p_T$




( $J/\psi$  @EIC)  $\circ$  [C. Flore, J.-P. Lansberg, H.-S. Shao, Y. Yedelkina (2020)]  
(Intrinsic Charm + valence) studies  $\circ\circ$  [NNPDF Collaboration (2022, 2023)]

## TetraQuark Explorations: Assets

$\{T_{4c}\} \ni X(6900) \Rightarrow$  Charm PDF @HL-LHC, FCC


Forward  $0^{++}$ ,  $1^{+-}$ ,  $2^{++}$  within the CGC

$\circ$  [F.G. C, A.V. Giannini, V.P. Gonçalves, Y.N. Lima (2025)]




$$\sigma(pp \rightarrow T_{4c}X) \propto g(x_1, Q^2) \otimes \mathcal{N}_A(x_2) \otimes D_{g/T_{4c}} + c(x_1, Q^2) \otimes \mathcal{N}_F(x_2) \otimes D_{c/T_{4c}}$$

Forward  $T_{4c}(1^{+-}) \Rightarrow$  Intrinsic Charm @pp, pA



$1^{+-}$  FFs

dilute-dense






# Highlights on Heavy-Flavor Fragmentation

We observe **hadrons**, whose momentum is different from that of the parent heavy quark. We usually write this as

$$FF(\text{hadron}) = FF(\text{quark}) \otimes FF(\text{quark} \rightarrow \text{hadron})$$

What are the characteristics of  $FF(\text{quark} \rightarrow \text{hadron})$ ?



The heavy quark has momentum  $P = mv_Q$ , the light quarks have momentum  $q = \Lambda v_q$ , with  $\Lambda$  a hadronic mass scale

For the binding we need  $v_Q \simeq v_q = v$ . We have then

$$P = zP + q \qquad mv = zmv + \Lambda v$$

and therefore

$$\langle z \rangle \simeq 1 - \frac{\Lambda}{m}$$

Hence,  $FF(\text{quark} \rightarrow \text{hadron})$  is peaked near  $z = 1$  and binding effects scale linearly in the heavy quark mass (Bjorken/Suzuki ( $\sim 1977$ ))

(NB. more rigorous derivations of this result are available)



# Quarkonia as Tools: Precision & Exploration

Precision determinations  
of QCD running coupling

$\Upsilon(1S)$  hadronic inclusive decays

[CLEO Collaboration (2006)] [🔗](#)

[N. Brambilla et al. (2007)] [🔗](#)

[A. Deur et al. (2016)] [🔗](#)

[ $\alpha_s(2019)$  ECT\* Trento WG (2019)] [🔗](#)

**QUARKONIA**  
QCD Hydrogen Atoms





# Quarkonia as Tools: Precision & Exploration

## QUARKONIA QCD Hydrogen Atoms

Precision determinations  
of QCD running coupling

$\Upsilon(1S)$  hadronic inclusive decays

[CLEO Collaboration (2006)] [link](#)

[N. Brambilla et al. (2007)] [link](#)

[A. Deur et al. (2016)] [link](#)

[ $\alpha_s(2019)$  ECT\* Trento WG (2019)] [link](#)

[G. Altarelli et al. (1998)] [link](#)

[J. Collins et al. (2022)] [link](#)

[A. Candido et al. (2023)] [link](#)

Forward inclusive reactions

Gluon PDF positivity  
at low  $x$  and low  $Q^2$



# Quarkonia as Tools: Precision & Exploration

## QUARKONIA QCD Hydrogen Atoms

Precision determinations  
of QCD running coupling

$\Upsilon(1S)$  hadronic inclusive decays

[CLEO Collaboration (2006)]

[N. Brambilla et al. (2007)]

[A. Deur et al. (2016)]

[ $\alpha_s(2019)$  ECT\* Trento WG (2019)]

[E. G. Ferreira (2015)]

[N. Brambilla et al. (2018)]

[E. G. Ferreira et al. (2019)]

Quarkonia in nuclear matter

Confined matter to  
QGP transition

[G. Altarelli et al. (1998)]

[J. Collins et al. (2022)]

[A. Candido et al. (2023)]

Forward inclusive reactions

Gluon PDF positivity  
at low  $x$  and low  $Q^2$



# Quarkonia as Tools: Precision & Exploration

Portals to exotic matter  
and BSM physics

Hadroquarkonium  
EW boson rare decays

- [\[S. Dubynskiy et al. \(2008\)\]](#)
- [\[F.-K. Guo et al. \(2018\)\]](#)
- [\[J. Ferretti et al. \(2020\)\]](#)
- [\[CMS Collaboration \(2019\)\]](#)

- [\[E. G. Ferreira \(2015\)\]](#)
- [\[N. Brambilla et al. \(2018\)\]](#)
- [\[E. G. Ferreira et al. \(2019\)\]](#)

Quarkonia in nuclear matter

Confined matter to  
QGP transition

Precision determinations  
of QCD running coupling

$\Upsilon(1S)$  hadronic inclusive decays


- [\[CLEO Collaboration \(2006\)\]](#)
- [\[N. Brambilla et al. \(2007\)\]](#)
- [\[A. Deur et al. \(2016\)\]](#)
- [\[ \$\alpha\_s\(2019\)\$  ECT\\* Trento WG \(2019\)\]](#)

- [\[G. Altarelli et al. \(1998\)\]](#)
- [\[J. Collins et al. \(2022\)\]](#)
- [\[A. Candido et al. \(2023\)\]](#)

Forward inclusive reactions

Gluon PDF positivity  
at low  $x$  and low  $Q^2$


**QUARKONIA**  
QCD Hydrogen Atoms






# Quarkonium Studies: Assets & Challenges

## Hadronic structure: Assets



EIC, LHCb, LHCspin

Boer-Mulders




 [A. Bacchetta, F.G. C., M. Radici, P. Tael (EPJC 2020)]




# Quarkonium Studies: Assets & Challenges

## Hadronic structure: Assets





EIC, LHCb, LHCspin

Boer-Mulders



[A. Bacchetta, F.G. C., M. Radici, P. Tael (EPJC 2020)]

[A. Bacchetta, F.G. C., M. Radici (EPJC 2024)]




EIC, LHCspin

Sivers




# Quarkonium Studies: Assets & Challenges

## Hadronic structure: Assets





EIC, LHCb, LHCspin

Boer-Mulders



[A. Bacchetta, F.G. C., M. Radici, P. Tael (EPJC 2020)]


[A. Bacchetta, F.G. C., M. Radici (EPJC 2024)]




EIC, LHCspin

Sivers

## Precision QCD: Challenges



Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity




Low x and low  $Q^2$

- [G. Altarelli et al. (1998)]
- [A. Candido et al. (2020)]
- [J. Collins et al. (2022)]
- [A. Candido et al. (2023)]




# Quarkonium Studies: Assets & Challenges

## Hadronic structure: Assets





EIC, LHCb, LHCspin

Boer-Mulders



[A. Bacchetta, F.G. C., M. Radici, P. Tael (EPJC 2020)]


[A. Bacchetta, F.G. C., M. Radici (EPJC 2024)]




EIC, LHCspin

Sivers

## Precision QCD: Challenges




Forward Onia  $\Rightarrow$  Gluon PDF & TMD Positivity




Low x and low  $Q^2$

- [G. Altarelli et al. (1998)]
- [A. Candido et al. (2020)]
- [J. Collins et al. (2022)]
- [A. Candido et al. (2023)]



$\gamma + g \rightarrow J/\psi + c \Rightarrow$  Valence Intrinsic Charm @EIC


Large x  
Moderate to large  $p_T$




( $J/\psi$  @EIC) [C. Flore, J.-P. Lansberg, H.-S. Shao, Y. Yedelkina (2020)]  
(Intrinsic Charm + valence) studies [NNPDF Collaboration (2022, 2023)]



# Highlights of Non-Relativistic QCD

 **NRQCD** → effective field theory → nonrelativistic (NR)  $Q\bar{Q}$  system


 Heavy-quark spinor fields as NR DOFs in the NRQCD Lagrangian


$$\mathcal{L}_{\text{NRQCD}} = \psi^\dagger \left( iD_0 + \frac{\vec{D}^2}{2m_Q} \right) \psi + \chi^\dagger \left( iD_0 - \frac{\vec{D}^2}{2m_Q} \right) \chi + \mathcal{L}_{\text{light}} + \delta\mathcal{L}$$





# Highlights of Non-Relativistic QCD

 **NRQCD**  $\rightarrow$  effective field theory  $\rightarrow$  nonrelativistic (NR)  $Q\bar{Q}$  system

 Heavy-quark spinor fields as NR DOFs in the NRQCD Lagrangian

$$\mathcal{L}_{\text{NRQCD}} = \psi^\dagger \left( iD_0 + \frac{\vec{D}^2}{2m_Q} \right) \psi + \chi^\dagger \left( iD_0 - \frac{\vec{D}^2}{2m_Q} \right) \chi + \mathcal{L}_{\text{light}} + \delta\mathcal{L}$$


 [cross section]  $=$  [short-distance coeffs.]  $\otimes$  [long-distance matrix elements]


$$d\sigma(H + X) = \sum_n d\hat{\sigma}(Q\bar{Q}[n] + X) \langle \mathcal{O}^H[n] \rangle$$






# Highlights of Non-Relativistic QCD

 **NRQCD** → effective field theory → nonrelativistic (NR)  $Q\bar{Q}$  system

 Heavy-quark spinor fields as NR DOFs in the NRQCD Lagrangian

$$\mathcal{L}_{\text{NRQCD}} = \psi^\dagger \left( iD_0 + \frac{\vec{D}^2}{2m_Q} \right) \psi + \chi^\dagger \left( iD_0 - \frac{\vec{D}^2}{2m_Q} \right) \chi + \mathcal{L}_{\text{light}} + \delta\mathcal{L}$$

 [cross section] = [short-distance coeffs.]  $\otimes$  [long-distance matrix elements]

$$d\sigma(H + X) = \sum_n d\hat{\sigma}(Q\bar{Q}[n] + X) \langle \mathcal{O}^H[n] \rangle$$

$$\begin{aligned} |\mathcal{Q}\rangle &= \mathcal{O}(1) |Q\bar{Q}[^3S_1^{(1)}]\rangle + \mathcal{O}(v) |Q\bar{Q}[^3P_J^{(8)}g]\rangle + \mathcal{O}(v^2) |Q\bar{Q}[^1S_0^{(8)}g]\rangle \\ &\quad + \mathcal{O}(v^2) |Q\bar{Q}[^3S_1^{(1,8)}gg]\rangle + \mathcal{O}(v^2) |Q\bar{Q}[^3D_J^{(1,8)}gg]\rangle + \dots \end{aligned}$$



**HF-NREVO**

**FOUNDING PILLARS**

# Heavy-Flavor Non-Relativistic evolution

**HF-NRevo**

Interpretation





# Heavy-Flavor Non-Relativistic evolution

**HF-NRevo**

Interpretation

Evolution



# Heavy-Flavor Non-Relativistic evolution

**HF-NRevo**

Interpretation

Evolution

Uncertainties





# Heavy-Flavor Non-Relativistic evolution

**HF-NRevo**

Interpretation

Evolution

Uncertainties

- NRQCD  $\Rightarrow$  proxy for initial-scale FF inputs
- A heavy-flavor-scheme based vision
- Low vs large  $p_T \Rightarrow$  FFNS vs VFNS fragmentation

# Heavy-Flavor Non-Relativistic evolution

**HF-NRevo**

Interpretation

Evolution

Uncertainties

- Proper DGLAP collinear evolution
- Thresholds at work for all partons
- Two-step strategy  $\Rightarrow$  EDevo  $\otimes$  ADevo



# Heavy-Flavor Non-Relativistic evolution

## HF-NRevo

Interpretation

Evolution

Uncertainties

- Initial-scale MHOUs  $\Rightarrow$  first systematic analysis
- Idea  $\Leftrightarrow$  **ThCovM**   & **MCscales**  MHOUs
- Future  $\Leftarrow$  extraction of NP LDMEs from data

**ALL-HEAVY PENTAQUARK**

**FRAGMENTATION**



# MultiModal Fragmentation to PentaQuarks

Mode 1

Direct MultiQuark  
Scalar S-wave  $P_{5c}$




# MultiModal Fragmentation to PentaQuarks

## Mode 1 Direct MultiQuark Scalar S-wave $P_{5c}$

Initial-scale input:  Suzuki model

**PQ5Q1.y** FFs:  $(Q \rightarrow P_{c\bar{c}ccc}) \otimes$  **APFEL++**

F-MHOUs [P] + F-NPWF [NP]








# MultiModal Fragmentation to PentaQuarks

## Mode 2 Scalar DiQuark Scalar S-wave $P_{5c}$

Initial-scale input:  Suzuki model

**PQ5Q1.y** FFs:  $(Q \rightarrow P_{c\bar{c}ccc}) \otimes$  **APFEL++**

F-MHOUs [P] + F-NPWF [NP]






# PQ5Q1.1\_cs\_n1o


JETHAD

v0.5.5








# PQ5Q1.1: Charm Fragmentation to PentaCharm




$[c \rightarrow P_{5c}]$  direct fragmentation





$[c \rightarrow P_{5c}]$  diquark fragmentation




# PQ5Q1.1: Gluon Fragmentation to PentaCharm



$[g \rightarrow P_{5c}]$  direct fragmentation




$[g \rightarrow P_{5c}]$  diquark fragmentation








# PQ5Q1.1: Energy Spectrum



$P_{5c}$  direct fragmentation




$P_{5c}$  diquark fragmentation




Direct



Dicharm




# PQ5Q1.1: Energy Spectrum




$P_{5c}$  direct fragmentation




$P_{5c}$  diquark fragmentation



$P_{5b}$  direct fragmentation



$P_{5b}$  diquark fragmentation



tCΦR-iD

DiCharm




# QUARKONIUM FRAGMENTATION


# From low energies to high energies

Low  $p_T \Rightarrow$  two-parton FFs  
Genuine higher-twist FFNS FFs

scalar




vector




High  $p_T \Rightarrow$  single-parton FFs  
Twist-2 VFNS FFs + power corrections

scalar





vector







# From low energies to high energies

Low  $p_T \Rightarrow$  two-parton FFs  
Genuine higher-twist FFNS FFs



High  $p_T \Rightarrow$  single-parton FFs  
Twist-2 VFNS FFs + power corrections



Collinear Fragmentation Correlator (CFC)

# From low energies to high energies



Low  $p_T \Rightarrow$  two-parton FFs  
Genuine higher-twist FFNS FFs

High  $p_T \Rightarrow$  single-parton FFs  
Twist-2 VFNS FFs + power corrections

## Matching low & high $p_T$

• TMD matching tails  $\Rightarrow$  different singularities

[\[D. Boer, J. Bor, L. Maxia, C. Pisano, F. Yuan \(2023\)\]](#)






# From low energies to high energies

Low  $p_T \Rightarrow$  two-parton FFs  
Genuine higher-twist FFNS FFs

High  $p_T \Rightarrow$  single-parton FFs  
Twist-2 VFNS FFs + power corrections

## Matching low & high $p_T$


scalar



- TMD matching tails  $\Rightarrow$  different singularities

[\[D. Boer, J. Bor, L. Maxia, C. Pisano, F. Yuan \(2023\)\]](#)

vector





- Short-distance  $\Leftrightarrow$  double-parton FFs (SCET)

[\[S. Fleming, A.K. Leibovich, T. Mehen, I.Z. Rothstein \(2012\)\]](#)

# From low energies to high energies

Low energy, low  $p_T$   
Short-distance production




Extra gluon due to Landau-Yang selection rule




# From low energies to high energies



Low energy, low  $p_T$   
Short-distance production



High energy, high  $p_T$   
Single-parton fragmentation



$p_T \gtrsim 10 \div 15 \text{ GeV}$   
(charmonia)



Extra gluon due to Landau-Yang selection rule

# HF-NRevo: Initial-scale NRCQD inputs




**Fig. 1** Left panel: a leading diagram for the fragmentation of a gluon to  $\mathcal{V}_Q [^3S_1^{(1)}]$  at  $\mathcal{O}(\alpha_s^3)$ . Central panel: one of the leading diagrams for the fragmentation of a constituent heavy quark ( $Q$ ) to a  $S$ -wave color singlet vector quarkonium  $\mathcal{V}_Q [^3S_1^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Right panel: a leading diagram for the fragmentation of a nonconstituent light ( $q$ ) or heavy ( $\bar{Q}$ ) quark to  $\mathcal{V}_Q [^3S_1^{(1)}]$  at  $\mathcal{O}(\alpha_s^4)$ . Violet blobs refer to the  $\mathcal{V}_Q [^3S_1^{(1)}]$  nonperturbative NRQCD LDME.




**Fig. 2** Left panel: a leading diagram for the fragmentation of a gluon to  $\mathcal{S}_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Central panel: one of the leading diagrams for the fragmentation of a constituent heavy quark ( $Q$ ) to a  $S$ -wave color singlet pseudoscalar quarkonium  $\mathcal{S}_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Right panel: a leading diagram for the fragmentation of a nonconstituent light ( $q$ ) or heavy ( $\bar{Q}$ ) quark to  $\mathcal{S}_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^3)$ . Green blobs refer to the  $\mathcal{S}_Q^- [^1S_0^{(1)}]$  nonperturbative NRQCD LDME.



# HF-NRevo: Initial-scale NRCQD inputs




**Fig. 1** Left panel: a leading diagrams for  $V_Q [^3S_1^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Middle panel: one of the leading diagrams for  $V_Q [^3S_1^{(1)}]$  at  $\mathcal{O}(\alpha_s^4)$ .

| Quarkonium ( $\mathcal{Q}$ )                                   | $g \rightarrow \mathcal{Q}$                                                                              | $Q \rightarrow \mathcal{Q}$                                                                                                                           | $q$ or $\bar{Q} \rightarrow \mathcal{Q}$                  |
|----------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------|
| $\mathcal{Q} \equiv V_Q [^1S_0^{(1)}] \equiv J/\psi, \Upsilon$ | $-\mathcal{O}(\alpha_s^3) \equiv \text{NLO}$<br>$\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ | $\mathcal{O}(\alpha_s^2) \equiv \text{LO}$<br>$\mathcal{O}(\alpha_s^3) \equiv \text{NLO}$<br>$\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ | $-\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ |





**Fig. 2** Left panel: a leading diagrams for  $S_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Middle panel: one of the leading diagrams for  $S_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^4)$ .

| Quarkonium ( $\mathcal{Q}$ )                                   | $g \rightarrow \mathcal{Q}$                                                                                                                           | $Q \rightarrow \mathcal{Q}$                                                                                                                           | $q$ or $\bar{Q} \rightarrow \mathcal{Q}$                                                                 |
|----------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|
| $\mathcal{Q} \equiv S_Q^- [^3S_1^{(1)}] \equiv \eta_c, \eta_b$ | $\mathcal{O}(\alpha_s^2) \equiv \text{LO}$<br>$\mathcal{O}(\alpha_s^3) \equiv \text{NLO}$<br>$\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ | $\mathcal{O}(\alpha_s^2) \equiv \text{LO}$<br>$\mathcal{O}(\alpha_s^3) \equiv \text{NLO}$<br>$\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ | $-\mathcal{O}(\alpha_s^3) \equiv \text{NLO}$<br>$\mathcal{O}(\alpha_s^4) \equiv \text{NNLO}$<br>$\vdots$ |

# HF-NRevo: Initial-scale NRCQD inputs



| $S_Q^- [^1S_0^{(1)}]$ quarkonium | $\mu_{F,0}(g \rightarrow S_Q^-)$ | $\mu_{F,0}(d, u, s \rightarrow S_Q^-)$ | $\mu_{F,0}(Q \rightarrow S_Q^-)$ | $\mu_{F,0}(\tilde{Q} \rightarrow S_Q^-)$ | $Q_0 \equiv \max(\{\mu_{F,0}\})$ |
|----------------------------------|----------------------------------|----------------------------------------|----------------------------------|------------------------------------------|----------------------------------|
| $\eta_c$                         | $2m_c$                           | $m_{d,u,s} + 2m_c$                     | $3m_c$                           | $m_b + 2m_c$                             | $m_b + 2m_c$                     |
| $\eta_b$                         | $2m_b$                           | $m_{d,u,s} + 2m_b$                     | $3m_b$                           | $m_c + 2m_b$                             | $3m_b$                           |





Fig. 2 Left panel: a leading diagram for the fragmentation of a gluon to  $S_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Central panel: one of the leading diagrams for the fragmentation of a constituent heavy quark ( $Q$ ) to a  $S$ -wave color singlet pseudoscalar quarkonium  $S_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Right panel: a leading diagram for the fragmentation of a nonconstituent light ( $q$ ) or heavy ( $\tilde{Q}$ ) quark to  $S_Q^- [^1S_0^{(1)}]$  at  $\mathcal{O}(\alpha_s^2)$ . Green blobs refer to the  $S_Q^- [^1S_0^{(1)}]$  nonperturbative NRQCD LDME.



# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!



$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q,\text{LO}}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2 \Gamma^{Q,\text{NLO}}(z)$$

$$(Q \rightarrow Q Q) \text{ at } \mu_0 = 3m_Q$$

(LO) [\[E. Braaten et al., Phys. Rev. D 48 \(1993\) 4230-4235\]](#)


(NLO) [\[X. Zheng et al., Phys. Rev. D 100 \(2019\) 1, 014005\]](#)

[\[My talk at the Higgs Centre, Edinburgh \(2023\)\]](#)

[\[F. G. C., M. Fucilla, Eur. Phys. J. C 82 \(2022\) 10, 929\]](#)

# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!




$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q, \text{LO}}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2 \Gamma^{Q, \text{NLO}}(z)$$

$(Q \rightarrow Q Q)$  at  $\mu_0 = 3m_Q$

(LO) [E. Braaten et al., Phys. Rev. D 48 (1993) 4230-4235]

(NLO) [X. Zheng et al., Phys. Rev. D 100 (2019) 1, 014005]

+



$$D_g^Q(z, 2m_Q) = \frac{5}{36(2\pi)^2} \alpha_s^3(2m_Q) \frac{|\mathcal{R}_Q(0)|^2}{m_Q^3} \int_0^z d\xi \int_{(\xi+z^2)/2z}^{(1+\xi)/2} d\tau \frac{1}{(1-\tau)^2(\tau-\xi)^2(\tau^2-\xi)^2} \\ \sum_{i=1}^2 z^i \left[ f_i^{(g)}(\xi, \tau) + g_i^{(g)}(\xi, \tau) \frac{1+\xi-2\tau}{2(\tau-\xi)\sqrt{\tau^2-\xi}} \ln \left( \frac{\tau-\xi+\sqrt{\tau^2-\xi}}{\tau-\xi-\sqrt{\tau^2-\xi}} \right) \right],$$

$(g \rightarrow Q gg)$  at  $\mu_0 = 2m_Q$

(LO) [A. Braaten, T.C Yuan, Phys. Rev. Lett. 71 (1993), 1673]


[My talk at the Higgs Centre, Edinburgh (2023)]

[F. G. C., M. Fucilla, Eur. Phys. J. C 82 (2022) 10, 929]



# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!




$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q, \text{LO}}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2 \Gamma^{Q, \text{NLO}}(z)$$

$$(Q \rightarrow Q Q) \text{ at } \mu_0 = 3m_Q$$

(LO) [E. Braaten et al., Phys. Rev. D 48 (1993) 4230-4235]

(NLO) [X. Zheng et al., Phys. Rev. D 100 (2019) 1, 014005]

+



$$D_g^Q(z, 2m_Q) = \frac{5}{36(2\pi)^2} \alpha_s^3(2m_Q) \frac{|\mathcal{R}_Q(0)|^2}{m_Q^3} \int_0^z d\xi \int_{(\xi+z^2)/2z}^{(1+\xi)/2} d\tau \frac{1}{(1-\tau)^2(\tau-\xi)^2(\tau^2-\xi)^2} \\ \sum_{i=1}^2 z^i \left[ f_i^{(g)}(\xi, \tau) + g_i^{(g)}(\xi, \tau) \frac{1+\xi-2\tau}{2(\tau-\xi)\sqrt{\tau^2-\xi}} \ln \left( \frac{\tau-\xi+\sqrt{\tau^2-\xi}}{\tau-\xi-\sqrt{\tau^2-\xi}} \right) \right],$$

$$(g \rightarrow Q gg) \text{ at } \mu_0 = 2m_Q$$

⊗

(LO) [A. Braaten, T.C Yuan, Phys. Rev. Lett. 71 (1993), 1673]

[My talk at the Higgs Centre, Edinburgh (2023)]

[F. G. C., M. Fucilla, Eur. Phys. J. C 82 (2022) 10, 929]

**NRFF1.0**

quarkonium FFs


=

**APFEL++/EKO**



# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!




$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q, \text{LO}}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2$$

(LO) [E. Braaten et al., Phys. Rev. D 48 (1993) 4230-4235]  
(NLO) [X. Zheng et al., Phys. Rev. D 100 (2019) 1, 014005]




$(Q \rightarrow Q Q)$  at  $\mu_0 = 3m_Q$

+



$$D_g^Q(z, 2m_Q) = \frac{5}{36(2\pi)^2} \alpha_s^3(2m_Q) \frac{|\mathcal{R}_Q(0)|^2}{m_Q^3} \int_0^z d\xi \int_{(\xi+z^2)/2z}^{(1+\xi)/2} d\tau \frac{1}{(1-\tau)^2} \sum_{i=1}^2 z^i \left[ f_i^{(g)}(\xi, \tau) + g_i^{(g)}(\xi, \tau) \frac{1+\xi-2\tau}{2(\tau-\xi)\sqrt{\tau^2-\xi}} \ln \left( \frac{\tau-\xi+\sqrt{\tau^2-\xi}}{\tau-\xi-\sqrt{\tau^2-\xi}} \right) \right]$$

(LO) [A. Braaten, T.C Yuan, Phys. Rev. Lett. 71 (1993), 1673]



$(g \rightarrow Q gg)$  at  $\mu_0 = 2m_Q$

⊗


[My talk at the Higgs Centre, Edinburgh (2023)]  
[F. G. C., M. Fucilla, Eur. Phys. J. C 82 (2022) 10, 929]

**NRFF1.0**  
quarkonium FFs = **APFEL++/EKO**



# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!




$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q, \text{LO}}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2$$



$(Q \rightarrow Q Q)$  at  $\mu_0 = 3m_Q$

(LO) [E. Braaten et al., Phys. Rev. D 48 (1993) 4230-4235]  
(NLO) [X. Zheng et al., Phys. Rev. D 100 (2019) 1, 014005]

+



$$D_g^Q(z, 2m_Q) = \frac{5}{36(2\pi)^2} \alpha_s^3(2m_Q) \frac{|\mathcal{R}_Q(0)|^2}{m_Q^3} \int_0^z d\xi \int_{(\xi+z^2)/2z}^{(1+\xi)/2} d\tau \frac{1}{(1-\tau)^2} \sum_{i=1}^2 z^i \left[ f_i^{(g)}(\xi, \tau) + g_i^{(g)}(\xi, \tau) \frac{1+\xi-2\tau}{2(\tau-\xi)\sqrt{\tau^2-\xi}} \ln \left( \frac{\tau-\xi+\sqrt{\tau^2-\xi}}{\tau-\xi-\sqrt{\tau^2-\xi}} \right) \right]$$



$(g \rightarrow Q gg)$  at  $\mu_0 = 2m_Q$

(LO) [A. Braaten, T.C Yuan, Phys. Rev. Lett. 71 (1993), 1673]




[My talk at the Higgs Centre, Edinburgh (2023)]  
 [F. G. C., M. Fucilla, Eur. Phys. J. C 82 (2022) 10, 929]

**NRFF1.0**  
quarkonium FFs


=

**APFEL++/EKO**



# Vector quarkonium from single-parton fragmentation

! Let us consider  $J/\psi$  and  $\Upsilon$  at large  $p_T \rightarrow$  initial-scale **heavy-quark** + **gluon** from **NRQCD**!




$$D_Q^Q(z, \mu_F \equiv \mu_0) = D_Q^{Q,LO}(z) + \frac{\alpha_s^3(\mu_R)}{m_Q^3} |\mathcal{R}_Q(0)|^2$$

(LO) [E. Braaten et al., Phys. Rev. D 48 (1993) 4230-4235]  
(NLO) [X. Zheng et al., Phys. Rev. D 100 (2019) 1, 014005]




$(Q \rightarrow Q Q)$  at  $\mu_0 = 3m_Q$

+



$$D_g^Q(z, 2m_Q) = \frac{5}{36(2\pi)^2} \alpha_s^3(2m_Q) \frac{|\mathcal{R}_Q(0)|^2}{m_Q^3} \int_0^z d\xi \int_{(\xi+z^2)/2z}^{(1+\xi)/2} d\tau \frac{1}{(1-\tau)^2} \sum_{i=1}^2 z^i \left[ f_i^{(g)}(\xi, \tau) + g_i^{(g)}(\xi, \tau) \frac{1+\xi-2\tau}{2(\tau-\xi)\sqrt{\tau^2-\xi}} \ln \left( \frac{\tau-\xi+\sqrt{\tau^2-\xi}}{\tau-\xi-\sqrt{\tau^2-\xi}} \right) \right]$$

(LO) [A. Braaten, T.C Yuan, Phys. Rev. Lett. 71 (1993), 1673]



$(g \rightarrow Q gg)$  at  $\mu_0 = 2m_Q$



[My talk at the Higgs Centre, Edinburgh (2023)]  
[F. G. C., M. Fucilla, Eur. Phys. J. C 82 (2022) 10, 929]



NRFF1.0


quarkonium FFs

=

APFEL++/EKO




# HF-NRevo: Initial-scale NRCQD inputs




Scalar  $\eta_c$

## HF-NRevo for $\eta_c[^1S_0^{(1)}]$ collinear fragmentation




# HF-NRevo: Initial-scale NRCQD inputs




Scalar  $\eta_c$

## HF-NRevo for $\eta_c[^1S_0^{(1)}]$ collinear fragmentation






# HF-NRevo: Initial-scale NRCQD inputs



Scalar  $\eta_c$


## HF-NRevo for $\eta_c[{}^1S_0^{(1)}]$ collinear fragmentation

Partonic  
hard factor




DGLAP A0evo

All-Order evolution (numeric)



DGLAP EDevo


Expanded Decoupled evolution



NRQCD

LDMEs

SDCs



$$\mathcal{D}_i^{\mathcal{Q}}(z, [n]) \otimes \langle \mathcal{O}^{\mathcal{Q}}([n]) \rangle$$

Energy

Evolution-ready scale


$$Q_0 \equiv m_b + 2m_c$$

$$3m_c$$

$$m_{\tilde{q}} + 2m_c$$

Time


# HF-NRevo: Initial-scale NRCQD inputs



Scalar  $\eta_b$


## HF-NRevo for $\eta_b[^1S_0^{(1)}]$ collinear fragmentation

Partonic  
hard factor




DGLAP A0evo

All-Order evolution (numeric)




DGLAP EDevo

Expanded Decoupled evolution



NRQCD

LDMEs



$$\mathcal{D}_i^{\mathcal{Q}}(z, [n]) \otimes \langle \mathcal{O}^{\mathcal{Q}}([n]) \rangle$$

Energy

Evolution-ready scale

$$Q_0 \equiv 3m_b$$


$$m_c + 2m_b$$

$$m_{\tilde{q}} + 2m_b$$

Time




# HF-NRevo: Initial-scale NRCQD inputs



Vector  $J/\psi$


## HF-NRevo for $J/\psi[{}^3S_1^{(1)}]$ collinear fragmentation

Partonic  
hard factor




DGLAP A0evo

All-Order evolution (numeric)



DGLAP EDevo

Expanded Decoupled evolution




$2m_c$

NRQCD

LDMEs

SDCs



$$\mathcal{D}_i^{\mathcal{Q}}(z, [n]) \otimes \langle \mathcal{O}^{\mathcal{Q}}([n]) \rangle$$


Energy

Evolution-ready scale

$$Q_0 \equiv 3m_c$$

Time


# HF-NRevo: Initial-scale NRCQD inputs



Vector  $\Upsilon$


## HF-NRevo for $\Upsilon[^3S_1^{(1)}]$ collinear fragmentation

Partonic  
hard factor




DGLAP A0evo

All-Order evolution (numeric)



DGLAP EDevo

Expanded Decoupled evolution




$2m_c$

NRQCD

LDMEs

SDCs



$$\mathcal{D}_i^{\mathcal{Q}}(z, [n]) \otimes \langle \mathcal{O}^{\mathcal{Q}}([n]) \rangle$$

Energy

Evolution-ready scale


$$Q_0 \equiv 3m_b$$

Time




# NRFF1.0: Gluon fragmentation to charmonia


Vector  $J/\psi$




$(g \rightarrow J/\psi)$  fragmentation channel



Scalar  $\eta_c$




$(g \rightarrow \eta_c)$  fragmentation channel




# NRFF1.0: Gluon fragmentation to bottomonia


Vector  $\Upsilon$




$(g \rightarrow \Upsilon)$  fragmentation channel



Scalar  $\eta_b$




$(g \rightarrow \eta_b)$  fragmentation channel






# NRFF1.0: Charm fragmentation to charmonia


Vector  $J/\psi$




$(c \rightarrow J/\psi)$  fragmentation channel



Scalar  $\eta_c$




$(c \rightarrow \eta_c)$  fragmentation channel




# NRFF1.0: Bottom fragmentation to bottomonia


Vector  $\Upsilon$



$(b \rightarrow \Upsilon)$  fragmentation channel



Scalar  $\eta_b$



$(b \rightarrow \eta_b)$  fragmentation channel

