

Space Charge Simulation for the Third Order Resonance Corrections at J-PARC MR

Space Charge 2026, 7th ICFA Mini-workshop

2026/9/30



Susumu Igarashi KEK/J-PARC Center

Contents

- Introduction
- Third order resonance correction for on-momentum particles
- Third order resonance correction for off-momentum particles
- Beam optics modulated by space charge effect
- Simulation results
- Summary


MR Design and Operation Modes

- Circumference 1567.5 m
- Three-fold symmetry
- Target Beam Power: 1.3 MW (for fast extraction)
- Fast extraction mode (FX) for the neutrino Facility: 1 turn extraction.
 - Cycle : 1.28 s $\rightarrow$ 1.16 s
 - Beam Intensity 2.7×10^{14} ppp $\rightarrow$ 3.3×10^{14} ppp
- Slow extraction mode (SX) for the hadron hall: 2 s – spill extraction.
 - Cycle : 4.24 s



Space Charge Tune Spread and Resonances

- MR Power 1.3 MW
- Number of protons:
4.15e13 protons per bunch
- Typical operation tune :
(21.35, 21.45)
- Beam loss reduction is necessary for the high intensity operation.
- Resonances of $3\nu_x = 64$ and $\nu_x + 2\nu_y = 64$ are corrected for the FX operation.



3rd Order Resonances Corrections with Trim Coils of Sextupole Magnets

- $v_x + 2v_y = 64$, (1+2)
- Beam survival recovers with SFA048: +0.5 A, SFA055: +0.2 A.
- $|G| = 0.076$

$$G_{1,2,64} = \frac{\sqrt{2}}{8\rho} b_x^{1/2} b_y k_2 \exp[i(f_x + 2f_y)]$$

- $3v_x = 64$, (3+0)
- Beam survival recovers with SFA048 +0.5 A, SFA055 0.0 A.
- $|G| = 0.055$

$$G_{3,0,64} = \frac{\sqrt{2}}{24\rho} b_x^{3/2} k_2 \exp[i(3f_x)]$$








- Simultaneous equations for 4 sextupole trim coils to correct both resonances.

$$\sum_{j=1}^4 \frac{\sqrt{2}}{24\pi} \beta_x^3(j) k_2(j) \cos[3\phi_x(j)] = \sum_{i=1}^2 \frac{\sqrt{2}}{24\pi} \beta_x^3(i) k_2(i) \cos[3\phi_x(i)]$$

$$\sum_{j=1}^4 \frac{\sqrt{2}}{24\pi} \beta_x^3(j) k_2(j) \sin[3\phi_x(j)] = \sum_{i=1}^2 \frac{\sqrt{2}}{24\pi} \beta_x^3(i) k_2(i) \sin[3\phi_x(i)]$$

$$\sum_{j=1}^4 \frac{\sqrt{2}}{8\pi} \beta_x^{\frac{1}{2}}(j) \beta_y(j) k_2(j) \cos[\phi_x(j) + 2\phi_y(j)] = \sum_{i=1}^2 \frac{\sqrt{2}}{8\pi} \beta_x^{\frac{1}{2}}(i) \beta_y(i) k_2(i) \cos[\phi_x(i) + 2\phi_y(i)]$$




$$\sum_{j=1}^4 \frac{\sqrt{2}}{8\pi} \beta_x^{\frac{1}{2}}(j) \beta_y(j) k_2(j) \sin[\phi_x(j) + 2\phi_y(j)] = \sum_{i=1}^2 \frac{\sqrt{2}}{8\pi} \beta_x^{\frac{1}{2}}(i) \beta_y(i) k_2(i) \sin[\phi_x(i) + 2\phi_y(i)]$$



Beam optics for off-momentum particles

- Beam optics for the off-momentum particles ($dp/p = \pm 0.2\%$, $\pm 0.4\%$,) were calculated with SAD with respect to the optics of on-momentum ($dp/p = 0$) particle .

$$G_{3,0,64} = \frac{\sqrt{2}}{24\rho} b_x^{3/2} k_2 \exp[i(3f_x)]$$



SFA magnets ●

group 1

group 2

QDR081
QFR082

group 3

group 4

group 5

group 6

s (m)


Resonance correction for 3+0 for off-momentum particles

$$G_{3,0,64} = \frac{\sqrt{2}}{24\rho} b_x^{3/2} k_2 \exp[i(3f_x)]$$



- Error at QDR081 and QFR082
- Resonance correction with trim coils of 4 SFA magnets
- Analysis is repeated for the 1+2 resonance.

SFA group 1	Trim Coil Current (A)
020	-1.6328
027	+2.8697
034	-4.7788
041	+3.2752

$G_{3,0}$ for $dp = 0$



$G_{3,0}$ for $dp = +0.2\%$



The real part and imaginary part of $\frac{d\Delta G_{3+0}}{d(\Delta p/p)}$ were derived for group 1.





Resonance correction for off-momentum particles

- Find the solution for the following simultaneous equations.

$$\sum_{i=1}^5 f_i \times \operatorname{Re}\left\{\frac{d\Delta G_{3+0}}{d(\Delta p/p)}(\text{group } i)\right\} = 0 \quad \sum_{i=1}^5 f_i \times \operatorname{Im}\left\{\frac{d\Delta G_{3+0}}{d(\Delta p/p)}(\text{group } i)\right\} = 0$$

$$\sum_{i=1}^5 f_i \times \operatorname{Re}\left\{\frac{d\Delta G_{1+2}}{d(\Delta p/p)}(\text{group } i)\right\} = 0 \quad \sum_{i=1}^5 f_i \times \operatorname{Im}\left\{\frac{d\Delta G_{1+2}}{d(\Delta p/p)}(\text{group } i)\right\} = 0$$


$$\sum_{i=1}^5 f_i = 1 \quad \left(\begin{array}{l} \sum_{i=1}^5 f_i \times \Delta G_{3+0}(\text{group } i)_{\text{cor}} = -\Delta G_{3+0 \text{ source}} \\ \sum_{i=1}^5 f_i \times \Delta G_{1+2}(\text{group } i)_{\text{cor}} = -\Delta G_{1+2 \text{ source}} \end{array} \right)$$




SFA	$\Delta G_{3+0} / (\Delta p/p)$	$\Delta G_{1+2} / (\Delta p/p)$	factor: f
group 1	11.09	10.64	-0.041718
group 2	3.25	2.36	0.242718
group 3	8.26	8.39	0.040839
group 4	10.22	11.83	0
group 5	0.71	1.52	0.671723
group 6	5.30	7.96	0.086440

Optics modulated by the space charge force

- Optics is modulated by the space charge force.
- It should be considered for the resonance corrections.



Initial test particles




- Optics modulation by space charge force was approximately calculated.
- Space charge tune shift is 0.06 for the 60π mm mrad particles.

Beam optics modulated by the space charge force

- For $dp=0\%$, space charge tune shift = -0.06
 - Start with the optics of (21.39, 21.39).
 - Multiply the scale factor to K1 values for the tune (21.33, 21.33).
- Same methods for the chromatic tune shifts of $dp/p = \pm 0.2\%$, $\pm 0.4\%$.



Beta modulation by space charge tune shift (21.33, 21.33) - (21.39, 21.39)



Space Charge Tracking Simulation

- Space charge tracking simulation program SCTR was made for J-PARC MR by Ohmi san.
- Particle in cell calculation
- Space charge potentials and kicks are calculated about 1000 elements in the circumference.
- Number of grid : $128 \times 128 \times 128$
- Grid size : $0.78 \text{ mm} \times 0.78 \text{ mm} \times 1 \text{ m}$
- Number of Macro Particles : 200,000
- Initial beam distribution
 - Transverse Gaussian rms emittance : $4\pi \text{ mm mrad}$
 - Longitudinal Bunching factor : 0.2


SCTR Space Charge Tracking Simulation
Particle Distribution



Simulation results

- Beam survival

SCTR MR Survival Ratio Collimator 65 π
 4.15e13 ppb Gauss 4 π Bf 0.2, Sext 50%



SFA	$\Delta G_{3+0} / (\Delta p/p)$	$\Delta G_{1+2} / (\Delta p/p)$	Beam Survival
group 1	11.09	10.64	0.9803
group 2	3.25	2.36	0.9916
group 3	8.26	8.39	
group 4	10.22	11.83	
group 5	0.71	1.52	0.9966
group 6	5.30	7.96	
20 mags	0	0	0.9966

Summary

- Simulation study indicated that the beam loss was reduced by correcting third order resonances for both on-momentum and off-momentum particles.
- The correction can be done with 5 groups of 4 sextupole magnets; total 20 sextupole magnets.
- We must consider the beam optics modulated by the space charge effect for the resonance correction.
- In this study we assume the error source. But...
 - We should be able to apply this method even if the error source is not known.
 - We should be able to measure the momentum deviation dependence of the resonance driving terms with the real beam.