

Evidence of Neutrino Emission from X-ray Bright Seyfert Galaxies with IceCube

Qinrui Liu^{1,2} & Shiqi Yu³
(for the IceCube collaboration)

ApJ 988 (2025) 1, 141 & ApJL 1000 (2026) 2, L37


1. Department of Physics, Simon Fraser University, Burnaby, BC V5A 1S6, Canada

2. Arthur B. McDonald Canadian Astroparticle Physics Research Institute, Kingston, ON K7L 3N6, Canada

3. Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA



Introduction

- A nearby Seyfert Galaxy, NGC 1068 has been identified as the most significant high-energy neutrino source (4.2σ) by IceCube [1].
- It sits in a Compton-thick environment with a high level of star formation. It is **dim in gamma-rays**, but is **bright in X-rays**.
- Neutrino emission at 1.5 - 15 TeV, **medium energies** for IceCube neutrino detection. **Soft best-fit spectrum** $E^{-3.2}$.
- Can we find more similar sources and examine models with current data?




Disk-corona Models

- AGN cores** optically thick for GeV-TeV gamma rays are one of the best neutrino source candidates [2,3].
- Acceleration of ions can happen via stochastic &/or magnetic reconnection processes.
- In Seyfert galaxies, a **magnetized corona** above the disk can be formed due to accretion and magnetic dissipation, providing sufficient targets for neutrino production and gamma-ray absorption.
- Disk-corona models can accommodate the total neutrino flux at medium energies $\mathcal{O}(\text{TeV}) - \mathcal{O}(10 \text{ TeV})$.
- Intrinsic X-ray luminosity** is a proxy for neutrino luminosity.



Method

Source Selection




- Seyfert Galaxies in the BAT AGN Spectroscopic Survey (BASS), which are bright in the intrinsic 2-10 keV X-ray flux.
- Northern Sky: 27 (+NGC 1068) sources.**
- Southern Sky: 14 sources**

Analyses

- Catalog Search:**
 - Fit the disk-corona model (optimistic scenario).
 - Fit the single-power-law spectrum.
- Stacking Search:**
 - Test the accumulated emission and improve the sensitivity if the model is correct.
 - Sources are weighted based on the predicted signal event number from each source.
- Binomial Test:**
 - Examine the significance of observing excesses from a subset of sources without assuming relative contributions.
- Data:**
 - Northern Sky:** same up-going track selection as used in the NGC 1068 identification analysis [1], with more data (9 + 1.7 yr)
 - Southern Sky:** Enhanced starting track event selection (9.5 yr) [4], which effectively suppresses the down-going atmospheric background.

On-going Follow-up



Combining multiple data samples (**tracks + cascades**) for an all-flavor neutrino emission study to increase the statistics and thus enhance the sensitivity.



The stacking search in the Southern Sky is expected to reach above 3σ [5].

Results



Northern Hemisphere



- 3 sources** have local significance above 3σ .
- Excess from **CGCG 420-015** and **NGC 4151** (excl. NGC 1068) at 2.7σ from the binomial test.
- No significant emission** in the stacking search (excl. NGC 1068). The upper limit constrains the collective emission to 30% of the prediction.

Southern Hemisphere

- 3σ excess in the stacking search (excl. Centaurus A).
- The top source - **Circinus Galaxy** has 3.1σ (power-law) and 2.7σ (model) (post-trial 1.8σ)



Takeway

- There should exist a **subset of sources** that can be described with an optimistic scenario of disk-corona models **similar to NGC 1068**.
- Moderate scenarios of disk-corona models will be testable with **next-generation neutrino telescopes**.
- In environments with high levels of column densities, there is a large **uncertainty** in the estimation of the **intrinsic X-ray flux**, which can significantly change neutrino emission expectations.
- Current data start to be able to test models. To find more sources and verify models, we need more **precise studies on the multi-wavelength emission** of the sources.
- More and more **X-ray-bright, gamma-ray-suppressed AGN** emerge as promising neutrino emitters. Northern Sky X-ray AGN binomial test: **11 sources 3.3σ** . See refs [6, 7].

References

- IceCube Collaboration, *Science* 378 (2022) 6619, 538-543
- Murase, Kimura & Meszaros, *125 (2020) 1, 011101*
- Kheirandish, Murase & Kimura, *ApJ* 922 (2021) 1, 45
- IceCube Collaboration, *PRD* 110 (2024) 2, 022001
- Manish Khanal et al. *APS Global Physics Summit 2026*
- IceCube Collaboration, *ApJ* 981 (2025) 2, 131
- IceCube Collaboration, *ApJL* 1000 (2026) 1, L26