No Shift, Sherlock

Shota Komatsu

Based on an upcoming paper with

José Calderón Infante (Caltech), Lucía Córdova (Amsterdam), Irene Valenzuela (CERN/IFT Madrid)

Swampland and bootstrap

- Swampland Program: Quantum gravity has imprints on IR EFT

  "In quantum gravity, not every low-energy EFT is allowed!"

- Two famous examples:

  No free parameter conjecture (absence of (-1)-form symmetry):

  All "parameters" in quantum gravity = VEV of some fields: $g \sim \langle \varphi \rangle$

  In string theory, $g_s = e^{\langle \phi \rangle}$

  No global symmetry conjecture :

  No global symmetry in quantum gravity: either gauged or broken in UV.

Swampland and bootstrap

- Swampland Program: Quantum gravity has imprints on IR EFT

  "In quantum gravity, not every low-energy EFT is allowed!"

- Two famous examples:

  No free parameter conjecture (absence of (-1)-form symmetry):

  All "parameters" in quantum gravity = VEV of some fields: $g \sim \langle \varphi \rangle$

  In string theory, $g_s = e^{\langle \phi \rangle}$

  No global symmetry conjecture :

  No global symmetry in quantum gravity: either gauged or broken in UV.

- More quantitative conjectures with possible implications to BSM

  Weak gravity conjecture, infinite distance conjecture,…

No global symmetry in quantum gravity

- One of the most (supposedly) solid swampland conjectures.

Coleman, Gidding, Strominger, Kallosh, Linde, Linde, Susskind, …. Banks, Seiberg,…+ many others

No global symmetry in quantum gravity

- One of the most (supposedly) solid swampland conjectures.

Coleman, Gidding, Strominger, Kallosh, Linde, Linde, Susskind, …. Banks, Seiberg,…+ many others

- Various indirect arguments:
  perturbative string theory, black hole entropy, etc….

- Symmetry violation from wormholes.

….Chen Lin, Hsin, Iliesiu, Yang, Milekhin, Tajdini, Belin, de Boer, Nayak, Sonner, Bah, Chen, Maldacena….

$$\overline{AA^*} \supset$$



No global symmetry in quantum gravity

- An argument based on AdS/CFT [Harlow, Ooguri 19]



Global symmetry in AdS QG is incompatible with entanglement wedge reconstruction (EWR).

No global symmetry in quantum gravity

- An argument based on AdS/CFT [Harlow, Ooguri 19]



Global symmetry in AdS QG is incompatible with entanglement wedge reconstruction (EWR).

- EWR works only in the semiclassical gravity regime, expected to be modified by stringy / QG corrections.

- But no-global-symmetry conjecture is believed to be valid beyond semiclassical GR.

- Can we prove no global symmetry conjecture in full QG regime by translating it to a CFT statement using AdS/CFT?

No global symmetry in AdS = ?? in CFT

- Well-known fact about AdS/CFT:

  Gauge symmetry in AdS QG $\leftrightarrow$ Conserved current on CFT

$$A_\mu \leftrightarrow J_\mu$$

- Global symmetry in AdS QG $\leftrightarrow$

  Global symmetry in CFT without conserved current

$$\mathcal{O}_Q \quad \rightarrow \quad e^{i\alpha Q}\mathcal{O}_Q \qquad \text{but no } J_\mu$$

No global symmetry in AdS = ?? in CFT

- Well-known fact about AdS/CFT:

  Gauge symmetry in AdS QG $\leftrightarrow$ Conserved current on CFT

  $$A_\mu \leftrightarrow J_\mu$$

- Global symmetry in AdS QG $\leftrightarrow$

  Global symmetry in CFT without conserved current

  $$\mathcal{O}_Q \quad \rightarrow \quad e^{i\alpha Q}\mathcal{O}_Q \qquad \text{but no } J_\mu$$

- **CFT counterpart of the no-global symmetry conjecture:**

  CFTs with stress tensor $T_{\mu\nu}$ and global symmetry G but no conserved current J_μ must be inconsistent.

No global symmetry in AdS = ?? in CFT

CFTs with stress tensor $T_{\mu\nu}$ and global symmetry G but no conserved current J_μ must be inconsistent.

- Clear CFT statements, but not necessarily easy to prove.
  - How do we characterize CFT with global symmetry without using conserved current?
  - As long as we look at finite number of correlators, it is difficult to distinguish "accidental" selection rules from genuine global symmetry.
  - Precisely speaking, there are counterexamples (e.g. multiple copies of Maxwell theories). $J_\mu = A^{\nu,(i)}\partial_\mu A_\nu^{(j)} - A^{\nu,(j)}\partial_\mu A_\nu^{(i)}$

- We don't know (yet) how to prove this in general. But there is actually one setup in which we can make progress.

Shift symmetry is easier

- Shift symmetry: theory is invariant under $\phi \to \phi + c$
- Clear distinction between gauged and global symmetries.
  Global shift symmetry: massless scalar
  Gauged shift symmetry: Higgs mechanism, massive scalar, no symmetry

Shift symmetry is easier

- Shift symmetry: theory is invariant under $\phi \to \phi + c$
- Clear distinction between gauged and global symmetries.
  Global shift symmetry: massless scalar
  Gauged shift symmetry: Higgs mechanism, massive scalar, no symmetry

CFT counterpart of shift symmetry in AdS

- Shift symmetry also changes the boundary value of ϕ: $\phi_{\rm bdy} \to \phi_{\rm bdy} + c$
- In AdS/CFT, $\phi_{\rm bdy}$ is exactly marginal coupling constant in CFT.

$$Z_{\phi_{\rm bdy}} = \left\langle e^{-\int \phi_{\rm bdy} \mathcal{O}} \right\rangle$$

$$\phi : \text{massless} \quad \mapsto \quad \mathcal{O} : \text{marginal } (\Delta = d)$$

Shift symmetry is easier

- Shift symmetry: theory is invariant under $\phi \to \phi + c$
- Clear distinction between gauged and global symmetries.
  Global shift symmetry: massless scalar
  Gauged shift symmetry: Higgs mechanism, massive scalar, no symmetry

CFT counterpart of shift symmetry in AdS

- Shift symmetry also changes the boundary value of ϕ: $\phi_{\rm bdy} \to \phi_{\rm bdy} + c$
- In AdS/CFT, $\phi_{\rm bdy}$ is exactly marginal coupling constant in CFT.

$$Z_{\phi_{\rm bdy}} = \left\langle e^{-\int \phi_{\rm bdy} \mathcal{O}} \right\rangle$$

$$\phi : \text{massless} \quad \mapsto \quad \mathcal{O} : \text{marginal } (\Delta = d)$$

- So shift symmetry in AdS corresponds to

  CFT which is invariant under the change of exactly marginal coupling constant $\phi_{\rm bdy} \to \phi_{\rm bdy} + c$

Shift symmetry is easier

- Shift symmetry: theory is invariant under $\phi \to \phi + c$
- Clear distinction between gauged and global symmetries.
  Global shift symmetry: massless scalar
  Gauged shift symmetry: Higgs mechanism, massive scalar, no symmetry

CFT counterpart of shift symmetry in AdS

- Shift symmetry also changes the boundary value of ϕ: $\phi_{\rm bdy} \to \phi_{\rm bdy} + c$
- In AdS/CFT, $\phi_{\rm bdy}$ is exactly marginal coupling constant in CFT.

$$Z_{\phi_{\rm bdy}} = \left\langle e^{-\int \phi_{\rm bdy} \mathcal{O}} \right\rangle$$

$$\phi : \text{massless} \quad \mapsto \quad \mathcal{O} : \text{marginal } (\Delta = d)$$

- So shift symmetry in AdS corresponds to

  CFT with a "trivial conformal manifold" (parametrized by $\phi_{\rm bdy}$), on which no CFT data changes

  or equivalently,

  CFT with an exactly marginal operator $\mathcal{O}$ and the deformation by $\mathcal{O}$ leaves all the CFT data invariant.

CFT counterpart of no shift in AdS

CFT "No Shift" conjecture

CFTs with a "trivial conformal manifold" (parametrized by $\phi_{\rm bdy}$), on which no CFT data changes are inconsistent.

or equivalently,

In CFT with an exactly marginal operator $\mathcal{O}$, the deformation by $\mathcal{O}$ always changes some CFT data.

CFT counterpart of no shift in AdS

CFT "No Shift" conjecture

CFTs with a "trivial conformal manifold" (parametrized by $\phi_{\rm bdy}$), on which no CFT data changes are inconsistent.

or equivalently,

In CFT with an exactly marginal operator $\mathcal{O}$, the deformation by $\mathcal{O}$ always changes some CFT data.

* It is important to consider CFT with stress tensor. Without stress tensor, there **are** examples of trivial conformal manifold.
(More on next slide)

Examples of trivial conformal mfd without $T_{\mu\nu}$

- Example 1: CFT with symmetry G and conformal defect breaks it down to H.

Drukker, Kong, Sakkas, ….

→ Defect conformal manifold G/H , every point on the manifold is equivalent.

$$\partial_\mu J^\mu = \tau(x_\parallel)\delta^{d-p}(x_\perp)$$

Tilt operator, $\Delta = p$, marginal



Defect

Defect

Symmetry operator
$(\phi \to \phi + c)$

Defect + $\exp\left(+c\int\tau\right)$

Examples of trivial conformal mfd without $T_{\mu\nu}$

- Example 1: CFT with symmetry G and conformal defect breaks it down to H. Drukker, Kong, Sakkas, ….

→ Defect conformal manifold G/H , every point on the manifold is equivalent.

$$\partial_\mu J^\mu = \tau(x_\parallel)\delta^{d-p}(x_\perp)$$

Tilt operator, $\Delta = p$, marginal

Example: BPS Wilson loop in $\mathcal{N} = 4$ SYM

$$\mathrm{Tr}\left[\mathrm{P}\exp\left(\int dt\, iA_\mu \dot{x}^\mu + \Phi_6|\dot{x}|\right)\right]$$

→ Choosing any other scalar Φ_I gives essentially the same defect. Trivial conformal manifold SO(6)/SO(5)

Examples of trivial conformal mfd without $T_{\mu\nu}$

- Example 1: CFT with symmetry G and conformal defect breaks it down to H. Drukker, Kong, Sakkas, ….

  → Defect conformal manifold G/H , every point on the manifold is equivalent.

$$\partial_\mu J^\mu = \tau(x_\parallel)\delta^{d-p}(x_\perp)$$

Tilt operator, $\Delta = p$, marginal

Example: BPS Wilson loop in $\mathcal{N} = 4$ SYM

$$\mathrm{Tr}\left[\mathrm{P}\exp\left(\int dt\, iA_\mu \dot{x}^\mu + \Phi_6|\dot{x}|\right)\right]$$

→ Choosing any other scalar Φ_I gives essentially the same defect. Trivial conformal manifold SO(6)/SO(5)

- Example 2: Free shift symmetric scalar in AdS without gravity.

  → obviously UV complete, generalized free field CFT, no stress tensor.

Plan

1. CFT argument for no shift symmetry
2. Generalization to higher form
3. Modular bootstrap
4. Conclusion

Plan

1. CFT argument for no shift symmetry
2. Generalization to higher form
3. Modular bootstrap
4. Conclusion

General strategy

- A proof by contradiction in two steps:

- **Step 0**: Assume ∃CFT with a trivial conformal manifold

- **Step 1**: Use a CFT bootstrap argument to show one can construct "identity conformal interface" for any two points on a trivial conformal manifold.

- **Step 2**: Show that the existence of the identity conformal interface between nearby CFTs contradicts conformal perturbation theory

Step 1: Trivial conformal mfd implies identity interface

Conformal interfaces

- Conformal interface $SO(d+1,1) \to SO(d,1)$



- Needs to satisfy various bulk-interface crossing eq



- Interface crossing eq: Bulk CFT data ⇒ Interface CFT data

Conformal interfaces

- Conformal interface $SO(d+1,1) \to SO(d,1)$



- Needs to satisfy various bulk-interface crossing eq



- Interface crossing eq: Bulk CFT data ⇒ Interface CFT data
- Microscopic construction of interfaces: imposing bc to bulk fields, coupling to localized dof….
- In the modern bootstrap, sol's to crossing = conformal interface

Trivial conformal mfd implies identity interface

- On trivial conformal manifold, CFT_1 and CFT_2 share the same bulk CFT data by assumption.
- Interface crossing eq: Bulk CFT data $\Rightarrow$ Interface CFT data

Trivial conformal mfd implies identity interface

- On trivial conformal manifold, CFT_1 and CFT_2 share the same bulk CFT data by assumption.
- Interface crossing eq: Bulk CFT data $\Rightarrow$ Interface CFT data
- 1-to-1 correspondence between sol's for interface bootstrap: interfaces within CFT_1 $\leftrightarrow$ interfaces connecting $CFT_{1,2}$



- In particular, it implies the "identity interface" between $CFT_{1,2}$ that corresponds to inserting nothing in CFT_1



Comment

- I have shown that, if no CFT data changes, one can construct “identity interface” between $CFT_{1,2}$.
- This sounds obvious, but it is actually a key point of the proof.
- **Contrapositive:** If there is no identity interface, CFT data changes.

Comment

- I have shown that, if no CFT data changes, one can construct “identity interface” between $CFT_{1,2}$.
- This sounds obvious, but it is actually a key point of the proof.
- **Contrapositive:** If there is no identity interface, CFT data changes.
- **Gravity counterpart:** If there is no tensionless domain wall, shift symmetry is broken.



- There are arguments pointing out difficulty of constructing such domain walls within gravitational EFT due to large back reaction.

Bah, Jefferson, Roumpedakis, Waddleton,…

- But such arguments do not tell anything about shift-symmetry breaking. So the claim is highly non-trivial from gravity viewpoint.
- In AdS_3/CFT_2, non-topologicalness $\leftrightarrow$ symmetry breaking (Later)

modular bootstrap

Step 2: Identity interface contradicts conformal perturbation theory

Identity interface for nearby CFTs

- Let's analyze the "identity interface" between nearby CFTs.

$$CFT_1 \quad | \quad CFT_2 \quad = CFT_1 + \delta\phi \int_{x\perp \geq 0} \mathcal{O}$$

- Used a bootstrap argument to show its existence without knowing microscopic constructions.
- But we do know
  - In the limit $CFT_{1,2}$ coincide, it reduces to "inserting nothing".
  - Infinitesimally, $CFT_2 = CFT_1 + \delta\phi \int \mathcal{O}$

Identity interface for nearby CFTs

- Let's analyze the "identity interface" between nearby CFTs.

$$CFT_1 \;\big|\; CFT_2 \;\big| \;= CFT_1 + \delta\phi \int_{x_\perp \geq 0} \mathcal{O}$$

- Used a bootstrap argument to show its existence without knowing microscopic constructions.
- But we do know
  - In the limit $CFT_{1,2}$ coincide, it reduces to "inserting nothing".
  - Infinitesimally, $CFT_2 = CFT_1 + \delta\phi \int \mathcal{O}$
- So infinitesimally,

"Identity interface" = $CFT_1 + \delta\phi \int_{x_\perp \geq 0} \mathcal{O} + \delta\rho \int_{x_\perp = 0} \mathcal{O}'$

Additional deformation localized at interface

- $\mathcal{O}'$ can be a local op. on the interface or some non-local expression on the interface.

Identity interface for nearby CFTs

- So infinitesimally,
  "Identity interface" = $CFT_1 + \delta\phi \int_{x_\perp \geq 0} \mathcal{O} + \delta\rho \int_{x_\perp = 0} \mathcal{O}'$

  Additional deformation localized at interface

- $\mathcal{O}'$ can be a local op. on the interface or some non-local expression on the interface.

- No deformation away from the interface is allowed since it will violate the stress tensor conservation in the bulk. $\partial_\mu T^{\mu\nu} \neq 0$

- (In CFT without stress tensor, such deformations are allowed)
  [Herzog, Shamir]

Identity interface is inconsistent

"Identity interface" =

$$CFT_1 + \delta\phi \int_{x_\perp \geq 0} \mathcal{O} + \delta\rho \int_{x_\perp = 0} \mathcal{O}'$$

- Consider "Displacement operator" D.

$$\partial_\mu T^{\mu\perp} = D(x_\parallel)\delta^{d-1}(x_\perp)$$

- For the deformation above, one can show [SK, Kusuki, Meineri, Ooguri '25]

$$D = \delta\phi\,\mathcal{O} + \delta\rho\,\partial_\perp \mathcal{O}' + (\text{higher order})$$

Identity interface is inconsistent

"Identity interface" =

$$CFT_1 + \delta\phi \int_{x_\perp \geq 0} \mathcal{O} + \delta\rho \int_{x_\perp = 0} \mathcal{O}'$$

- Consider "Displacement operator" D.

$$\partial_\mu T^{\mu\perp} = D(x_\parallel)\delta^{d-1}(x_\perp)$$

- For the deformation above, one can show [SK, Kusuki, Meineri, Ooguri '25]

$$D = \delta\phi\,\mathcal{O} + \delta\rho\,\partial_\perp \mathcal{O}' + (\text{higher order})$$

- But for the identity interface, $D = 0$.

- The only way to get $D = 0$ is to set $\mathcal{O} \propto \partial_\perp \mathcal{O}'$. But $\mathcal{O}$ is a conformal primary. **Contradiction.**

- Thus, the identity interface contradicts with conformal perturbation. Hence, the shift symmetry must be broken.

Plan

1. CFT argument for no shift symmetry
2. Generalization to higher form
3. Modular bootstrap
4. Conclusion

Generalization to higher form symmetry

- 1-form symmetries of Maxwell theory.

$$dF = 0\,, \quad d * F = 0$$

- In QG, they are expected to be violated by the presence of charged particles.

Related to charge completeness conjecture

- Can we show the absence of 1-form symmetries of Maxwell theory in AdS QG?

Generalization to higher form symmetry

- If there was (electric) 1-form symmetry, one can construct the codim-2 topological interface in AdS.



- By pushing them to boundary, one gets a monodromy defect (co-dimension 2 + codimension 1)



- 1-form symmetry in the bulk predicts that the monodromy defect is topological.

CFT counterpart of no 1-form symmetry in AdS

CFT "No 1-form" conjecture

In CFT with stress tensor, monodromy defects cannot be topological.



Generalization to higher form symmetry



- Working in $\alpha \ll 1$ regime, one can show that the displacement operator is nonzero.

$$D \propto j$$

Contradiction to topologicalness.

- One can also argue that any local deformation at the defect (edge of the circle) cannot remedy this.

Thus, the topological monodromy defect cannot arise in CFT.

Plan

1. CFT argument for no shift symmetry
2. Generalization to higher form
3. Modular bootstrap
4. Conclusion

Bootstrap estimate on 1-form symmetry violation



- Consider a torus partition function with $e^{i\alpha \int *j}$ inserted.
- Untwisted channel (L is time)

$$I(\alpha) = \sum_a e^{-E_a L + i\alpha Q_a}$$

- Twisted channel (L is space)

$$I(\alpha) \overset{L\to 0}{\sim} e^{-E_0(\alpha)/L}$$

$E_0(\alpha)$: Energy of ground state in twisted sector

- Equating both sides and taking derivatives:

$$\partial_\alpha^2 E_0(\alpha)\big|_{\alpha=0} = \lim_{L\to 0} L e^{\frac{c}{12L}} \sum_a Q_a^2 e^{-E_a L}$$

Non-topologicalness $\leftrightarrow$ violation of 1-form symmetry (existence of charged op)

Bootstrap estimate on 1-form symmetry violation

$$\partial_\alpha^2 E_0(\alpha)\big|_{\alpha=0} = \lim_{L\to 0} L e^{\frac{c}{12L}} \sum_a Q_a^2 e^{-E_a L}$$

- $E_0(\alpha)$ can be computed by perturbation in α .

$$\int_{x_1\le x, y\le x_2} dxdy\langle j(x)j(y)\rangle = C_J \int_{x_1\le x, y\le x_2} dxdy \frac{1}{|x-y|^2} = C_J \log|x_1 - x_2|$$

$$\Rightarrow E_0(\alpha) = C_J \alpha^2 + \cdots$$

- Quantitative estimate on the violation of 1-form symmetry

$$\langle Q^2\rangle \overset{\Delta\gg 1}{\sim} C_J\sqrt{\frac{\Delta}{c}}$$

- Analysis seems generalizable for bulk 0-form symmetry.

[In progress]

Conclusion

- CFT-based argument for no shift symmetry in AdS QG.
  - Works beyond the semi-classical gravity regime.
  - But works only for shift symmetry + spontaneously broken symmetries.

- Conformal interfaces seem useful for analyzing swampland-related question. [SK, Kusuki, Meineri, Ooguri]

- Monodromy defects also seem important.

  ⇒ A systematic bootstrap analysis of monodromy defect?
  Application to weak gravity conjecture?

- In holographic theories, we expect $\langle Q^2 \rangle \overset{\Delta \geq \frac{c}{6}}{\simeq} C_J \sqrt{\frac{\Delta}{c}}$ Black hole interpretation?

- Quantitative bootstrap bound in higher dimensions?

- Other related swampland conjectures? Infinite distance conjecture?

- Non-spontaneously broken symmetries?