

# Perturbative unitarity bounds on field-space curvature in de Sitter spacetime

Tomoya Inada (UTokyo, Japan)

## Perturbative Unitarity Bounds

Theoretical constraints on EFT parameters from **unitarity** as a consistency condition


We investigate unitarity bounds for a field-space curvature model

$$\mathcal{L} = -\frac{1}{2}(\partial\phi)^2 - \frac{1}{2}(\partial\sigma)^2 - H^2\sigma^2 + \frac{1}{4f^2}\sigma^2(\partial\phi)^2$$

in **de Sitter spacetime**.


We derive the bounds using **momentum-space purity  $\gamma$**  rather than scattering amplitudes.

(following an approach developed at Cambridge!)




# The Path to the NUT Solution: A Coordinate-Free Approach

## 1 Newman–Penrose (NP) Formalism



## 2 Newman–Unti–Tamburino (NUT) Solution



# HOW CAN GRAVITON FLUCTUATIONS INDUCE A SPREAD OF THE LIGHTCONE?

Scalar field  $\phi$  coupled to thermal gravitons  $\Rightarrow$  corrections to scalar field commutator

$$P\left([\phi(x), \phi(0)] \neq 0\right) \propto \exp\left(\frac{-(\vec{x}^2 - t^2)^2}{2\text{Var}(\vec{x}^2)}\right)$$

Thermal contribution

$$\text{Var}(\vec{x}^2) = \frac{16 G_N T t^3}{3}$$

Vacuum contribution

$$\text{Var}(\vec{x}^2) \sim t^2 \log\left(\frac{t}{R}\right)$$

$$\frac{\text{Var}(\vec{x}^2)}{\langle \vec{x}^2 \rangle^2} \sim \frac{1}{t}$$

Still a perturbative effect

At  $T=300$  K,  
wait  $10^4$  yrs for the lightcone  
spread to reach a meter

Black Hole application

$$t \sim t_P \left(\frac{M}{M_P}\right)^{5/3}$$

Time duration for the lightcone  
spread to reach the BH size

# Reheating after axion inflation

Tenta Tsuji


KEK Theory Center

$$\ddot{\phi} + 3H\dot{\phi} + \Gamma_{\phi}\dot{\phi} + V'(\phi) = 0$$

thermal friction from thermal bath

## The completion of Reheating?


[2503.01228, JCAP07(2025)002]



We study this question in axion inflation set up

# Thermal Equilibrium Open EFT of de Sitter and Warm Inflation

Wenqi Yu (HKUST)



# How many Nambu–Goldstone modes?

## Goldstone theorem

SSB  $\implies$  gapless Nambu–Goldstone modes (NGMs)

But in general  $N_{\text{NGM}} \leq N_{\text{BG}}$ .

## Known reduction mechanisms

Two reduction mechanisms are known:

$$\langle [Q, Q'] \rangle \neq 0 \implies \mathcal{L}_{\text{eff}} \supset \pi \dot{\pi}' \quad (\text{BWM pairing})$$

$$[P, Q] \sim Q' \implies \pi' \sim \partial \pi \quad (\text{inverse Higgs})$$

## New mechanism: **geometric pairing**

$$[P, Q] \sim P \implies \mathcal{L}_{\text{eff}} \supset \pi \dot{\pi}'$$

A general counting formula for NGMs in many-body systems follows.