



UNIVERSITY OF MARYLAND, COLLEGE PARK



Probing Instanton Dynamics in the Pion Vector Form Factor with Gradient Flow


Vaibhav Chahar, Piotr Korcyl

Jagiellonian University, Poland

43rd International Symposium on Lattice Field Theory (Lattice 2026)

Introduction

- **QCD** is a non-Abelian gauge theory based on the $SU(3)$ color symmetry group. It describes interactions between quarks and gluons.
- The QCD vacuum is not empty, as it contains topologically non-trivial gluon field configurations.
- Instantons are localized, classical solutions of the Euclidean Yang–Mills equations.
- They describe **tunneling** between distinct classical QCD vacua.



R. Jackiw and C. Rebbi, *Phys. Rev. Lett.* 37, 172 (1976)

Wilson Flow (Gradient Flow)

- The QCD vacuum on the lattice is full of UV noise, tiny fluctuations that obscure physical, topological structures.
- Wilson flow is a process that evolves lattice gauge fields $U_\mu(x)$ along a continuous “flow time” t by solving:

$$\frac{dU_\mu(x,t)}{dt} = -g_0^2 \left\{ \frac{\partial S_g[U(t)]}{\partial U_\mu(x,t)} \right\} U_\mu(x,t), \quad U_\mu(x,0) = U_\mu(x),$$

- The gauge fields become smoother as t increases, with small-scale noise diffused away.
- Applications in Lattice Field Theory:
 - Topological Charge and Susceptibility
 - Scale Setting
 - Improve Signal to noise

Wilson Flow and Instantons


- Each instanton carries a topological charge corresponding to tunneling between degenerate QCD vacua.

$$Q = \frac{1}{32\pi^2} \int d^4x G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a$$

- At **small flow times**, UV fluctuations are suppressed, but physical long-distance structures remain intact.
- At **moderate flow**, instanton-like objects appear as **coherent lumps** in the topological charge.
- If the flow time is too long, instantons may shrink and annihilate (over-smoothing), so an **optimal range of flow time** is chosen.
- In this work, we study hadronic matrix elements directly at finite Wilson flow time $t > 0$, and understand how they change as the flow time is varied.

Instantons and Pion Form Factors

- The **Instanton Liquid Model (ILM)** provides a mathematical framework to describe this complex vacuum by modeling it as an ensemble of interacting instantons and anti-instantons.
- One particularly insightful observable is the **pion electromagnetic form factor**, which probes the internal structure and dynamics of the pion as a bound state of quarks.
- The Pure Instanton Liquid Model gives a higher value of the pion form factor and does not agree with lattice qcd and experiments.
- We study the pion form factor for different values for t .




Pion Form Factors from Lattice

$$O_{\pi}(x,t) = \bar{d}(x,t) \gamma_5 u(x,t),$$

$$O_{\pi}^{\dagger}(x,t) = \bar{u}(x,t) \gamma_5 d(x,t).$$

$$C_{\pi}^{2\text{pt}}(t, \mathbf{p}) = \sum_{\mathbf{x}} e^{-i\mathbf{p} \cdot \mathbf{x}} \langle 0 | O_{\pi}(\mathbf{x}, t) O_{\pi}^{\dagger}(0, 0) | 0 \rangle.$$

$$C_{\pi}^{3\text{pt}}(t, t_{\text{sep}}, \mathbf{p}_f, \mathbf{p}_i) = \sum_{\mathbf{x}, \mathbf{y}} e^{-i\mathbf{p}_f \cdot (\mathbf{x} - \mathbf{y})} e^{-i\mathbf{p}_i \cdot \mathbf{y}} \langle 0 | O_{\pi}(\mathbf{x}, t_{\text{sep}}) J_{\mu}(\mathbf{y}, t) O_{\pi}^{\dagger}(0, 0) | 0 \rangle.$$



$$R(t_{\text{sink}}, \tau, \vec{p}, \vec{p}) = \frac{C_{3\text{pt}}^{\mathcal{O}}(\tau, \vec{p}, \vec{p})}{C_{2\text{pt}}(t_{\text{sink}}, \vec{p})} \sqrt{\frac{C_{2\text{pt}}(t_{\text{sink}} - \tau, \vec{p}) C_{2\text{pt}}(\tau, \vec{p}) C_{2\text{pt}}(t_{\text{sink}}, \vec{p})}{C_{2\text{pt}}(t_{\text{sink}} - \tau, \vec{p}) C_{2\text{pt}}(\tau, \vec{p}) C_{2\text{pt}}(t_{\text{sink}}, \vec{p})}}.$$

$$R(t_{\text{sink}}, \tau, p', p) = \frac{\langle \pi(p') | O(\tau) | \pi(p) \rangle}{2}$$

$$2\sqrt{E_{\vec{p}} E_{\vec{p}'}} \times Z_V \times \langle \pi(\vec{p}') | \mathcal{O}_V^{\mu}(0) | \pi(\vec{p}) \rangle = (p_f + p_i)_{\mu} \times f_{\pi\pi}(-q^2)$$



Lattice Setup

- The study uses publicly available PACS-CS gauge field ensemble.
- • The Fermionic part o(a)-improved Wilson action with $N_f = 2 + 1$ dynamical quarks and the gauge part is the Iwasaki gauge action.
- Bare parameters for the ensemble

β	a [fm]	k_l	L/a	T/a	c_{sw}	N_G	M_π	Z_v
1.96	0.0907(13)	0.13754	32	64	1.715	200	409.7(7)	0.7354(37)


Setting up the bare parameters

- Rather than performing ADDITIONAL simulations at vanishing quark mass, we attempt to introduce a massive renormalization and improvement scheme, extracting the relevant parameters directly from our available ensemble at $M_\pi \simeq 400$ MeV.
- Preliminary: set $c_{\text{sw}} = 1.0$. A more precise analysis requires tuning c_{sw} and c_A for example using PCAC relations.




Setting up the bare parameters

We see a significant difference between the form factors at tree-level and flow time = $5t_0$



Setting up the bare parameters



Setting up the bare parameters

- PCAC relation

$$\langle \partial_\mu A_\mu(x) \mathcal{O}(y) \rangle = 2m \langle P(x) \mathcal{O}(y) \rangle \quad \text{where, } A_\mu(x) = \bar{u}(x) \gamma_\mu \gamma_5 d(x), P(x) = \bar{u}(x) \gamma_5 d(x)$$

For simplicity, $\mathcal{O}(y) = P^\dagger(y)$

$$\frac{1}{2} (\partial_\mu^* + \partial_\mu) \langle A_\mu^I(x) P^\dagger(y) \rangle = 2m \langle P(x) P^\dagger(y) \rangle$$

$$A_\mu^I(x) = A_\mu(x) + c_A \frac{1}{2} (\partial_\mu^* + \partial_\mu) P(x)$$

We set $\mu = 0$

$$\overrightarrow{\partial_0} \langle \{ A_0(x) + c_A \overrightarrow{\partial_0} P(x) \} P^\dagger(y) \rangle = 2m \langle P(x) P^\dagger(y) \rangle$$

- With symmetric derivative $\overleftrightarrow{\partial_0} = \frac{1}{2} (\partial_0^* + \partial_0)$

$$\overrightarrow{\partial_0} \langle A_0(x) P^\dagger(y) \rangle + c_A \overrightarrow{\partial_0} \overrightarrow{\partial_0} \langle P(x) P^\dagger(y) \rangle = 2m \langle P(x) P^\dagger(y) \rangle$$

- Now we have ratio


$$m(x, y; c_A, c_{SW}, m_\pi) = \frac{\overrightarrow{\partial_0} A_d \langle A_0(x) P^\dagger(y) \rangle + c_A \overrightarrow{\partial_0} \overrightarrow{\partial_0} \langle P(x) P^\dagger(y) \rangle}{\langle P(x) P^\dagger(y) \rangle}.$$

Setting up the bare parameters



$$m(x,y;c_A,c_{SW},m_\pi)=\frac{\overline{\partial}_0\langle A_0(x)P^\dagger(y)\rangle+c_A\overline{\partial}_0\overline{\partial}_0\langle P(x)P^\dagger(y)\rangle}{\langle P(x)P^\dagger(y)\rangle}.$$

- The ratio is plotted with c_A
- Choose the points in the cone that lie on the hyperdiagonal of the lattice.
- The cone is defined using **physical distances**.
- Then choose the points that lie inside the shaded frustum.
- The larger the angle θ , the more points to consider
- Calculate the variance of $m(y;c_{SW},c_A)$ using points lying in the frustum of various c_{SW} and M_π .


$$\text{Var}(m)=\frac{1}{N-1}\sum_{y\in\mathcal{C}}(m(y)-\overline{m})^2$$



Results




Results: preliminary



Results: preliminary



- Once we fix c_{sw} we will know all necessary parameters and could estimate the form factor.
- The algorithm can be applied at $t > 0$.



Preliminary results for parameter tuning at $t = 0$, for 20 gauge configurations.

Results: preliminary

- Can matrix elements at large flow time be extracted using the same procedure as at $t = 0$?
- Model calculations suggest that the standard extraction procedure remains valid at finite flow time.
- After tuning κ for each flow time, the pion mass is consistent across flows. However, excited-state contributions are enhanced at larger flow times.



Summary and Outlook

- Tune the parameters c_{sw} , κ
- After tuning the parameters, calculation of form factors with higher values of flow time will be used.
- Extrapolation to $a \rightarrow 0$
- Use more lattice ensembles for broader results.

Thank you