

# Drift Properties of Noble Gases in Negative Ion Drift TPCs


---

ELIZABETH TILLY – WELLESLEY COLLEGE

*WORK CARRIED OUT DURING PHD AT THE UNIVERSITY OF NEW MEXICO*

CPAD – OCTOBER 2026

# Directionality for Rare-Event Searches





S. Vahsen et al Annu. Rev. Nucl. Part. Sci. 2021. 71 189-224 [[2102.04596 \(arxiv.org\)](https://arxiv.org/abs/2102.04596)]

# Noble Gases as Targets

*Far from a comprehensive list...*

- **Gaseous He, Ar, Xe and CF<sub>4</sub> for WIMP DM searches**
  - J.F. Castel *et al.* JINST **19** C05029 (2024)
  - F.D. Amaro *et al.* Particles **4** 343-353 (2021)
  - S.E. Vahsen *et al.* (2020) arxiv:2008.12587
  - C. Deaconu *et al.* Phys. Rev. D **95** 122002 (2017)
  - K. Nakamura *et al.* JINST **7** C02023 (2012)
- **Gaseous Xe for neutrinoless double beta decay**
  - K. Mistry, D. R. Nygren, (2025) arxiv:2512.19290
  - I.G. Irastorza *et al.* JCAP **01** 033 (2016)
- **Gaseous He, Ar, Xe, CF<sub>4</sub> in Migdal Effect searches**
  - D. Yi *et al.* Nature **649**, 580–583 (2026)
  - H.M. Araújo *et al.* (MIGDAL collaboration), Astropart. Phys. **151** 102853 (2023)
  - K.D. Nakamura *et al.* PTEP **1** 013C01 (2021)
- **Gaseous Ar in collider-based neutrino experiments**
  - R.M. Amarinei, *et al.* JINST **18** P12001 (2023)
  - A. Abed Abud, *et al.* (2022) arXiv:2203.06281



*And the list goes on...*

# Negative Ion Drift Low-Pressure Gas TPCs


Lower diffusion








Longer Tracks



Better Directional Resolution



# Advantages of Negative Ion Drift (NID)



| 150 Torr CF <sub>4</sub> + X torr CS <sub>2</sub> |        |
|---------------------------------------------------|--------|
| CS <sub>2</sub> (Torr)                            | σ(μm)  |
| 0                                                 | ~400   |
| 2.9                                               | 133.53 |
| 4.2                                               | 126.10 |
| 5.4                                               | 125.09 |

| 45 Torr CF <sub>4</sub> + X torr CS <sub>2</sub> |          |
|--------------------------------------------------|----------|
| 0                                                | ~550     |
| 4                                                | ~150-200 |

Gains for these tracks are on the order of 10<sup>5</sup>!

# Thermal Drift

- At **low electric fields**, ions are **nearly in thermal equilibrium** with the gas. As a consequence:

- Diffusion width evolves according to the following

$$\sigma^2 = \frac{2kT}{e} \frac{L}{E}$$

- Mobility is constant with E/P


$$\mu = \frac{v_d}{E}$$

- At higher fields (i.e. more energetic systems):

- Mobility and diffusion are **larger at higher fields**
- Diffusion becomes **axially dependent**



$v_d$  = drift velocity,  $\mu$  = mobility,  $E$  = Electric Field,  $e$  = ion charge,  
 $kT$  = thermal energy,  $L$  = drift length,  $\sigma$  = diffusion width



Plots created using [desmos.com](https://desmos.com)



Electric Field

# Optimizing/Testing Different Gas Mixtures

GOAL: Explore gas mixtures with high quantities of the **target gas** maintaining **negative ion benefits**

- Target Gases: **Xe, Ar, He, CF<sub>4</sub>** NID gases: **SF<sub>6</sub> and CS<sub>2</sub>**
  - Of interest to Migdal searches, Directional Dark Matter searches, Neutrino Studies, etc
- Properties of interest for Noble Gas NID TPCs:
  - **Diffusion = Track broadening from stochastic motion within the gas** → NID is narrower
  - **Mobility = Proportionality constant between drift velocity and field** → NID is slower
  - **Gain = Stable operation and amplification capability** → electron stripping is required for avalanche amplification in NID but NID gases increase stability
  - **Attachment Length = Mean distance before electrons are captured in the gas** → Must have short attachment length


# The Detector





N.S. Phan *et al* 2017 *JINST* **12** P02012

\*\* THGEM we used is 400  $\mu$ m thick and has 10x10 cm active area, a pitch of 0.5mm, and a hole diameter of 0.3mm


# Relative Gains



- Strongly electronegative SF<sub>6</sub> and CS<sub>2</sub> ions must be stripped before amplification
- CS<sub>2</sub>-based mixtures had higher gains, in general, than SF<sub>6</sub>-based mixtures
- $\frac{\sigma}{\mu} \approx 20\%$  for CS<sub>2</sub> and 25% for SF<sub>6</sub>
- Stable operation resolving 5.9 keV peak from <sup>55</sup>Fe with as little as 3 Torr SF<sub>6</sub> and 5 Torr CS<sub>2</sub>!



# Laser-based Measurements



# Mobility

$$\mu_o = \frac{v_d}{E} \frac{N}{N_o}$$


- $N$  and  $N_o$  are gas density in detector and at STP,  $v_d$  is drift velocity,  $E$  is Electric Field

- Mobilities are roughly 2 orders of magnitude lower than in the pure noble gases
- Many minority species were noted with mobilities around ~10-20% of the primary species

**Mobility at 600 V/cm and 3 Torr SF<sub>6</sub>**

| Gas | P <sub>total</sub><br>(Torr) | ED* $\mu_o$<br>(cm <sup>2</sup> s <sup>-1</sup> V <sup>-1</sup> ) | NID $\mu_o$<br>(cm <sup>2</sup> s <sup>-1</sup> V <sup>-1</sup> ) |
|-----|------------------------------|-------------------------------------------------------------------|-------------------------------------------------------------------|
| Xe  | 20                           | 188                                                               | <b>0.77</b>                                                       |
| Ar  | 50                           | 325                                                               | <b>1.64</b>                                                       |
| He  | 200                          | 796                                                               | <b>9.03</b>                                                       |

\* S. Biagi (1995). <https://magboltz.web.cern.ch/magboltz/>



# Mobility


$$\mu_o = \frac{v_d}{E} \frac{N}{N_o}$$

- $N$  and  $N_o$  are gas density in detector and at STP,  $v_d$  is drift velocity,  $E$  is Electric Field

- Mobilities are roughly 2 orders of magnitude lower than in the pure noble gases
- Many minority species were noted with mobilities around ~10-20% of the primary species

| Mobility at 600 V/cm and 3 Torr SF <sub>6</sub> |                           |                                                                |                                                                |
|-------------------------------------------------|---------------------------|----------------------------------------------------------------|----------------------------------------------------------------|
| Gas                                             | P <sub>total</sub> (Torr) | ED* $\mu_o$ (cm <sup>2</sup> s <sup>-1</sup> V <sup>-1</sup> ) | NID $\mu_o$ (cm <sup>2</sup> s <sup>-1</sup> V <sup>-1</sup> ) |
| Xe                                              | 20                        | 188                                                            | 0.77                                                           |
| Ar                                              | 50                        | 325                                                            | 1.64                                                           |
| He                                              | 200                       | 796                                                            | 9.03                                                           |

\* S. Biagi (1995). <https://magboltz.web.cern.ch/magboltz/>




# Waveform Shape


Fitting Functions Used to model Diffusion width,  $\sigma$  (and attachment length,  $\lambda = 1/L$ )

- Gaussian:  $G(t) = \exp\left[-\frac{(t-m)^2}{\sqrt{2\sigma^2}}\right]$
- Exponentially Modified Gaussian:  $EMG(t) = \frac{L}{2} \exp\left[\frac{L}{2}(2m + L\sigma^2 - 2t)\right] \operatorname{erfc}\left[\frac{m+L\sigma^2-t}{\sqrt{2\sigma^2}}\right]$
- Step + Gaussian:  $SG(t) = \frac{h}{2}\left(1 - \operatorname{erfc}\left[\frac{x-m}{\sqrt{2\sigma^2}}\right]\right) + G(t, m, \sigma)$


25+25 Torr Ar+CS<sub>2</sub>



40+10 Torr Ar+CS<sub>2</sub>



25+25 Torr Ar+SF<sub>6</sub>




# Longitudinal Diffusion

- $\sigma_z = D'_L \sqrt{58.3 \text{ cm}}$ 
  - $D'_L$  is longitudinal diffusion coefficient used by Garfield and Magboltz. It relates to the normal Diffusion coefficient,  $D$ , by  $D' = \sqrt{2D/v_d}$
- Diffusion can be suppressed by ~ an order of magnitude with as little as 1 Torr  $\text{SF}_6$  or 5 Torr  $\text{CS}_2$
- In many cases it's nearly thermal!

| Diffusion Coefficient at 600 V/cm and 3 Torr $\text{SF}_6$ |                           |                                                            |                                                          |
|------------------------------------------------------------|---------------------------|------------------------------------------------------------|----------------------------------------------------------|
| Gas                                                        | P <sub>total</sub> (Torr) | ED $D'_{\text{long}}^*$ ( $\mu\text{m}/\sqrt{\text{cm}}$ ) | NI $D'_{\text{long}}$ ( $\mu\text{m}/\sqrt{\text{cm}}$ ) |
| Xe                                                         | 20                        | 941                                                        | <b>165</b>                                               |
| Ar                                                         | 50                        | 1143                                                       | <b>154</b>                                               |
| He                                                         | 200                       | 1063                                                       | <b>~77</b>                                               |

\* S. Biagi (1995). <https://magboltz.web.cern.ch/magboltz/>




# Attachment Length in CS<sub>2</sub> Mixtures

- Attachment could only be measured in a subset of the data
- CS<sub>2</sub> + Ar and CF<sub>4</sub> mixtures had large attachment lengths!

Consequences of high attachment for tracking detectors:

1. Exponentially modified waveforms along drift axis
2. “Mixed Drift” waveforms if drift length too short
  - Leads to waveform distortion
  - Leads to mixed gain for ED and NID components of the waveform



# Conclusions


- Negative ion drift properties in mixtures of Xe, Ar, He, and CF<sub>4</sub> with either SF<sub>6</sub> or CS<sub>2</sub> have been measured with very low partial pressures of the NID component
- Stable gain for Fe-55 peak noted in many of the mixtures
- Diffusion was suppressed by ~1 order of magnitude
- Mobility was slowed by 2 orders of magnitude
- In some cases, attachment length could also be measured!

*Special thanks to my collaborators on this project:*  
*Dinesh Loomba and Drew Gibson and the UNM group*

| Diffusion Coefficient at 600 V/cm<br>and 3 Torr SF <sub>6</sub> |                              |                                                |                                   |
|-----------------------------------------------------------------|------------------------------|------------------------------------------------|-----------------------------------|
| Gas                                                             | P <sub>total</sub><br>(Torr) | ED D' <sub>long</sub> <sup>*</sup><br>(μm/√cm) | NI D' <sub>long</sub><br>(μm/√cm) |
| Xe                                                              | 20                           | 941                                            | 165                               |
| Ar                                                              | 50                           | 1143                                           | 154                               |
| He                                                              | 200                          | 1063                                           | ~77                               |

\* S. Biagi (1995). <https://magboltz.web.cern.ch/magboltz/>

Fermilab/DOE  
GIRA Award  
Awarded to E.Tilly



# Backups

---

# Notable Papers Regarding NID


- C.J. Martoff *et al* 2000 *NIMA* **440** 335 ([https://doi.org/10.1016/S0168-9002\(99\)00955-9](https://doi.org/10.1016/S0168-9002(99)00955-9))
  - The original NID Paper
- N.S. Phan *et al* 2017 *JINST* **12** P02012 (<https://doi.org/10.1088/1748-0221/12/02/P02012>)
  - Established SF<sub>6</sub> as an ideal NID gas
- J.B.R. Battat *et al* 2017 *AstroPart Phys* **91** 65 (<https://doi.org/10.1016/j.astropartphys.2017.03.007>)
  - DRIFT Experiment results using minority NI species to fiducialize leading to 54 day zero-background run

# The Future of these studies

1. Paper coming soon!
2. Development of a novel low-pressure NI TPC
  - 10cm x 10cm 2D strips at 200  $\mu\text{m}$  pitch
    - Gas Electron Multiplier (GEM) at 140  $\mu\text{m}$  pitch to amplify the signal from the tracks
  - Front-end electronics assembled, adapted, tested and characterized with collaborators at Boston University




Ed Kearns Alex Madorsky Lila Smith




# Light Yield in NID

50 Torr CF4:




45 Torr CF4 +  
4.3 Torr CS2



Courtesy of D. Loomba RD51 Collab. Meeting 6/14/22

# Minority Species Area





# Measuring Electron Energy

- Under a very simple assumption:

$$\frac{A_{SF_5^-}}{A_{SF_6^-}} = \frac{\sigma_{SF_5^-}}{\sigma_{SF_6^-}}$$

- We can use the area ratios to get approximate electron energies (pre-capture)
  - Note: different electronegativities mean these two species probably have different gains
    - Thus, this is more likely an upper limit on the energy



# Dealing with Water

- Small amounts of water in detector lead to 'water-shelves' on left side of waveform
- Compromises measurement of minority area and diffusion sigma of main peak
- Include 1 or 2 step functions in fit:

$$S_{\pm}(x) = A \frac{h_{\pm}}{2} \left( 1 - \operatorname{erf} \left( \pm \frac{x - \mu}{\sqrt{2}\sigma} \right) \right)$$



$$G(x) = A \exp \left( -\frac{(x - \mu)^2}{2\sigma^2} \right)$$

- For minority area:

$$F(x) = S_{+}(x) + S_{-}(x) + G(x)$$

- For main peak diffusion:

$$F(x) = S_{+}(x) + G(x)$$




# Exponentially Modified Gaussian


$$F(x) = \frac{A\lambda}{2} \exp\left(\frac{\lambda}{2}(2\mu + \lambda\sigma^2 - 2x)\right) \operatorname{erfc}\left(\frac{m + \lambda\sigma^2 - x}{\sqrt{2}\sigma^2}\right)$$

$\lambda$  = mean free path of attachment (i.e. attachment length)

$\sigma$  = diffusion




Fit Residuals




# Negative Ion Species Cross Sections


Standard capture and dissociation process for  $\text{SF}_6^-$ :




Standard dissociation processes for  $\text{CF}_4^-$ :



$\text{CS}_2$  likely forms NI in a similar process as  $\text{SF}_6^-$ .




L. G. Christophorou, J. K. Olthoff, Int. J. of Mass Spectrom. 205 (1) 27–41 (2001)




I. Iga, M. V. V. S. Rao, S. K. Srivastava, and J. E. Nogueira, Z. Phys. D 24, 111 (1992).

# Select Averaged Waveforms with Attachment



40 + 5 Torr CF<sub>4</sub> + CS<sub>2</sub>



40 + 10 Torr Ar + CS<sub>2</sub>





# Energy Resolution



# Aside: Attachment in CF<sub>4</sub>

- By measuring the exponential portion of the waveform, we reconstruct measured attachment to within 80%

2024.Jul.03 - 50 torr CF<sub>4</sub> - Nlrns



L. G. Christophorou, J. K. Olthoff, M. V. Rao, *J. Phys. Chem. Ref. Data* 25, 1341–1388 (1996)  
 S. Biagi (1995). <https://magboltz.web.cern.ch/magboltz/>

# Bridging the Gap Between Exponential Fit and EMG

- Exponential fits on  $\text{CF}_4$  waveforms reproduced Christophorou's recommended attachment length to within 8%
- Measurements made in  $\text{Ar}+\text{CS}_2$  found very good agreement between exponential and EMG fits →
- EMG fits were chosen instead of exponential fits in order to measure both attachment and diffusion width
  - Exponential fits could not be verified on waveforms with  $\lambda < 5\text{mm}$ 
    - This region falls roughly in our EMG 'good-fit' region, thus it was straightforward to use EMG results

