

Spatial resolution of IHEP AC-LGADs with different process and structure design

Friday 8 September 2023 12:30 (20 minutes)

AC-coupled Low- Gain Avalanche Detectors (AC-LGAD) are designed as detectors with 100% fill factor for high precision 4D-tracking, which have been studied and researched by many institutes including BNL, FBK et al. Institute of High Energy Physics (IHEP) has also done many researches on AC-LGAD. First IHEP AC-LGAD sensors with a pitch of 2000 μm and AC pad of 1000 μ show time resolution better than 20ps, and spatial resolution better than 16 μm . Testing results show that as the N+ doping dose decreases from 10 P to 0.2 P, the spatial resolution is reduced from 33 μm to 15 μm . Details of the results will be discussed. Second version of IHEP AC-LGAD be fabricated, lower than 0.2P n+ layer dose be used for improve the spatial resolution and sensors with different pad-pitch structures also be fabricated. Testing results show that as decreasing the n+ layer dose from 0.2P to 0.02P, the spatial resolution of AC-LGAD can be lower than 8 μm . Strips with different pad-pitch structures be fabricated and studied, results show that pad-pitch structures will also affect the spatial resolution. Testing results of IHEP AC-LGAD v2 sensors tested by using laser system will also be shown.

Your name

Mei Zhao

Institute

Institute of High Energy Physics of the Chinese Academy of Sciences

Email address

mei.zhao@cern.ch

Presenter: LI, Mengzhao (Chinese Academy of Sciences (CN))

Session Classification: Closing session

Track Classification: Position Sensitive Fast Timing Detectors