

Early Physics Opportunities at DUNE

Sungbin Oh

Hanyang University

KPS 2026 Fall Meeting: Focus: Deep Underground Neutrino Experiment

23 October 2026




한양대학교
HANYANG UNIVERSITY







How Early?



How Early?



How Early?



Challenges in the Neutrino Oscillation Physics Program

Near Detector

$$N_{\text{ND}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}(E_{\nu}) \sigma(E_{\nu}) \epsilon(E_{\nu}) dE_{\nu}$$

Far Detector

$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}(E_{\nu}) P_{\text{osc.}}(E_{\nu}) \sigma(E_{\nu}) \epsilon(E_{\nu}) dE_{\nu}$$

Challenges in the Neutrino Oscillation Physics Program

Near Detector

$$N_{\text{ND}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}(E_{\nu}) \sigma(E_{\nu}) \epsilon(E_{\nu}) dE_{\nu}$$

Far Detector

$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}(E_{\nu}) P_{\text{osc.}}(E_{\nu}) \sigma(E_{\nu}) \epsilon(E_{\nu}) dE_{\nu}$$

Original Flux	Obs.	Cross section	Detector effect
------------------	------	------------------	--------------------

Reality is complicated

Challenges in the Neutrino Oscillation Physics Program

Near Detector

$$N_{\text{ND}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}^{\text{ND}}(E_{\nu}) P_{\text{osc.}}^{\text{ND}}(E_{\nu}) \sigma(E_{\nu}) \epsilon^{\text{ND}}(E_{\nu}) dE_{\nu}$$

Far Detector

$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}^{\text{FD}}(E_{\nu}) P_{\text{osc.}}^{\text{FD}}(E_{\nu}) \sigma(E_{\nu}) \epsilon^{\text{FD}}(E_{\nu}) dE_{\nu}$$

Reality is EVEN more complicated

Challenges in the Neutrino Oscillation Physics Program

Far Detector


$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}^{\text{FD}}(E_{\nu}) P_{\text{osc.}}^{\text{FD}}(E_{\nu}) \sigma(E_{\nu}) \epsilon^{\text{FD}}(E_{\nu}) dE_{\nu}$$

Challenges in the Neutrino Oscillation Physics Program



Far Detector

$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \underbrace{\Phi_{\nu}^{\text{FD}}(E_{\nu})}_{\text{The Movable Near Detector Component (ND-LAr)}} \underbrace{P_{\text{osc.}}^{\text{FD}}(E_{\nu})}_{\text{will constrain uncertainties} \rightarrow \text{But not "Early"}}$$

The Movable Near Detector Component (ND-LAr)
will constrain uncertainties → **But not "Early"**



Instruments 5 (2021) 4, 31



Challenges in the Neutrino Oscillation Physics Program

Far Detector

$$N_{\text{FD}}(E_{\nu}^{\text{reco.}}) = \int \Phi_{\nu}^{\text{FD}}(E_{\nu}) P_{\text{osc.}}^{\text{FD}}(E_{\nu}) \sigma(E_{\nu}) \epsilon^{\text{FD}}(E_{\nu}) dE_{\nu}$$


Final state particles
kinematics: O

$$\sigma(E_{\nu}) \epsilon(E_{\nu}) = \int \frac{d\sigma(E_{\nu})}{dO} \epsilon(O | E_{\nu}) dO$$

We can tackle this **right now!**

Hadrons are Important!


- For charged-current ν_e -argon interactions at the DUNE far detector
- Fraction of hadronic energy deposition is significant: $\sim 30 - 40\%$



Alexander Friedland, Shirley Weishi Li, *Phys.Rev.D* 99 (2019) 3, 036009

Hadrons are Important!


- For charged-current ν_e -argon interactions at the DUNE far detector
- Fraction of hadronic energy deposition is significant: $\sim 30 - 40\%$



Alexander Friedland, Shirley Weishi Li, *Phys.Rev.D* 99 (2019) 3, 036009



Hadrons are Challenging!

- Understanding their **propagation** in liquid argon
 - Measurements of hadron-argon interactions are sparse



Hadrons are Challenging!



- Understanding their **propagation** in liquid argon
 - Measurements of hadron-argon interactions are sparse
- Information lost from inelastically interacting hadrons
 - Energy **reconstruction**
 - Particle identification (**PID**)



Opportunities for Hadrons: ProtoDUNE!



- Hosted by the CERN Neutrino Platform
- DUNE far detector prototypes
- Exposed to momentum-analyzed hadron beams
- Run 1 with the beam (2018): produced many results, with more coming soon
- Run 2 with the beam (2024 - 2025): data is available for further studies!

Phys.Rev.D 110 (2024) 9, 092011
Phys.Rev.D 113 (2026) 5, 052004
Phys.Rev.D 113 (2026) 11, 112002
arXiv:2511.11925, submitted to *PRL*



Opportunities for Hadrons: (1) Interaction with Argon


- Charged pion–argon interaction cross-section measurements using the ProtoDUNE Run 1 data
 - First measurements in this kinetic energy region
 - A first measurement at the Δ -resonance peak would be very interesting
- Proton–argon interaction cross-section measurement: large uncertainties at low kinetic energy
 - Improved proton track end-point reconstruction will be very helpful!




Y. Liu, S.B. Oh et al. DUNE collaboration,
arXiv:2511.11925 (submitted to PRL)

Opportunities for Hadrons: (1) Interaction with Argon


- Exclusive cross-section measurements are also very important
 - Direct impact on neutrino energy resolution and its uncertainty
- Below are results from ProtoDUNE Run 1 data
 - More data from Run 2 is available now for further studies in exclusive channels



(a) Absorption



(b) Charge Exchange




(c) Other Interactions



J. Calcutt et al. DUNE collaboration, *Phys.Rev.D* 113 (2026) 11, 112002

Opportunities for Hadrons: (2) Reconstruction and PID

- Energy reconstruction for inelastically interacting charged pions
 - A novel method was developed using the ProtoDUNE Run 1 data
 - Uses both position and calorimetric information




S.B. Oh et al. DUNE collaboration, *JINST* 20 (2025) 02, P02021





Opportunities for Hadrons: (2) Reconstruction and PID

- Energy reconstruction for inelastically interacting charged pions
 - A novel method was developed using the ProtoDUNE Run 1 data
 - Uses both position and **calorimetric** information





S.B. Oh et al. DUNE collaboration, *JINST* 20 (2025) 02, P02021




Opportunities for Hadrons: (2) Reconstruction and PID

- Energy reconstruction for inelastically interacting charged pions
 - A novel method was developed using the ProtoDUNE Run 1 data
 - Uses both position and **calorimetric** information





Opportunities for Hadrons: (2) Reconstruction and PID

- Energy reconstruction for inelastically interacting charged pions
 - A novel method was developed using the ProtoDUNE Run 1 data
 - Uses both position and **calorimetric** information
 - **Multiple scattering angle** is a promising variable
 - Mass- and energy-dependent; EM only (μ) vs. EM + Strong ($p, \pi^\pm$); excellent angular resolution $\sim \mathcal{O}(\text{mrad.})$





Geant4 Simulation for 14 cm segment in LAr



Opportunities at Near Detector Prototype: The 2x2 Demonstrator




- Demonstration of ND-LAr design with 4 modules at $\sim 60\%$ scale: pixelated charge readout!
- Installed in the Minerva detector hall
- Run 1 (2024): NuMI data (~ 4.5 days, $\sim 30k \nu$) and cosmic data
- Run 2 (2025): calibration data (neutron & γ sources, ^{220}Rn injection ...)
- Run 3 (2026 - 2027): more NuMI and cosmic data



DUNE collaboration, *Instruments* 10 (2026) 1, 18

Opportunities at Near Detector Prototype: The 2x2 Demonstrator


- ν – Ar studies using the NuMI beam
- Testing beyond the standard model scenarios
- Reconstruction & calibration studies





DUNE collaboration, *Instruments* 10 (2026) 1, 18

Opportunities in FD1: (1) Core-Collapse Supernovae

- Core-collapse supernovae produce neutrino bursts
 - Observed only once: SN 1987A, 12 electron events by Kamiokande-II
- The next neutrino burst from a core-collapse supernova will be observed by many neutrino telescopes
 - DUNE far detector modules are also neutrino telescopes





Credit:
NASA, ESA, CSA, M. Matsuura (Cardiff University),
R. Arendt (NASA's Goddard Spaceflight Center & University of Maryland, Baltimore County),
C. Fransson (Stockholm University), J. Larsson (KTH Royal Institute of Technology), A. Pagan (STScI)



Hirata, K et al., *Phys.Rev.Lett.* 58 (1987) 1490-1493


Opportunities in FD1: (1) Core-Collapse Supernovae

- DUNE has a unique sensitivity to ν_e from the neutrino burst: $\nu_e \text{CC}$ ($\nu_e + \text{Ar} \rightarrow e^- + {}^{40}\text{K}^*$)
 - A big advantage for probing the early neutrinos from the burst!





- But there are many challenges (= opportunities)

- Cross-section modeling
 - No measurement yet!
 - SBND: ν_e from μDAR
 - Dedicate event generator (MARLEY)
- Triggering and data streaming systems
- Reconstruction: low-energy electrons and de-excitation photons
- Supernova pointing
 - Improving $\nu_e + e^- \rightarrow \nu_e + e^-$ selection purity is the key



Opportunities in FD1: (2) Atmospheric Neutrinos

- Atmospheric neutrinos offer rich physics
 - $E_\nu < 1$ GeV: upward-going ν_e provides sensitivity to δ_{CP}
 - $E_\nu \approx \mathcal{O}(1$ GeV): sensitivity to mass ordering
 - Main background for beyond the standard model searches
 - e.g., dark matter and baryon number violation






- Neutrino reconstruction study for FD-HD was published in July 2026, *Eur.Phys.J.C* 86 (2026) 7, 904
- Opportunities in FD-HD sensitivity and a full suite of FD-VD (1st module) studies

Opportunities in FD1: (3) Baryon Number Violations

- DUNE has low hadron energy thresholds, mm-scale position resolution, and precise calorimetry
- Could provide world-class sensitivities to baryon number violation searches

- A recent study for $n - \bar{n}$ oscillation using a TransformerCVN architecture achieved 3× the signal efficiency at the same background rejection, relative to the DUNE TDR
- Scaling by $\sqrt{3}$ leads to a $\tau_{n-\bar{n}}$ limit of $\sim 10^9$ s (400 kt·yr)

K. Yu @ Neutrino 2026



Summary

- DUNE construction is progressing well
 - There are physics opportunities before the beam-based science program starts
- **Hadrons** are important for the DUNE long-baseline neutrino oscillation physics program
 - About 30% to 40% of neutrino interaction energy is deposited by hadrons
 - Hadron propagation & interaction modeling, energy reconstruction and particle identification are challenging
 - There are **imminent physics opportunities** with the DUNE FD prototype detectors to tackle them
- There are also **physics opportunities with the first DUNE FD module**
 - To realize DUNE's unique sensitivity to ν_e from core-collapse supernova neutrino bursts
 - Modeling and reconstruction of $\nu_e + \text{Ar} \rightarrow e^- + {}^{40}\text{K}^*$ events
 - Triggering and data streaming systems
 - Selection purity for $\nu_e + e^- \rightarrow \nu_e + e^-$ process is key for pointing performance
 - Atmospheric neutrinos offer rich physics; first far detector module (FD-VD) studies are wide open
 - A recent $n - \bar{n}$ oscillation study shows promising performance; analyzing the first FD data will be exciting

Back Up