




# Near Detectors for Long Baseline Oscillation Experiments

Asher Kaboth  
10 April 2019

# CP Violation

Results shown are for with reactor constraint; T2K only in back up


Posterior probability density



- There is a hint from T2K that  $\delta_{CP}$  is not  $0/\pi$
- Should we trust this result?


# Disappearance Sector

- NOvA results prefer non-maximal mixing
- Very narrow result in  $\Delta m^2_{32}$
- Should we trust this result?



Prefer NH at  $1.8\sigma$   
(T2K similar)

# Oscillation Experiments in a Nutshell



# Why Is This Hard?

QCD RUINS EVERYTHING

Non-perturbative!


Ancient data!

Axial currents!

Effective parameters!

A-scaling is hard!

# Oscillation Experiments in a Nutshell






$$N_n = \Phi_n \times \sigma_n \times \epsilon_n$$

$$N_f = \Phi_f \times \sigma_f \times \epsilon_f \times P(\nu_\alpha \rightarrow \nu_\beta)$$

How many  $\nu_\beta$  here?



# Neutrino Beam

QCD Lives Here



# Hadronic Uncertainties



- Long baseline experiments use associated hadronic production experiments (e.g. NA61/SHINE) to constrain pion/kaon production
- Still need to extrapolate from phase space of associated experiments to full beam line






# Beam Uncertainties


Super-K





ND280: Neutrino Mode,  $\nu_{\mu}$



SK: Antineutrino Mode,  $\bar{\nu}_{\mu}$




# $\nu$ -N Cross Section Model




Uncertainties come from underlying model parameters and normalizations


Charged current quasi-elastic




Charged current multinucleon



Deep Inelastic Scattering




Charged Current 1π





# Mode vs Topology






# T2K Off-Axis Near Detector


Primary Interaction Material: Carbon  
Secondary Interaction Materials:  
Oxygen, Lead, Brass, Argon

Interaction in P0D

Interaction in FGD1




Interaction in ECal




Strategy:

- Parameterize underlying models
- Select data samples to optimize constraints
- Propagate uncertainties through parameters



CC0 $\pi$



CC1 $\pi^+$






CC other



# ND280 $\nu$ -mode samples

FGD1 samples shown;  
FGD2 similar



PRELIMINARY



PRELIMINARY

- Three samples allow sensitivity to different beam energies and cross section interaction modes
- High statistics in neutrino mode provide strong constraints

# CC0 $\pi$ Samples

Before analysis

After analysis



PRELIMINARY



PRELIMINARY

- Clear that data is in better agreement after the analysis
- Adjustment comes through all the modes
- T2K is no longer statistically limited at the near detector!



# Propagation of Uncertainty



SK FHC  $\nu_\mu$  Flux



Error source	1-Ring $\mu$		1-Ring $e$		
	FHC	RHC	FHC	RHC	FHC CC1 $\pi$
Beam	4.3%	4.1%	4.4%	4.2%	4.4%
Cross-section (constr. by ND280)	4.7%	4.0%	4.8%	4.1%	4.1%
Cross-section (all)	5.6%	4.4%	8.4%	6.2%	5.6%
Beam + Cross-section (constr. by ND280)	3.3%	3.3%	3.3%	3.1%	4.0%
Beam + Cross-section (all)	4.4%	2.9%	7.7%	5.7%	5.6%
New $E_b$ fake data parameter	3.2%	1.3%	7.2%	4.1%	2.8%
SK+FSI+SI	3.3%	2.9%	4.1%	4.3%	16.6%
<b>Total</b>	<b>5.5%</b>	<b>4.4%</b>	<b>8.8%</b>	<b>7.3%</b>	<b>17.8%</b>


# 'Fake Data' Analyses

- Generate 'fake data' from alternative models
- Perform full analysis
- Example: Binding energy in nuclei
- Check if the analysis is sensitive to this




# NOvA ND

Primary Interaction Material: Carbon




Scintillator units

Muon catcher



# NOvA ND





## Strategy:

- Unfold ND data to predict true energy spectrum
- Apply Far/Near ratio and oscillations
- Fold back to reconstructed energy
- Systematics are applied as variations on the true-reconstructed matrices




# NOvA ND




- Split into bins of hadronic energy
- This reflects different energy resolutions

# Propagation of Uncertainty


- Statistical uncertainty still dominates for NOvA
- Nevertheless, as datasets increase, this will become increasingly important



# Problems: Phase Space





- Near detectors typically have a restricted phase space relative to their far detectors



- Uncertainties in  $Q^2$  can badly affect this!

# Problems: Energy Spectrum





- Near and far do not see identical fluxes
- Different modes have different energy resolution/biases
- If this is wrong—can produce biases in osc. parameters




# Problem: Model Degeneracy

- Example: can shift energy from protons to neutrons and the ND spectrum looks fine via other model compensations
- Impact on oscillation contours is large




# Next Generation



- Systematic uncertainties have a huge effect on the sensitivity of future LBL experiments
- We have to do much better!



# New Concepts: PRISM



- Use the off-axis effect to generate different energy neutrino beams
- Systematics are low 'bin-to-bin'

# New Concepts: PRISM


- Can construct 'monoenergetic' beams
- Can construct 'pre-oscillated' beams
- Both HK and DUNE have plans for a PRISM detector






# New Concepts: Low Threshold


- Upgrade to T2K ND280 to increase efficiency at high angles
- New detector target with much finer granularity



- Replace (most of) P0D with **Scintillator Detector** + **2 High-Angle TPCs** + **TOF**
- Improve acceptance for large angle tracks
- Keep current “tracker” [2 FGDs + 3 TPCs] (& upstream part of P0D) as well as ECal, magnet & SMRD



# New Concepts: Low Threshold





- Resolution only gets us so far—to do better, need lower density
- Proposal: High pressure gas TPC
- Begin to distinguish low energy hadrons—better mode determination, model constraint





# New Concepts: Low Threshold

- DUNE plans to build a HPgTPC a part of the near detector complex
- Prototype detectors are underway both at FNAL and RHUL




# New Concepts: Beam

- Major problem is the fact that near detectors measure flux times cross section
- Separation of the two is desirable!
- Enter electron scattering!





# New Concepts: Beam



- MINERvA has show a proof-of-concept analysis of this technique
- Difficulty lies in separating events from intrinsic beam nue, tiny cross section
- Future experiments thinking about ways to include this measurement in NDs

# Conclusions

- The difficulty of QCD modeling produces complex challenges for neutrino physics
- The current generation of near detectors do a great job for their experiments
- Novel techniques and analyses are needed to drive the next generation of experiments