

# Neutrino Interferometry for New Physics Search

TK, MPLA27(2012)1230024  
IceCube, Nature Physics 14(2018)961  
IceCube, To be published (2019)

## Motivation

- String theory
- Loop quantum gravity
- Horava-Lifshitz gravity
- Lee-Wick theory
- Non-commutative field theory
- Supersymmetry
- etc

## Physics

- Lorentz violating field
- Quantum foam
- Neutrino-dark matter coupling
- Neutrino-dark energy coupling
- Neutrino-torsion coupling
- Neutrino velocity  $\neq c$
- Violation of equivalent principle
- etc


Check the poster  
“IceCube HESE 7.5-yr data  
by Kareem Farrag

Find us on Facebook,  
“Institute of Physics Astroparticle Physics”  
<https://www.facebook.com/IOPAPP>

Teppei Katori  
Queen Mary University of London  
IoP APP-HEPP meeting, Imperial College London, UK, April 8, 2019



# Neutrino interferometry – Atmospheric neutrinos




Neutrino oscillation is a nature interferometer. Any extra interactions in the Lagrangian contribute the phase shift

For 20 TeV up-going atmospheric neutrinos ( $L \sim 12700\text{km}$ ), detectable phase shift by neutrino is

$$\bar{\psi} a^\mu \gamma_\mu \psi, \quad a \sim 10^{-24} \text{ GeV}$$

If anomalous coupling with neutrinos in vacuum cause a phase shift in similar order, we can see it from **spectrum distortion of atmospheric neutrinos**



Effective Hamiltonian with new physics operators

Teppei Kat 
$$H \sim \frac{m^2}{2E} + \hat{a}^{(3)} - E \cdot \hat{c}^{(4)} + E^2 \cdot \hat{a}^{(5)} - E^3 \cdot \hat{c}^{(6)} \dots$$

# Neutrino interferometry – Atmospheric neutrinos

dim.	method	type	sector	limits	ref.
3	CMB polarization	astrophysical	photon	$\sim 10^{-43}$ GeV	[6]
	He-Xe comagnetometer	tabletop	neutron	$\sim 10^{-34}$ GeV	[10]
	torsion pendulum	tabletop	electron	$\sim 10^{-31}$ GeV	[12]
	muon g-2	accelerator	muon	$\sim 10^{-24}$ GeV	[13]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(3)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(3)})  < 2.9 \times 10^{-24}$ GeV (99% C.L.) $< 2.0 \times 10^{-24}$ GeV (90% C.L.)	this work
4	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-38}$	[7]
	Laser interferometer	LIGO	photon	$\sim 10^{-22}$	[8]
	Sapphire cavity oscillator	tabletop	photon	$\sim 10^{-18}$	[5]
	Ne-Rb-K comagnetometer	tabletop	neutron	$\sim 10^{-29}$	[11]
	trapped $\text{Ca}^+$ ion	tabletop	electron	$\sim 10^{-19}$	[14]
neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(4)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(4)})  < 3.9 \times 10^{-28}$ (99% C.L.) $< 2.7 \times 10^{-28}$ (90% C.L.)	this work	
5	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-34}$ $\text{GeV}^{-1}$	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-22}$ to $10^{-18}$ $\text{GeV}^{-1}$	[9]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(5)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(5)})  < 2.3 \times 10^{-32}$ $\text{GeV}^{-1}$ (99% C.L.) $< 1.5 \times 10^{-32}$ $\text{GeV}^{-1}$ (90% C.L.)	this work
6	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-31}$ $\text{GeV}^{-2}$	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-42}$ to $10^{-35}$ $\text{GeV}^{-2}$	[9]
	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-31}$ $\text{GeV}^{-2}$	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(6)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(6)})  < 1.5 \times 10^{-36}$ $\text{GeV}^{-2}$ (99% C.L.) $< 9.1 \times 10^{-37}$ $\text{GeV}^{-2}$ (90% C.L.)	this work
7	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-28}$ $\text{GeV}^{-3}$	[7]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(7)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(7)})  < 8.3 \times 10^{-41}$ $\text{GeV}^{-3}$ (99% C.L.) $< 3.6 \times 10^{-41}$ $\text{GeV}^{-3}$ (90% C.L.)	this work
8	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-46}$ $\text{GeV}^{-4}$	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(8)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(8)})  < 5.2 \times 10^{-45}$ $\text{GeV}^{-4}$ (99% C.L.) $< 1.4 \times 10^{-45}$ $\text{GeV}^{-4}$ (90% C.L.)	this work

TABLE I: Comparison of attainable best limits of SME coefficients in various fields.

Effective Hamiltonian with new physics operators

Teppei Kat 
$$H \sim \frac{m^2}{2E} + \hat{a}^{(3)} - E \cdot \hat{c}^{(4)} + E^2 \cdot \hat{a}^{(5)} - E^3 \cdot \hat{c}^{(6)} \dots$$



# Neutrino interferometry – Atmospheric neutrinos

dim.	method	type	sector	limits	ref.
3	CMB polarization	astrophysical	photon	$\sim 10^{-43}$ GeV	[6]
	He-Xe comagnetometer	tabletop	neutron	$\sim 10^{-34}$ GeV	[10]
	torsion pendulum	tabletop	electron	$\sim 10^{-31}$ GeV	[12]
	muon g-2	accelerator	muon	$\sim 10^{-24}$ GeV	[13]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(3)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(3)}) $ $< 2.9 \times 10^{-24}$ GeV (99% C.L.) $< 2.0 \times 10^{-24}$ GeV (90% C.L.)	this work
4	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-38}$	[7]
	Laser interferometer	LIGO	photon	$\sim 10^{-22}$	[8]
	Sapphire cavity oscillator	tabletop	photon	$\sim 10^{-18}$	[5]
	Ne-Rb-K comagnetometer	tabletop	neutron	$\sim 10^{-29}$	[11]
	trapped $\text{Ca}^+$ ion	tabletop	electron	$\sim 10^{-19}$	[14]
neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(4)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(4)}) $ $< 3.9 \times 10^{-28}$ (99% C.L.) $< 2.7 \times 10^{-28}$ (90% C.L.)	this work	
5	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-34}$ $\text{GeV}^{-1}$	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-22}$ to $10^{-18}$ $\text{GeV}^{-1}$	[9]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(5)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(5)}) $ $< 2.3 \times 10^{-32}$ $\text{GeV}^{-1}$ (99% C.L.) $< 1.5 \times 10^{-32}$ $\text{GeV}^{-1}$ (90% C.L.)	this work
6	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-31}$ $\text{GeV}^{-2}$	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-42}$ to $10^{-35}$ $\text{GeV}^{-2}$	[9]
	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-31}$ $\text{GeV}^{-2}$	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(6)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(6)}) $ $< 1.5 \times 10^{-36}$ $\text{GeV}^{-2}$ (99% C.L.) $< 9.1 \times 10^{-37}$ $\text{GeV}^{-2}$ (90% C.L.)	this work
7	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-28}$ $\text{GeV}^{-3}$	[7]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(7)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(7)}) $ $< 8.3 \times 10^{-41}$ $\text{GeV}^{-3}$ (99% C.L.) $< 3.6 \times 10^{-41}$ $\text{GeV}^{-3}$ (90% C.L.)	this work
8	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-46}$ $\text{GeV}^{-4}$	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(8)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(8)}) $ $< 5.2 \times 10^{-45}$ $\text{GeV}^{-4}$ (99% C.L.) $< 1.4 \times 10^{-45}$ $\text{GeV}^{-4}$ (90% C.L.)	this work

TABLE I: Comparison of attainable best limits of SME coefficients in various fields.

Atomic experiments have strong limits on lower order couplings (renormalizable). Neutrino oscillations are doing good

Effective Hamiltonian with new physics operators

$$H \sim \frac{m^2}{2E} + \hat{a}^{(3)} - E \cdot \hat{c}^{(4)} + E^2 \cdot \hat{a}^{(5)} - E^3 \cdot \hat{c}^{(6)} \dots$$

# Neutrino interferometry – Atmospheric neutrinos

dim.	method	type	sector	limits	ref.
3	CMB polarization	astrophysical	photon	$\sim 10^{-43}$ GeV	[6]
	He-Xe comagnetometer	tabletop	neutron	$\sim 10^{-34}$ GeV	[10]
	torsion pendulum	tabletop	electron	$\sim 10^{-31}$ GeV	[12]
	muon g-2	accelerator	muon	$\sim 10^{-24}$ GeV	[13]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(3)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(3)})  < 2.9 \times 10^{-24}$ GeV (99% C.L.) $< 2.0 \times 10^{-24}$ GeV (90% C.L.)	this work
4	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-38}$	[7]
	Laser interferometer	LIGO	photon	$\sim 10^{-22}$	[8]
	Sapphire cavity oscillator	tabletop	photon	$\sim 10^{-18}$	[5]
	Ne-Rb-K comagnetometer	tabletop	neutron	$\sim 10^{-29}$	[11]
	trapped $\text{Ca}^+$ ion	tabletop	electron	$\sim 10^{-19}$	[14]
neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(4)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(4)})  < 3.9 \times 10^{-28}$ (99% C.L.) $< 2.7 \times 10^{-28}$ (90% C.L.)	this work	
5	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-34}$ GeV <sup>-1</sup>	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-22}$ to $10^{-18}$ GeV <sup>-1</sup>	[9]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(5)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(5)})  < 2.3 \times 10^{-32}$ GeV <sup>-1</sup> (99% C.L.) $< 1.5 \times 10^{-32}$ GeV <sup>-1</sup> (90% C.L.)	this work
6	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-31}$ GeV <sup>-2</sup>	[7]
	ultra-high-energy cosmic ray	astrophysical	proton	$\sim 10^{-42}$ to $10^{-35}$ GeV <sup>-2</sup>	[9]
	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-31}$ GeV <sup>-2</sup>	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(6)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(6)})  < 1.5 \times 10^{-36}$ GeV <sup>-2</sup> (99% C.L.) $< 9.1 \times 10^{-37}$ GeV <sup>-2</sup> (90% C.L.)	this work
7	GRB vacuum birefringence	astrophysical	photon	$\sim 10^{-28}$ GeV <sup>-3</sup>	[7]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{a}_{\mu\tau}^{(7)}) ,  \text{Im}(\hat{a}_{\mu\tau}^{(7)})  < 8.3 \times 10^{-41}$ GeV <sup>-3</sup> (99% C.L.) $< 3.6 \times 10^{-41}$ GeV <sup>-3</sup> (90% C.L.)	this work
8	gravitational Cherenkov radiation	astrophysical	gravity	$\sim 10^{-46}$ GeV <sup>-4</sup>	[15]
	neutrino oscillation	atmospheric	neutrino	$ \text{Re}(\hat{c}_{\mu\tau}^{(8)}) ,  \text{Im}(\hat{c}_{\mu\tau}^{(8)})  < 5.2 \times 10^{-45}$ GeV <sup>-4</sup> (99% C.L.) $< 1.4 \times 10^{-45}$ GeV <sup>-4</sup> (90% C.L.)	this work

TABLE I: Comparison of attainable best limits of SME coefficients in various fields.


Astrophysical observations have strong limits on higher order couplings non-renormalizable). Neutrino oscillations are better tools there

Effective Hamiltonian with new physics operators

$$H \sim \frac{m^2}{2E} + \hat{a}^{(3)} - E \cdot \hat{c}^{(4)} + E^2 \cdot \hat{a}^{(5)} - E^3 \cdot \hat{c}^{(6)} \dots$$

# Neutrino interferometry - Astrophysical high-energy neutrinos

Combination of longer baseline and higher energy makes extra-terrestrial neutrino to be the most sensitive source of fundamental physics.




# Neutrino interferometry – Astrophysical high-energy neutrinos

Combination of longer baseline and higher energy makes extra-terrestrial neutrino to be the most sensitive source of fundamental physics.

Vacuum Cherenkov radiation and Blazar neutrino ToF can limit new physics of neutrino.

Neutrino mixing properties of astrophysical neutrinos can push this limit further.

- In principle, we can do it without knowing neutrino flavours at the production
- This is the most sensitive, as long as we assume new physics cause neutrino mixing



# Neutrino interferometry - Astrophysical high-energy neutrinos

Any new physics can end up in the effective Hamiltonian

$$h_{\text{eff}} = \frac{1}{2E} U^\dagger M^2 U + \sum_n \left( \frac{E}{\Lambda_n} \right)^n \tilde{U}_n^\dagger O_n \tilde{U}_n = V^\dagger \Delta V$$

neutrino oscillation formula

$$P_{\alpha \rightarrow \beta}(L) = 1 - 4 \sum_{i>j} \text{Re}(V_{\alpha i}^* V_{\beta i}^* V_{\alpha j} V_{\beta j}) \sin^2 \left( \frac{\Delta_{ij}}{2} L \right) + 2 \sum_{i>j} \text{Im}(V_{\alpha i}^* V_{\beta i}^* V_{\alpha j} V_{\beta j}) \sin(\Delta_{ij} L)$$

neutrino mixing formula

$$P_{\alpha \rightarrow \beta}(L \rightarrow \infty) \sim 1 - 2 \sum_{i>j} \text{Re}(V_{\alpha i}^* V_{\beta i}^* V_{\alpha j} V_{\beta j}) = \sum_i |V_{\alpha i}|^2 |V_{\beta i}|^2$$

Information of small contamination of new physics appears on **neutrino flavour mixings**

At high energy, neutrino mass term is suppressed




# Neutrino interferometry - Astrophysical high-energy neutrinos

There are 3 astrophysical neutrino production models

- i. pion decay dominant model, 1:2:0
- ii. electron neutrino dominant model, 1:0:0
- iii. muon neutrino dominant model, 0:1:0
- iv. tau neutrino dominant model, 0:0:1

Initial flavour ratio is modified on the Earth due to neutrino mixing



# Neutrino interferometry - Astrophysical high-energy neutrinos


There are 3 astrophysical neutrino production models

- i. pion decay dominant model, 1:2:0
- ii. electron neutrino dominant model, 1:0:0
- iii. muon neutrino dominant model, 0:1:0
- iv. tau neutrino dominant model, 0:0:1

Initial flavour ratio is modified on the Earth due to neutrino mixing

All possible flavour ratio is confined in a small space.

If you measure flavour ratio outside of this, you win!



# Neutrino interferometry - Astrophysical high-energy neutrinos

There are 3 astrophysical neutrino production models

- i. pion decay dominant model, 1:2:0
- ii. electron neutrino dominant model, 1:0:0
- iii. muon neutrino dominant model, 0:1:0
- iv. tau neutrino dominant model, 0:0:1


Initial flavour ratio is modified on the Earth due to neutrino mixing

All possible flavour ratio is confined in a small space.

If you measure flavour ratio outside of this, you win!




E.g.) dim-3 new physics operator  
 $\sim 10^{-24}$  GeV (just below IceCube atmospheric neutrino limit)




# New IceCube data (2018)

## Blazar neutrino


- IC170922A and TXS 0506+056
- Optical coincidence
- Clustering from this direction



IceCube, Science361(2018)147



WORK IN PROGRESS



[https://charge.wisc.edu/icecube/wipac\\_store.aspx](https://charge.wisc.edu/icecube/wipac_store.aspx)



**IceCube IC170922 t-shirt (Crew-Neck)**  
\$18.00  
The front side features an image of "IC170922" and the IceCube logo on the back.  
Heathered navy, crewneck, rinspun cotton/polyester, Available in unisex sizes S-2XL. Runs small.

Support IceCube!



## New flavour ratio measurement

- Likelihood is very shallow and fit confuse between  $\nu_e$  and  $\nu_\tau$
- New flavour ratio result has some power to distinguish between  $\nu_e$  and  $\nu_\tau$

# IceCube flavor ratio test for new physics (to be published)

We start to exclude possible new physics in Planck scale signal region

- This moment, we can exclude only 2 scenarios
- dimension-3 vacuum operator limit  $(\bar{\psi}a^\mu\gamma_\mu\psi, \bar{\psi}b^\mu\gamma_\mu\gamma_5\psi) \sim 10^{-25} \text{ GeV}$
- dimension-4 vacuum operator limit  $(\bar{\psi}c^{\mu\nu}\gamma_\mu\partial_\nu\psi, \bar{\psi}d^{\mu\nu}\gamma_\mu\gamma_5\partial_\nu\psi) \sim 10^{-33}$
- dimension-6 vacuum operator limit  $\sim 10^{-40} \text{ GeV}^2$





# Conclusion

Neutrino interferometry is a powerful technique to look for new physics if new physics couple with neutrinos and they cause neutrino mixings.

Spectrum distortion of atmospheric neutrino is used to look for new physics.

Astrophysical neutrino mixing sensitivity reaches to naïve expectation of Planck scale. However, in this moment, the sensitivity is limited. We need more statistics and better particle identification algorithm to find new physics.

**Thank you for your attention!**

