



Contribution ID: 91

Type: **Oral (Non-Student) / orale (non-étudiant)**

## **Weak measurements with an ensemble quantum processor.**

*Thursday 19 June 2014 10:00 (15 minutes)*

NMR ensemble quantum processors can be used for implementing a universal quantum circuit with a limited number of qubits using current technologies. The computation is carried out with high fidelity over a large number of processors (individual molecules) in parallel and the readout corresponds to ensemble averages of single-qubit observables. This type of readout poses a obstacle when the algorithm requires post-processing on individual outcomes, for example in the case of weak measurements with post-selection. We present a method for overcoming the challenge of post-selection in an ensemble processor and show experimental results of the first weak measurement in NMR. The experiment involves 3 qubits but can be extended to larger systems of up to 12 qubits using conventional methods. These extensions may be useful for carrying out weak measurements experiments that cannot be done with the more common optical setup.

- [arXiv:1311.5890](https://arxiv.org/abs/1311.5890)

**Author:** BRODUTCH, Aharon (Univesity of Waterloo)

**Co-authors:** Dr LU, Dawei (University of Waterloo); Mr LI, Hang (Tsinghua University); Mr LI, Jun (University of science and technology China); Prof. LAFLAMME, Raymond (University of Waterloo)

**Presenter:** BRODUTCH, Aharon (Univesity of Waterloo)

**Session Classification:** (R1-6) Quantum Information Theory - DTP-DAMOPC-DCMMP / Théorie de l'information quantique - DPT-DPAMPC-DPMCM

**Track Classification:** Division of Atomic, Molecular and Optical Physics, Canada / Division de la physique atomique, moléculaire et photonique, Canada (DAMOPC-DPAMPC)