



Contribution ID: 146
compétition)

Type: **Oral (Student, In Competition) / Orale (Étudiant(e), inscrit à la**

Dissociative Excitation of Thymine by Electron Impact

Monday 16 June 2014 15:00 (15 minutes)

A crossed electron-gas beam system coupled to a VUV spectrometer has been used to investigate the dissociation of thymine ($C_5H_6N_2O_2$) into excited atomic fragments in the electron-impact energy range from threshold to 375 eV. A special stainless steel oven is used to vaporize the thymine and form it into a beam where it is intersected by a magnetically collimated electron beam, typical current 50 μA . The main features in the spectrum are the H Lyman series lines. The probability of extracting excited C or N atoms from the ring is shown to be very small. In addition to spectral data, excitation probability curves as a function of electron energy will be presented for the main emission features. Possible dissociation channels and excitation mechanisms in the parent molecule will be discussed.

The authors thank NSERC (Canada) for financial support.

Author: TIESSEN, Collin (U)

Co-authors: Dr MCCONKEY, J W (University of Windsor); Dr HEIN, Jeff (Jet Propulsion Laboratory, California Institute of Technology); Mr TROCCHI, Joshua (University of Windsor); Dr KEDZIERSKI, W (University of Windsor)

Presenter: TIESSEN, Collin (U)

Session Classification: (M1-7) Ultrafast Imaging and Spectroscopy I - DAMOPC / Imagerie ultrarapide et spectroscopie I - DPAMPC

Track Classification: Division of Atomic, Molecular and Optical Physics, Canada / Division de la physique atomique, moléculaire et photonique, Canada (DAMOPC-DPAMPC)