

Contribution ID: 243

Type: Oral (Non-Student) / orale (non-étudiant)

## Track-based alignment of the ATLAS Inner Detector: New extensions and expected performance for the next physics data run

Wednesday 18 June 2014 14:30 (15 minutes)

The Large Hadron Collider (LHC) at CERN is the world's largest particle accelerator, colliding proton beams at

unprecedented centre-of-mass energies. ATLAS is a multipurpose experiment that records the products of LHC

collisions. In order to reconstruct the trajectories of particles produced in these collisions, ATLAS is equipped with a precision tracking system, the Inner Detector (ID). The ID alignment procedure ensures an accurate determination of the position and orientation of the detector's components, such that detector mis-alignments do not degrade the physics performance. During the current shutdown, the ATLAS experiment will upgrade its ID with the installation of a new, additional pixel layer, named Insertable B-Layer (IBL). It will be installed between the existing pixel detector and a new smaller diameter beam-pipe. The IBL will ensure excellent tracking, vertexing and b-tagging performance during the LHC run II phase, but also imposes new challenges to the ID alignment. We report on the track-based alignment approaches, their new extensions, and the expected performance for the next physics data run.

Author: DANNINGER, Matthias (University of British Columbia (CA))

Presenter: DANNINGER, Matthias (University of British Columbia (CA))

**Session Classification:** (W2-8) Energy Frontier: ATLAS at ~14 TeV - PPD-DIMP / Frontière d'énergie: ATLAS à ~14 TeV - PPD-DPIM

Track Classification: Particle Physics / Physique des particules (PPD)