



Contribution ID: 185
compétition)

Type: **Oral (Student, In Competition) / Orale (Étudiant(e), inscrit à la**

A search at Super-Kamiokande for low mass dark matter candidates in the T2K neutrino beam

Friday 20 June 2014 12:30 (15 minutes)

The T2K neutrino beam is produced by colliding 30 GeV protons with a graphite target, and some dark sector models predict that a dark matter candidate could be created in the collision. This massive and neutral particle could scatter off a nucleon in Super-Kamiokande, a 50 kilotonne water Cherenkov detector. Similar to the neutral current quasi-elastic neutrino-oxygen interaction, the dark matter candidate could interact with the oxygen nucleus. As the nucleus de-excites, 6 MeV gamma-rays are emitted which can be efficiently detected by Super-Kamiokande. The longer time of flight for a dark matter candidate, compared to a neutrino, allows separation between the dark matter induced signal and the neutrino induced background. In the intense global effort to measure dark matter, this complementary search investigates the sub-GeV mass range where other experiments have reduced sensitivity.

Author: NANTAIS, Corina (U)

Presenter: NANTAIS, Corina (U)

Session Classification: (F-PLN1) - CAP Best Student Presentations Final Competition / Session plénière - Compétition finale de l'ACP pour les meilleures communications étudiantes

Track Classification: Particle Physics / Physique des particules (PPD)