

Contribution ID: 794 compétition)

Type: Oral (Student, In Competition) / Orale (Étudiant(e), inscrit à la

## Double-beta decay half-life of 96Zr –nuclear physics meets geochemistry

Monday 15 June 2015 15:00 (15 minutes)

Double-beta (\beta\beta) decay measurements are a class of nuclear studies with the objective of detecting the neutrinoless (0\nu) decay variants. Detection of a 0\nu\beta\beta decay would prove the neutrino to be massive and to be its own anti-particle (i.e., a Majorana particle). A key parameter in the detection of the 0\nu\beta\beta decay is the energy, or Q-value, of the decay.

 $\{96\}$ Zr is of particular interest as a double-beta decay candidate. A geochemical measurement of its \beta\beta decay half-life by measuring an isotopic anomaly of the  $\{96\}$ Mo daughter in ancient zircon samples yielded a value of  $0.94(32)x10^{19}$  yr [1]. More recently, the NEMO collaboration measured the half-life directly to be  $2.4(3)x10^{19}$  yr [2], twice as long as the geochemical measurement. As the geochemical result could be contaminated by a sequence of two single \beta-decays, the first being a 4-fold unique forbidden \beta-decay of  $\{96\}$ Zr to the 44 keV J $\{pi\}$ =5<sup>+</sup> excited state in  $\{96\}$ Nb, followed by the 23 h \beta-decay of  $\{96\}$ Nb to  $\{96\}$ Mo, further study is mandated. Depending on the Q-value for the first decay, the estimated half-life could be of the same order as the one for the \beta\beta-decay [3]. However, the key parameter is the Q-value for the single \beta-decay, which enters in leading order as Q $\{13\}$  into the phase-space factor of the decay.

Such a study is being carried out at the TRIUMF TITAN experiment and at the University of Calgary Isotope Science Lab. At TITAN we are measuring the Q-values for the ^{96}Zr to ^{96}Mo \beta\beta-decay and for the ^{96}Zr to ^{96}Nb single \beta-decay, with the goal of reaching a precision near 0.1 keV. At the UCalgary ISL, we are repeating the measurement of the ^{96}Mo isotopic anomaly using modern equipment and techniques. Combined, these measurements will remove a long-standing discrepancy of the two independent ^{96}Zr \beta\beta-decay half-life measurements.

[1] M. E. Wieser and J. R. De Laeter, Phys. Rev. C 64, 024308 (2001).

[2] NEMO-3 Collaboration, Nucl. Phys. A 847, 168-179 (2010).

[3] J. Suhonen, Univ. Jyväskylä, private communication.

## Author: MAYER, Adam (University of Calgary)

**Co-authors:** FREKERS, Dieter (Institut fuer Kernphysik); DILLING, Jens (triumf/UBC); WIESER, Michael (University of Calgary); THOMPSON, Robert (University of Calgary)

Presenter: MAYER, Adam (University of Calgary)

**Session Classification:** M1-6 Neutrinoless Double-beta Decay I (PPD-DNP) / Double désintégration beta sans neutrino I (PPD-DPN)

Track Classification: Nuclear Physics / Physique nucléaire (DNP-DPN)