



Contribution ID: 762
compétition)

Type: **Oral (Student, In Competition) / Orale (Étudiant(e), inscrit à la**

Dense Plasma Focus for Short-Lived Isotope Activation

Tuesday 16 June 2015 17:00 (15 minutes)

Short-lived radioisotopes (SLRs) are used for medical applications including positron emission tomography (PET). The required activity for N-13 for PET is about 4 GBq for a myocardial blood perfusion assessment. Dense plasma focus (DPF) has been considered as a low cost method for producing SLRs as an alternative to conventional cyclotron facilities. A low energy dense plasma focus has been built and optimized at the University of Saskatchewan to study the feasibility of SLRs production, in particular N-13 using energetic deuteron ion beams produced in a dense plasma focus. X-ray detectors and a Faraday cup have been used to characterize the DPF properties, particularly the ion beam energy based on time-of-flight measurements. The preliminary results have shown generation of ions with energies up to 2 MeV, well exceeding the threshold energy for N-13 production (328 keV). Electrical signals have been used for circuit analyses in order to interpret the anomalous plasma resistance and plasma inductance during the pinch phase. Simulation of N-13 activation using deuteron beam has been carried out.

Author: Mr BEHBAHANI, R. A. (University of Saskatchewan)

Co-authors: Dr HIROSE, Akira (University of Saskatchewan); XIAO, Chijin (Univ. of Saskatchewan); Dr WOLFE, Sean (Plasmionique Inc.); Mr CHUNG, seanne (University of Saskatchewan)

Presenter: Mr BEHBAHANI, R. A. (University of Saskatchewan)

Session Classification: T3-10 Special session to honour Dr. Akira Hirose II (DPP) / Session spéciale en l'honneur du Dr Akira Hirose II (DPP)

Track Classification: Plasma Physics / Physique des plasmas (DPP)