



Contribution ID: 674

Type: Oral (Non-Student) / orale (non-étudiant)

Inverse melting in a simple 2D liquid

Tuesday 16 June 2015 14:45 (15 minutes)

We employ several computer simulation techniques to study the phase behaviour of a simple, two dimensional monodisperse system of particles interacting through a core-softened potential comprising a repulsive shoulder and an attractive square well. This model was previously constructed and used to explore anomalous liquid behaviour in 2D and 3D, including liquid-liquid phase separation [1]. The calculated phase diagram includes six crystal phases in addition to the liquid and gas. Interestingly, we find that one of the melting curves exhibits inverse melting, for which the liquid freezes to a crystal upon isobaric heating over a very small range of pressure [2]. We find that the range of inverse melting can be enlarged by increasing the extent of the repulsive shoulder, and show that despite occurring in 2D, the melting transition is first order and to a liquid, rather than to a hexatic or quasicrystal phase [3]. As this range increases, the topology of the phase diagram changes systematically until it breaks, leading to even more crystal phases appearing.

[1] A. Scala, M. R. Sadr-Lahijany, N. Giovambattista, S. V. Buldyrev, and H. E. Stanley, Phys. Rev. E 63, 041202 (2001).

[2] A. M. Almudallal, S. V. Buldyrev, and I. Saika-Voivod, J. Chem. Phys. 137, 034507 (2012).

[3] A. M. Almudallal, S. V. Buldyrev, and I. Saika-Voivod, J. Chem. Phys. 140, 144505 (2014).

Author: ALMUDALLAL, Ahmad (Memorial University of Newfoundland)

Co-authors: Prof. SAIKA-VOIVOD, Ivan (Memorial University of Newfoundland); Prof. BULDYREV, Sergey (Yeshiva University)

Presenter: ALMUDALLAL, Ahmad (Memorial University of Newfoundland)

Session Classification: T2-2 Condensed Matter Theory (DCMMP-DTP) / Théorie de la matière condensée (DPMCM-DPT)

Track Classification: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)