



Contribution ID: 1314

Type: Oral (Non-Student) / orale (non-étudiant)

Shape dependence of two-cylinder Renyi entropies for a free boson lattice field theory

Thursday 16 June 2016 14:30 (15 minutes)

We study the bipartite von Neumann and $\alpha = 2$ Renyi entanglement entropies for a system of free bosons put on a torus cut into two cylinders. The torus is formed out of the $L \times L$ square lattice, and the entropy is supposed to scale as $S = aL + c\gamma(L_A/L) + \dots$, with L_A being the size of the partitioned region. c and a are some constants and the function $\gamma(L_A/L)$ that is not known analytically is supposed to reflect some universality. We compute $\gamma(L_A/L)$ numerically and compare the results to several candidate functions derived from Quantum Lifshitz model, anti de-Sitter gravity in $3 + 1$ dimensions, and an Extensive Mutual Information model. Using lattices of different size, we explore the finite-size-scaling behaviour of the residuals for each fit, to attempt to discern which function most effectively describes the thermodynamic limit of the free boson system.

Authors: COOK, Caleb (Perimeter Institute for Theoretical Physics); DALIDOVICH, Denis (Perimeter Institute for Theoretical Physics); LANTAGNE-HURTUBISE, Etienne (Perimeter Institute for Theoretical Physics); HAYWARD-SIERENS, Lauren (Perimeter Institute for Theoretical Physics); CHOJNACKI, Leilee (Perimeter Institute for Theoretical Physics); MELKO, Roger (University of Waterloo); VLAAR, Tiffany (Perimeter Institute for Theoretical Physics)

Presenter: DALIDOVICH, Denis (Perimeter Institute for Theoretical Physics)

Session Classification: R2-5 Quantum Information and Quantum Optics (DCMMP-DAMOPC) / Information quantique et optique quantique (DPMCM-DPAMPC)

Track Classification: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)