

Contribution ID: 2162

of Physicists

Canadian Association

Association canadienne des physiciens et physiciennes

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Enhanced α -Transfer population of the 2_{ms}^+ mixed-symmetry state in 52 Ti

Wednesday 13 June 2018 14:30 (15 minutes)

The residual nucleon-nucleon interaction plays a crucial role in nuclear structure physics. In spherical eveneven nuclei the quadrupole interaction leads to characteristic low-lying 2+ states of proton-neutron mixed symmetry character, decaying via M1 transitions to the proton-neutron symmetric states. We have calculated the associated M1 transition strengths in the 52 Ti nucleus within the ab initio valence-space in-medium similarity renormalization group based on NN and 3N forces from chiral effective field theory. In this framework we also construct an effective valence-space M1 operator, which captures nonperturbatively many-body excitations outside the valence space. Our calculations well describe the established mixed-symmetry state in 52 Ti nucleus. Thus, ab-initio calculations are able to describe fundamental low-lying collective excitations in nuclei.\\

In order to investigate the microscopic structure of the mixed symmetry state in ⁵²Ti nucleus even further, we run an experiment on the ⁵²Ti nucleus, populated via the alpha transfer reaction ⁴⁸Ca(¹²C,⁸Be)⁵²Ti using a ⁴⁸Ca beam from the Maier-Leibnitz-Laboratory in Munich. Gamma rays of populated states were detected with the high-granularity MINIBALL array of HPGe detectors, and charged particles were detected using a highly segmented DSSD silicon detector, allowing to select the channel of interest via a multiplicity coincidence condition. In first order, ⁵²Ti can be described as the coupling of a ⁴He nucleus to the doubly-magic ⁴⁸Ca core. In the frame work of the interacting boson model 2 (IBM-2), Alonso *et al.*[1] have shown that the population of the MS 2_{ms}^+ state is strictly forbidden. This prediction is also confirmed in our new shell model calculation using the full fp space and effective interactions. Alpha transfer spectroscopic factors were evaluated with a cluster model using the oxbash shell model code. In contrast to the theoretical predictions, we experimentally find an exceptionally strong population of the well-established 2_{ms}^+ mixed-symmetry state in ⁵²Ti relative to the population of the 2_1^+ state. We discuss the impact of our findings to our microscopic understanding of the proton-neutron interaction and shell structure in this interesting region of the nuclear chart.

References

[1] C.E. Alonso and J.M. Arias, Phys. Rev. C 78, 017301 (2008).

Author: ALI, Fuad A. (Univerity of Guelph)

Co-authors: MUECHER, Dennis (University of Guelph); BILDSTEIN, V. (Department of Physics, University of Guelph, Guelph, Ontario); GARRETT, Paul (University of Guelph); SVENSSON, Carl (University of Guelph); BROWN, B. Alex (Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan, USA.); HOLT, J. D. (TRIUMF 4004 Wesbrook Mall, Vancouver, British Columbia V6T 2A3, Canada); KILIC, A. I (University of Guelph)

Presenter: ALI, Fuad A. (Univerity of Guelph)

Session Classification: W3-8 Nuclear Structure II (DNP) | Structure nucléaire II (DPN)

Track Classification: Nuclear Physics / Physique nucléaire (DNP-DPN)