



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 210

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Ab-initio calculations of structure factors for dark matter searches

Wednesday, June 9, 2021 5:10 PM (10 minutes)

We present the first ab-initio calculations of the structure factors for elastic spin-dependent WIMP scattering off ^{19}F , ^{27}Al , ^{23}Na , ^{27}Al , ^{29}Si , ^{73}Ge , ^{127}I and $^{129,131}\text{Xe}$. A set of established two- (NN) and three-nucleon (3N) interactions derived from chiral effective field theory (EFT) are used for nuclear interaction, including N^3LO -level NN + N^2LO -level 3N, N^4LO -level NN + N^2LO -level 3N, and N^2LO -level NN+3N with the (1232)-isobar degrees of freedom. Within the same chiral EFT framework, we employ corresponding WIMP-nucleus currents at the one-body level and also include the effects from axial-vector two-body currents. We then apply the ab-initio in-medium similarity renormalization group to construct valence-space Hamiltonians and consistently transformed operators of nuclear responses. By combining the newly developed frameworks, natural orbitals and expressing the 3N force with a very large basis size, we obtain basis-space converged structure factors in heavy nuclei. This work paves the path toward a true first-principles calculation of the structure factor for WIMP scattering in all nuclei relevant for ongoing searches. All results are publicly available in a Jupyter notebook.

Author: Dr HU, Baishan (TRIUMF)

Co-authors: Prof. HOLT, Jason (TRIUMF and McGill University); Dr MIYAGI, Takayuki (TRIUMF); Mr PADUA, José (TRIUMF); Dr STROBERG, Ragnar (University of Washington)

Presenter: Dr HU, Baishan (TRIUMF)

Session Classification: W4-5 Theory I (DNP) / Théorie I (DPN)

Track Classification: Nuclear Physics / Physique nucléaire (DNP-DPN)