



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 335

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Is New Physics Needed to Explain the ATOMKI Anomaly?

Thursday 10 June 2021 12:55 (10 minutes)

Do we really need a hypothetical gauge boson, “X17”, to explain the famous ATOMKI measurements? Or can there be some interplay between the theoretical and experimental effects? We show that the bump in the ${}^8\text{Be}(18.15) \rightarrow {}^8\text{Be} + e^+ + e^-$ decay data can be reproduced within the Standard Model by adding the full set of second-order corrections and the interference terms to the Born-level decay amplitudes, and demonstrate how experimental selection and acceptance bias exacerbate the apparent difference between the experimental data and the Born-level prediction.

Authors: Prof. ALEKSEJEVS, Aleksandrs (Memorial University of Newfoundland); BARKANOVA, Svetlana (Grenfell Campus of Memorial University); KOLOMENSKY, Yury G (University of California, Berkeley); SHEFF, Benjamin (University of Michigan)

Presenter: Prof. ALEKSEJEVS, Aleksandrs (Memorial University of Newfoundland)

Session Classification: R2-5 Theory III (DNP) / Théorie III (DPN)

Track Classification: Nuclear Physics / Physique nucléaire (DNP-DPN)