



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 461

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Mode-coupling instability of two-dimensional complex plasma crystals in asymmetric capacitively-coupled radio-frequency discharges

Tuesday 8 June 2021 15:35 (15 minutes)

The dependence of the mode-coupling instability threshold in two-dimensional complex plasma crystals is studied. It is shown that for a given microparticle suspension at a given discharge power there exist two thresholds in pressure. Above a specific pressure p_{\max} , the monolayer is always in the crystal phase. Below a specific pressure p_{\min} , the crystalline monolayer undergoes the mode-coupling instability and the monolayer is in the fluid phase. In between p_{\min} and p_{\max} , the crystal will be in the fluid phase when increasing the pressure from below p_{\min} until it reaches p_{\max} where it recrystallises, while it remains in the crystal phase when decreasing the pressure from above p_{\max} until it reaches p_{\min} . A simple auto-consistent sheath model can explain the melting threshold as a function of pressure and rf power due the changes of the sheath electric field and the microparticle charges leading to the crossing of the compressional in-plane phonon mode and the out-of plane phonon mode.

Author: COUEDEL, Lenaic (University of Saskatchewan)

Co-authors: Dr NOSENKO, Vladimir (Institut für Materialphysik im Weltraum, Deutsches Zentrum für Luft- und Raumfahrt (DLR)); Dr ZHDANOV, Sergey (Institut für Materialphysik im Weltraum, Deutsches Zentrum für Luft- und Raumfahrt (DLR))

Presenter: COUEDEL, Lenaic (University of Saskatchewan)

Session Classification: TS-3 Plasma Physics Symposium (DPP) / Symposium de physique des plasmas (DPP)

Track Classification: Symposia Day (DPP) - Low temperature plasmas/Fusion plasmas (magnetic and inertial confinement)/ Laser plasmas/Basic plasmas