



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 264 Type: **Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)**

## **(G\*) Harvesting Entanglement inside a Black Hole**

*Thursday 10 June 2021 17:03 (3 minutes)*

We carry out the first investigation of the entanglement and mutual information harvesting protocols for detectors moving on freely falling trajectories that cross the horizon of a black hole. We consider two pointlike Unruh-DeWitt detectors in different combinations of free-falling and static trajectories in (1+1)-dimensional Schwarzschild black hole spacetime and compare the results. We find that (i) correlations harvested between free-falling and static detectors are always less than those of the familiar two-static-detector case (ii) there is a large kinematic component to the correlations; (iii) correlations can be harvested purely from the quantum vacuum, with evidence that this is possible even when the detectors are causally disconnected by the event horizon; (iv) the previously known 'entanglement shadow' near the horizon is indeed absent for the two free-falling-detector case since the relative gravitational redshift of the detectors remains finite as the horizon is crossed, in accordance with the equivalence principle.

**Author:** GALLOCK-YOSHIMURA, Kensuke (University of Waterloo)

**Co-authors:** TJOA, Erickson (University of Waterloo); MANN, Robert (University of Waterloo)

**Presenter:** GALLOCK-YOSHIMURA, Kensuke (University of Waterloo)

**Session Classification:** R4-2 Quantum Information: Theory (DTP) / Information quantique: théorie (DPT)

**Track Classification:** Theoretical Physics / Physique théorique (DTP-DPT)