



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 44

Type: **Invited Speaker** / **Conférencier(ère) invité(e)**

(I) Magneto-optical Kerr effect and signature of the chiral anomaly in a Weyl semimetal in magnetic field

Wednesday 9 June 2021 16:45 (5 minutes)

One striking property of the Landau level spectrum of a Weyl semimetal (WSM) is the existence of a chiral Landau level in which the electrons propagate unidirectionally along the magnetic field. This linearly dispersive level profoundly influences the optical properties of the WSM especially if it originates from a tilted Weyl cone. In this talk, we compare the behavior of the magneto-optical Kerr effect (MOKE) in a WSM with that of a normal (i.e. non-topological) metal. We obtain the Kerr angle from the optical conductivity tensor $\sigma_{\alpha\beta}(\omega)$ using the minimal model of a WSM developed in Ref. [1] which has four tilted Weyl nodes related by mirror and time-reversal symmetry. In the Voigt configuration, a large peak of the Kerr angle occurs at the plasmon frequency. We show that the blueshift in frequency of this peak with increasing magnetic field is a signature of the chiral anomaly in the MOKE.

[1] S. Bertrand, Jean-Michel Parent, R. Côté, and I. Garate, Phys. Rev. B 100, 075107 (2019).

Author: Prof. CÔTÉ, René (Université de Sherbrooke)

Co-authors: Mr PARENT, Jean-Michel (Université de Sherbrooke); Prof. GARATE, Ion (Université de Sherbrooke)

Presenter: Prof. CÔTÉ, René (Université de Sherbrooke)

Session Classification: W4-2 Theory and Condensed Matter (DTP/DCCMP) / Théorie et matière condensée (DPT/DPMCM)

Track Classification: Theoretical Physics / Physique théorique (DTP-DPT)