


# Measurement of the very rare $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay at CERN SPS

Bob Velghe<sup>\*</sup>, for the NA62 Collaboration

2021 CAP Virtual Congress  
June 9, 2021

# $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ - In the Standard Model

Can be represented by penguin and box diagrams:



It is a Flavor Changing Neutral Current: the decay is extremely suppressed.

Theoretical uncertainties well under control: QCD and electroweak corrections, hadronic matrix element related to the  $K^+ \rightarrow \pi^0 e^+ \nu_e$  decay.

[F. Mescia, C. Smith, 07'] [J. Brod, M. Gorbahn, 08'] [J. Brod, M. Gorbahn, E. Stamou, 11']

In terms of the CKM parameters:


$$\begin{aligned} \mathcal{B}_{\text{SM}}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) &= (8.39 \pm 0.30) \times 10^{-11} \left[ \frac{|V_{cb}|}{40.7 \times 10^{-3}} \right]^{2.8} \left[ \frac{\gamma}{73.2^\circ} \right]^{0.74} \\ &= (8.4 \pm 1.0) \times 10^{-11} \end{aligned}$$

CKM uncertainties:  $|V_{cb}| \approx 9.9\%$ ,  $\gamma \approx 6.7\%$ . [A. J. Buras et al, 15']


# $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ – Beyond the Standard Model

$K^+ \rightarrow \pi^+ \nu \bar{\nu}$  has been studied in many BSM scenarios. A few examples:

- **$Z'$  models**, [A. J. Buras, F. De Fazio, J. Girrbach, 13'] [A. J. Buras, D. Buttazzo, R. Kneijens, 15'] [J. Aebischer, A. J. Buras, J. Kumar, 20']
- **Leptoquark models**, [C. Bobeth, A. J. Buras, 18'] [S. Fajfer, N. Košnik, L. Vale Silva, 18']
- **Supersymmetry**, [G. Isidori et al, 06'] [T. Blažek, P. Maták, 14'] [M. Tanimoto, K. Yamamoto, 16']
- **Lepton Flavour Violation models**. [M. Bordone et al, 17']




[A. J. Buras et al, 15']



[M. Bordone et al, 17']

# $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ - Experimental Program



$$\mathcal{B}_{\text{Exp.}}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \left(1.73^{+1.15}_{-1.05}\right) \times 10^{-10} \text{ (E787/E949)}$$

Decay at rest [E787/949 Collaboration, 08']

$$\mathcal{B}_{\text{Exp.}}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \left(0.48^{+0.72}_{-0.48}\right) \times 10^{-10} \text{ (NA62 2016 + 2017)}$$




Decay in flight [NA62 Collaboration, 20']

This talk: NA62 2018 results [NA62 Collaboration, 21']

# $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ - Decay in Flight

Signal:  $K^+$  associated to a  $\pi^+$  and missing energy.


- Identification of  $K$  and  $\pi$ ,
- Multi-track event rejection,
- Vetoes for  $\gamma$  and  $\mu$ , rejection  $> \mathcal{O}(10^7)$ ,
- $\mathcal{O}(100 \text{ ps})$  timing for  $K - \pi$  matching,
- Excellent kinematic reconstruction  $\rightarrow m_{\text{miss.}}^2 = (P_{K^+} - P_{\pi^+})^2$ .



The main kaon backgrounds are  $K^+ \rightarrow \mu^+ \nu_\mu (\gamma)$ ,  $K^+ \rightarrow \pi^+ \pi^0 (\gamma)$ ,  $K^+ \rightarrow \pi^+ \pi^+ \pi^-$  and  $K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$ .

# NA62 Experiment at the CERN SPS – In 2018

Beam: 75 GeV/c  $\pm$  1%, K,  $\pi$  and  $p$  (6:70:23), particle rate  $\approx$  500 MHz.




KTAG →	Kaon tagging,	CHOD →	Event multiplicity,
GTK →	Beam tracker,	LKr, MUV1, MUV2 →	Particle ID,
CHANTI →	Charged particles veto,	LAV, IRC, SAC →	Photon vetos,
STRAW →	Spectrometer,	MUV3 →	Muon vetos,
RICH →	Particle ID,	MUV0, HASC →	Off-acceptance vetos.

Note: 2018 data set divided into two subsets: S1 (20%) w/o COL and S2 (80%) w/ COL.

For details about the detector, see [NA62 collaboration, 17']

$K^+$  decay selection without PID and photon/multi-track rejection:



Pion momentum restricted to  $15 < p_{\pi^+} < 35$  GeV/c (region 1) and  $15 < p_{\pi^+} < 45$  GeV/c (region 2). The analysis of the S2 subset has been done in 5 GeV/c bins.

$K^+ \rightarrow \pi^+\pi^0$  from the control trigger chain are used for normalization.

$$\text{SES} = \frac{\mathcal{B}(K^+ \rightarrow \pi^+\pi^0) \cdot A_{\pi\pi}}{D \cdot N_{\pi\pi} \cdot A_{\pi\nu\bar{\nu}} \cdot \epsilon_{\text{RV}} \cdot \epsilon_{\text{trig}}^{\text{PNN}}} \propto \frac{1}{N_K \cdot \epsilon_{\pi\nu\bar{\nu}}}$$

The expected number of signal events is then

$$N_{\pi\nu\bar{\nu}}^{\text{exp.}} = \frac{\mathcal{B}_{\text{SM}}(K^+ \rightarrow \pi^+\nu\bar{\nu})}{\text{SES}}$$



		Subset S1 (w/o COL)	Subset S2 (w/ COL)
CTRL Trig. downscale	D	400	400
	$N_{\pi\pi} \times 10^{-7}$	3.14	11.6
Acceptance (MC)	$A_{\pi\pi} \times 10^2$	$7.62 \pm 0.77$	$11.77 \pm 1.18$
Acceptance (MC)	$A_{\pi\nu\bar{\nu}} \times 10^2$	$3.95 \pm 0.40$	$6.37 \pm 0.64$
Trig. efficiency	$\epsilon_{\text{trig}}^{\text{PNN}}$	$0.89 \pm 0.05$	$0.89 \pm 0.05$
Random veto	$\epsilon_{\text{RV}}$	$0.66 \pm 0.01$	$0.66 \pm 0.01$
	$\text{SES} \times 10^{10}$	$0.54 \pm 0.04$	$0.14 \pm 0.01$
	$N_{\pi\nu\bar{\nu}}^{\text{exp}}$	$1.56 \pm 0.10 \pm 0.19_{\text{ext}}$	$6.02 \pm 0.39 \pm 0.72_{\text{ext}}$

$\epsilon_{\text{RV}}$  encodes the random vetos caused by accidental activity in the detector.



# Background – Kaon Decays – $K^+ \rightarrow \pi^+ \pi^0$

Data driven estimation:  $N_{\text{decay}}^{\text{exp.}} = N_{\text{bkg.}} \cdot f_{\text{kin.}}(\text{region})$  where  $N_{\text{bkg.}}$  is the number of event in bkg. region after the signal selection and  $f_{\text{kin.}}$  the kinematic rejection factor for a given control/signal region.




This approach is used for the  $K^+ \rightarrow \pi^+ \pi^0$ ,  $K^+ \rightarrow \mu^+ \nu_\mu$  and  $K^+ \rightarrow \pi^+ \pi^+ \pi^-$  modes.  $K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$  and other less problematic decays rely on MC simulations.

<sup>†</sup> In the case of  $K^+ \rightarrow \pi^+ \pi^+ \pi^-$   $f_{\text{kin.}}$  is estimated from MC.


## Background – Upstream Processes

Data-driven estimation:  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  selection except for no  $\pi/K$  association and distance of closest approach (CDA)  $> 4$  mm.





This family of processes is controlled by a "box" cut in the collimator plane (subset S1) / a specialized boosted decision tree (BDT) (subset S2).

The figure shows the (expected) and observed number of background events in the **control regions**, after the signal selection. Signal regions were masked during the analysis.



Background	Expected (S1)	Expected (S2)
$\pi^+\pi^0$	$0.23 \pm 0.02$	$0.52 \pm 0.05$
$\mu^+\nu$	$0.19 \pm 0.06$	$0.45 \pm 0.06$
$\pi^+\pi^-e^+\nu$	$0.10 \pm 0.03$	$0.41 \pm 0.10$
$\pi^+\pi^+\pi^-$	$0.05 \pm 0.02$	$0.17 \pm 0.08$
$\pi^+\gamma\gamma$	$< 0.01$	$< 0.01$
$\pi^0 J^+\nu$	$< 0.001$	$< 0.001$
Upstream	$0.54^{+0.39}_{-0.21}$	$2.76^{+0.90}_{-0.70}$
Total	$1.11^{+0.40}_{-0.22}$	$4.31^{+0.91}_{-0.72}$

# Observed $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ Candidates



Categories S3 to S8 correspond to the six 5 GeV/c bins of the S2 subset.

After unblinding the 16', 17' and 20' data sets, a total of 20 candidates were found, consistent with the expectations. The corresponding branching ratio is

$$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \left( 10.6_{-3.4}^{+4.0} \Big|_{\text{stat.}} \pm 0.9_{\text{sys.}} \right) \times 10^{-11} \text{ at } 68\% \text{ CL},$$

in agreement with the SM value  $(8.4 \pm 1.0) \times 10^{-11}$ .

	Expected signal	Expected background	Observed candidates
2016	$0.267 \pm 0.20_{\text{sys.}} \pm 0.32_{\text{ext.}}$	$0.15 \pm 0.093$	1
2017	$2.16 \pm 0.13_{\text{sys.}} \pm 0.26_{\text{ext.}}$	$1.46 \pm 0.30$	2
16' + 17' + 18'	$10.01 \pm 0.42_{\text{sys.}} \pm 1.19_{\text{ext.}}$	$7.03_{-0.82}^{+1.05}$	20

Data taking will resume in July 2021. The beam line was modified during LS2 to suppress the dominant upstream background processes.

Rich complementary program: searches for dark photons, heavy neutral leptons, beam dump mode, etc.

