



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 3185 Type: **Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)**

## **(G\*) Toward understanding the nuclear efficiency threshold of bubble chamber detectors**

*Monday 6 June 2022 16:45 (15 minutes)*

A bubble chamber using fluorocarbons or liquid noble gases is a competitive technology to detect a low-energy nuclear recoil due to elastic scattering of weakly interacting massive particle (WIMP) dark matter. It consists of pressure and a temperature-controlled vessel filled with a liquid in the superheated state. Bubble nucleation from liquid to vapor phase can only occur if the energy deposition is above a certain energy threshold, described by the “heat-spike” Seitz Model. The nucleation efficiency of low-energy nuclear recoils in superheated liquids plays a crucial role in interpreting results from direct searches for WIMPs-dark matter. In this research, we used molecular dynamics simulation to study the bubble nucleation threshold, and we performed a Monte Carlo simulation using SRIM to obtain the nuclear recoil efficiency curve. The goal is to construct a physics model to explain the discrepancy observed between the experimental results and the current Seitz model. The preliminary results will be presented and compared with existing experimental data of bubble chamber detectors.

**Author:** LI, Xiang (University of Alberta)

**Presenter:** LI, Xiang (University of Alberta)

**Session Classification:** M3-8 Dark Matter Experiment II (PPD) | Experiences de matière sombre II (PPD)

**Track Classification:** Technical Sessions / Sessions techniques: Particle Physics / Physique des particules (PPD)