

# A $t\bar{t}$ Cross Section Measurement at $\sqrt{s} = 5$ TeV With the ATLAS Detector

ATLAS-CONF-2022-031

CAP 2022 - Hamilton

Pekka Sinervo (Supervisor)<sup>1</sup> **Sahibjeet Singh<sup>1</sup>**

<sup>1</sup>University of Toronto

June 6, 2022





# Outline

- 1 Motivations
- 2 Analysis Overview
- 3 Jet Energy Scale Correction
- 4 Results
- 5 Conclusions

# Motivations

- 257 pb<sup>-1</sup> of data collected by ATLAS in 2017 with  $\sqrt{s} = 5.02$  TeV
- Allows for a measurement of  $\sigma_{t\bar{t}}$  with a lower number of average interactions per crossing ( $\mu$ )
  - $\mu \sim 2$  at 5.02 TeV compared to  $\mu \sim 30$  at 13 TeV
- $\sigma_{t\bar{t}}$  measured at low  $\sqrt{s}$  affects gluon PDF at high Bjorken  $x$
- CMS measured and combined  $\sigma_{t\bar{t}}$  in the dilepton and single-lepton channels - [arXiv:2112.09114](https://arxiv.org/abs/2112.09114)
  - $\sigma_{t\bar{t}} = 62.6 \pm 4.1(\text{stat.}) \pm 3.0(\text{syst. + lumi.}) \text{ pb } (\pm 7.9\%)$




ATLAS-CONF-2022-031

# Analysis Strategy

- Measure  $\sigma_{t\bar{t}}$  in single-lepton and dilepton channels separately then combine
- Dilepton channel:
  - Cut-and-count method used to measure  $\sigma_{t\bar{t}}$  in both SF and OF dilepton events
  - CONF note released in 2021 -  
[ATLAS-CONF-2021-003](#)
- Single-lepton channel:
  - A Boosted Decision Tree (BDT) trained to separate signal and background
  - Binned profile-likelihood fit of BDT output used to extract  $\sigma_{t\bar{t}}$
- Results combined using the Convino tool
  - [arXiv:1706.01681](#)

Top Pair Branching Fractions





# Dilepton Channel

ATLAS-CONF-2022-031

$$\begin{aligned} N_{1,m}^{\ell\ell} &= L\sigma_{t\bar{t}} \epsilon_{\ell\ell} 2\epsilon_b^{\ell\ell} (1 - C_b^{\ell\ell} \epsilon_b^{\ell\ell}) f_{1,m}^{\ell\ell, t\bar{t}} + \sum_{k=\text{bkg}} s_1^k f_{1,m}^{\ell\ell,k}, \\ N_{2,m}^{\ell\ell} &= L\sigma_{t\bar{t}} \epsilon_{\ell\ell} C_b^{\ell\ell} (\epsilon_b^{\ell\ell})^2 f_{2,m}^{\ell\ell, t\bar{t}} + \sum_{k=\text{bkg}} s_2^k f_{2,m}^{\ell\ell,k} \end{aligned}$$

- Single lepton trigger, 2 OS leptons with  $p_T > 18$  GeV
- Jet  $p_T > 25$  GeV with 1 or 2 b-tags
- Cuts on  $m_{ll}$  and  $E_T^{\text{miss}}$
- Double tagging formalism used for both SF and OF events for the first time!
- Measure  $\sigma_{t\bar{t}}, \epsilon_b^{ll}, S_1^Z, S_2^Z$




# Single-lepton Channel

ATLAS-CONF-2022-031







- Exactly one electron or muon candidate
- Lepton  $p_T > 25$  GeV and  $|\eta| < 2.5$
- $\geq 2$  jets with  $p_T > 20$  GeV and  $|\eta| < 2.5$
- Events classified into 6 regions based on number of jets and b-tagged jets
- Cuts on MET and  $m_T^W$  applied to reduce mis-identified lepton background

REGION NAME	JET MULTIPLICITY	<i>b</i> -JET MULTIPLICITY
$\ell+2j \geq 1b$	2	$\geq 1$
$\ell+3j$ 1b	3	1
$\ell+3j$ 2b	3	2
$\ell+4j$ 1b	$\geq 4$	1
$\ell+4j$ 2b	4	2
$\ell+5j$ 2b	$\geq 5$	2



# BDT Distributions

- Good agreement between prediction and data in the BDT distributions for the 6 regions - [ATLAS-CONF-2022-031](#)



# 5 TeV JES Calibration

- Measure and correct the JES at  $\sqrt{s} = 5.02$  TeV using the  $Z + \text{jet}$  balance method
- Select events with a back-to-back topology and  $Z \rightarrow l^+l^-$  decay

## Topology Requirements

SFOS lepton pair with  $p_T > 20$  GeV


$81 < m_{ll} < 101$  GeV

$\geq 1$  jet with  $p_T > 10$  GeV

2nd jet  $p_T < \max(10 \text{ GeV}, 0.1 p_T^{\text{ref}})$

$$p_T^{\text{ref}} = p_T^Z |\cos \Delta\phi(Z, \text{jet})|$$

$$|\Delta\phi(Z, \text{jet})| > 2.8$$



$$p_T^{\text{ref}} = p_T(Z) * |\cos \Delta\phi(Z, j_1)|$$

- Data-MC ratio plots of the average response  $\langle r \rangle = \langle p_T^{\text{jet}} / p_T^{\text{ref}} \rangle$  used to measure shift in JES and its uncertainty

# Combination

- Convino tool used to combine Dilepton and single-lepton results - [arXiv:1706.01681](https://arxiv.org/abs/1706.01681)
- Minimize  $\chi^2 = \chi_S^2 + \chi_U^2 + \chi_P^2$ 
  - $\chi_S^2$ : The statistical uncertainty of each measurement
  - $\chi_U^2$ : The correlations between systematic uncertainties and any constraints from data
  - $\chi_P^2$ : Gaussian penalty term for systematic uncertainties and encodes prior information on correlations between systematic uncertainties

# Results

- SL:  $68.2 \pm 0.9(\text{stat.}) \pm 2.9(\text{syst.}) \pm 1.1(\text{lumi.}) \pm 0.2(\text{beam})$  pb
- DL:  $65.7 \pm 4.5(\text{stat.}) \pm 1.6(\text{syst.}) \pm 1.2(\text{lumi.}) \pm 0.2(\text{beam})$  pb
- Comb:  $67.5 \pm 0.9(\text{stat.}) \pm 2.3(\text{syst.}) \pm 1.1(\text{lumi.}) \pm 0.2(\text{beam})$  pb



ATLAS-CONF-2022-031

- Predicted  $\sigma_{t\bar{t}} = 68.2$  pb
- 4.5%  $\sigma_{t\bar{t}}$  uncertainty is the **most precise** single-lepton measurement at any energy!
- 3.9% overall uncertainty after combination

Category	Dilepton	$\delta\sigma_{t\bar{t}} [\%]$	Combination
$t\bar{t}$ generator <sup>†</sup>	1.2	1.0	0.8
$t\bar{t}$ parton-shower/hadronisation <sup>*+†</sup>	0.3	0.9	0.7
$t\bar{t}$ $h_{\text{damp}}$ and scale variations <sup>†</sup>	1.0	1.1	0.8
$t\bar{t}$ parton-distribution functions <sup>†</sup>	0.2	0.2	0.2
Single-top background	1.1	0.8	0.6
$W/Z+\text{jets}$ background*	0.8	2.4	1.8
Diboson background	0.3	0.1	< 0.1
Misidentified leptons*	0.7	0.3	0.3
Electron identification/isolation	0.8	1.2	0.8
Electron energy scale/resolution	0.1	0.1	< 0.1
Muon identification/isolation	0.6	0.2	0.3
Muon momentum scale/resolution	0.1	0.1	0.1
Lepton-trigger efficiency	0.2	0.9	0.7
Jet-energy scale/resolution	0.1	1.1	0.8
$\sqrt{s} = 5.02$ TeV JES correction	0.1	0.6	0.5
Jet-vertex tagging	< 0.1	0.2	0.2
Flavour tagging	0.1	1.1	0.8
$E_T^{\text{miss}}$	0.1	0.4	0.3
Simulation statistical uncertainty*	0.2	0.6	0.5
Data statistical uncertainty*	6.8	1.3	1.3
Total systematic uncertainty	3.1	4.2	3.7
Integrated luminosity	1.8	1.6	1.6
Beam energy	0.3	0.3	0.3
Total uncertainty	7.5	4.5	3.9

# Effect of 5.02 TeV Measurement on Gluon PDF

- Effect of combined  $\sigma_{t\bar{t}}$  at  $\sqrt{s} = 5.02$  TeV on the new ATLASpdf21 added using xFitter tool
- New data prefers harder gluon for  $x > 0.1$



ATLAS-CONF-2022-031

# Conclusions

- The most precise  $\sigma_{t\bar{t}}$  measurement in the single-lepton channel is now the  $\sqrt{s} = 5.02$  TeV measurement!
- The single  $\sigma_{t\bar{t}}$  at  $\sqrt{s} = 5.02$  TeV has a visible effect on the gluon PDF at high  $x$  when added to ATLASpdf21!
- Special runs like the  $\sqrt{s} = 5$  TeV campaign are a largely untapped pool of interesting physics in the top-sector
- Unique challenges with such runs (eg: dedicated calibrations needed) but also unique opportunities!
  - Heavy-ion program at the LHC planned for run 3!