



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 3819

Type: Oral (Non-Student) / Orale (non-étudiant(e))

## A Penrose-type inequality with angular momenta for black holes

*Thursday 22 June 2023 11:00 (15 minutes)*

The Penrose inequality places a lower bound on the mass of a black hole spacetime in terms of the area of a cross-section of the event horizon. The heuristic argument for the inequality is based upon the standard picture of gravitational collapse and it has been rigorously proved in the setting of time-symmetric initial data. We will discuss the derivation of a Penrose-type inequality with angular momenta for four dimensional, biaxially symmetric, maximal, asymptotically flat initial data sets  $(M, g, k)$  for the Einstein equations with fixed angular momenta and horizon inner boundary.

### Keyword-1

general relativity

### Keyword-2

black holes

### Keyword-3

**Author:** KUNDURI, Hari (McMaster University, Mathematics and Physics)

**Presenter:** KUNDURI, Hari (McMaster University, Mathematics and Physics)

**Session Classification:** (DTP) R2-2 Frontiers in Theoretical Physics | Frontières de la physique théorique (DPT)

**Track Classification:** Technical Sessions / Sessions techniques: Theoretical Physics / Physique théorique (DTP-DPT)